## **PCT**

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:		(11) International Publication Number:	WO 98/22494
C07K 5/00, 5/06, 237/22	A2	(43) International Publication Date:	28 May 1998 (28.05.98)
(21) International Application Number: PC	T/US97/208	56 Sheridan Road, Oakland, C	A 94618 (US). MABRY,
(22) International Filing Date: 21 November 19	997 (21.11.9	77) Thomas, E. [US/US]; 8104 Hi—	

US

US

US

US

(71) Applicants (for all designated States except US): ATHENA NEUROSCIENCES, INC. [US/US]; 800 Gateway Boulevard, South San Francisco, CA 94080 (US). ELI LILLY & COMPANY [US/US]; Lilly Corporate Center, Indianapolis,

22 November 1996 (22.11.96)

28 February 1997 (28.02.97)

28 February 1997 (28.02.97)

28 February 1997 (28.02.97)

(72) Inventors; and

(30) Priority Data:

08/755,442

08/808,528

08/807,528

08/807,427

IN 46285 (US).

(75) Inventors/Applicants (for US only): AUDIA, James, E. [US/US]; 6449 Lakeside Woods Circle, Indianapolis, IN 46278 (US). BRITTON, Thomas, C. [US/US]; 4700 Royal Oak Lane, Carmel, IN 46033 (US). DROSTE, James, J. [US/US]; 7430 Wood Stream Drive, Indianapolis, IN 46254 (US). FOLMER, Beverly, K. [US/US]; 26 Currant Drive, Newark, DE 19702 (US). HUFFMAN, George, W. [US/US]; 12 Maple Crest Drive, Carmel, IN 46032 (US). JOHN, Varghese [US/US]; 1722 18th Avenue, San

Francisco, CA 94122 (US). LATIMER, Lee, H. [US/US]; 56 Sheridan Road, Oakland, CA 94618 (US). MABRY, Thomas, E. [US/US]; 8104 Hi-Vu Drive, Indianapolis, IN 46227 (US). NISSEN, Jeffrey, S. [US/US]; 4348 Oil Creek Drive, Indianapolis, IN 46268 (US). PORTER, Warren, J. [US/US]; 8037 Lieber Road, Indianapolis, IN 46260 (US). REEL, Jon, K. [US/US]; 14701 Alsong Court, Carmel, IN 46032 (US). THORSETT, Eugene, D. [US/US]; 571 Buena Vista Street, Moss Beach, CA 94038 (US). TUNG, Jay, S. [US/US]; 2224 Semeria Avenue, Belmont, CA 94002 (US). WU, Jing [CN/US]; Apartment 502, 10 De Sabla Road, San Mateo, CA 94402 (US). EID, Clark, Norman [US/US]; 53 Dundee Drive, Cheshire, CT 06410 (US). SCOTT, William, Leonard [US/US]; 144 Buckingham Drive, Indianapolis, IN 46208 (US).

- (74) Agents: SWISS, Gerald, F. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).
- (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished upon receipt of that report.

(54) Title: METHODS AND COMPOUNDS FOR INHIBITING  $\beta$ -AMYLOID PEPTIDE RELEASE AND/OR ITS SYNTHESIS

#### (57) Abstract

Disclosed are compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. Also disclosed pharmaceutical compositions comprising a compound which inhibits  $\beta$ -amyloid peptide release and/or its synthesis as well as methods for treating Alzheimer's disease both prophylactically and therapeutically with such pharmaceutical compositions.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT .	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	Œ	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	ΙL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	is	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	Pſ	Portugal		
Cυ	. Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	Li	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# METHODS AND COMPOUNDS FOR INHIBITING $\beta$ -AMYLOID PEPTIDE RELEASE AND/OR ITS SYNTHESIS

## CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of the following U.S. Provisional Applications:

- 1. U.S. Provisional Application No. 60/\_\_\_,\_\_, which was converted pursuant to 37 C.F.R. §1.53(b)(2)(ii) from U.S. Patent Application No. 08/755,442, filed November 22, 1996;
  - 2. U.S. Provisional Application No. 60/\_\_\_\_,\_\_\_, which was converted pursuant to 37 C.F.R. §1.53(b)(2)(ii) from U.S. Patent Application No. 08/808,528, filed February 28, 1997;
- 3. U.S. Provisional Application No. 60/\_\_\_\_, which was converted pursuant to 37 C.F.R. §1.53(b)(2)(ii) from U.S. Patent Application No. 08/807,528, filed February 28, 1997; and
  - 4. U.S. Provisional Application No. 60/\_\_\_,\_\_, which was converted pursuant to 37 C.F.R. §1.53(b)(2)(ii) from U.S. Patent Application No. 08/807,427, filed February 28, 1997.

Each of these applications are incorporated herein by reference in their entirety.

#### BACKGROUND OF THE INVENTION

#### 20 Field of the Invention

15

This invention relates to methods which inhibit cellular  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. This invention also relates to pharmaceutical compositions comprising such compounds as well as methods for inhibiting release of

25  $\beta$ -amyloid peptide.

#### BACKGROUND OF THE INVENTION

## Field of the Invention

This invention relates to methods which inhibit cellular  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. This invention also relates to pharmaceutical compositions comprising such compounds as well as methods for inhibiting release of  $\beta$ -amyloid peptide.

#### References

5

15

20

The following publications, patents and patent applications are cited in this application as superscript numbers:

- Glenner, et al., "Alzheimer's Disease: Initial Report of the Purification and Characterization of a Novel Cerebrovascular Amyloid Protein", *Biochem. Biophys. Res. Commun.*, 120:885-890 (1984).
- Glenner, et al., "Polypeptide Marker for Alzheimer's Disease and its Use for Diagnosis", U.S. Patent No. 4,666,829 issued May 19, 1987.
- Selkoe, "The Molecular Pathology of Alzheimer's Disease", Neuron, 6:487-498 (1991).
- Goate, et al., "Segregation of a Missense Mutation in the Amyloid Precursor Protein Gene with Familial Alzheimer's Disease", *Nature*, 349:704-706 (1990).
- Chartier-Harlan, et al., "Early-Onset Alzheimer's Disease Caused by Mutations at Codon 717 of the β-Amyloid Precursor Proteing Gene", Nature, 353:844-846 (1989).
  - Murrell, et al., "A Mutation in the Amyloid Precursor Protein Associated with Hereditary Alzheimer's Disease", *Science*, 254:97-99 (1991).
  - Mullan, et al., "A Pathogenic Mutation for Probable Alzheimer's Disease in the APP Gene at the N-Terminus of β-Amyloid, Nature Genet., 1:345-347 (1992).

40

35

PCT/US97/20804

8 Schenk, et al., "Methods and Compositions for the Detection of Soluble β-Amyloid Peptide", International Patent Application Publication No. WO 94/10569, published 11 May 1994. 9 5 Selkoe, "Amyloid Protein and Alzheimer's Disease", Scientific American, pp. 2-8, November, 1991. 10 Losse, et al., Tetrahedron, 27:1423-1434 (1971). 11 10 Citron, et al., "Mutation of the  $\beta$ -Amyloid Precursor Protein in Familial Alzheimer's Disease Increases  $\beta$ -Protein Production, Nature, 360:672-674 (1992). 12 Hansen, et al., "Reexamination and Further Development of a Precise and Rapid Dye Method for Measuring Cell Growth/Cell 15 Kill", J. Immun. Meth., 119:203-210 (1989). 13 P. Seubert, Nature (1992) 359:325-327 14 20 Johnson-Wood et al., PNAS USA (1997) 94:1550-1555 15 Tetrahedron Letters, 34(48), 7685 (1993))

25 All of the above publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

#### State of the Art

30

35

40

Alzheimer's Disease (AD) is a degenerative brain disorder characterized clinically by progressive loss of memory, cognition, reasoning, judgment and emotional stability that gradually leads to profound mental deterioration and ultimately death. AD is a very common cause of progressive mental failure (dementia) in aged humans and is believed to represent the fourth most common medical cause of death in the United States. AD has been observed in races and ethnic groups worldwide and presents a major present and future public health problem. The disease is currently estimated to affect about two to three

million individuals in the United States alone. AD is at present incurable. No treatment that effectively prevents AD or reverses its symptoms and course is currently known.

5

10

15

The brains of individuals with AD exhibit characteristic lesions termed senile (or amyloid) plaques, amyloid angiopathy (amyloid deposits in blood vessels) and neurofibrillary tangles. Large numbers of these lesions, particularly amyloid plaques and neurofibrillary tangles, are generally found in several areas of the human brain important for memory and cognitive function in patients with AD. Smaller numbers of these lesions in a more restrictive anatomical distribution are also found in the brains of most aged humans who do not have clinical AD. Amyloid plaques and amyloid angiopathy also characterize the brains of individuals with Trisomy 21 (Down's Syndrome) and Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type (HCHWA-D). At present, a definitive diagnosis of AD usually requires observing the aforementioned lesions in the brain tissue of patients who have died with the disease or, rarely, in small biopsied samples of brain tissue taken during an invasive neurosurgical procedure.

20

The principal chemical constituent of the amyloid plaques and vascular amyloid deposits (amyloid angiopathy) characteristic of AD and the other disorders mentioned above is an approximately 4.2 kilodalton (kD) protein of about 39-43 amino acids designated the  $\beta$ -amyloid peptide ( $\beta$ AP) or sometimes A $\beta$ , A $\beta$ P or  $\beta$ /A4.  $\beta$ -Amyloid peptide was first purified and a partial amino acid sequence was provided by Glenner, et al. The isolation procedure and the sequence data for the first 28 amino acids are described in U.S. Patent No. 4,666,829<sup>2</sup>.

30

25

Molecular biological and protein chemical analyses have shown that the  $\beta$ -amyloid peptide is a small fragment of a much larger precursor protein (APP), that is normally produced by cells in many tissues of various animals,

WO 98/22494 PCT/US97/20804

-- 5 --

5

10

15

20

25

30

including humans. Knowledge of the structure of the gene encoding the APP has demonstrated that  $\beta$ -amyloid peptide arises as a peptide fragment that is cleaved from APP by protease enzyme(s). The precise biochemical mechanism by which the  $\beta$ -amyloid peptide fragment is cleaved from APP and subsequently deposited as amyloid plaques in the cerebral tissue and in the walls of the cerebral and meningeal blood vessels is currently unknown.

Several lines of evidence indicate that progressive cerebral deposition of B-amyloid peptide plays a seminal role in the pathogenesis of AD and can precede cognitive symptoms by years or decades. See, for example, Selkoe<sup>3</sup>. The most important line of evidence is the discovery that missense DNA mutations at amino acid 717 of the 770-amino acid isoform of APP can be found in affected members but not unaffected members of several families with a genetically determined (familial) form of AD (Goate, et al.4; Chartier-Harlan, et al.5; and Murrell, et al.6) and is referred to as the Swedish variant. A double mutation changing lysine595-methionine596 to asparagine595-leucine596 (with reference to the 695 isoform) found in a Swedish family was reported in 1992 (Mullan, et al.7). Genetic linkage analyses have demonstrated that these mutations, as well as certain other mutations in the APP gene, are the specific molecular cause of AD in the affected members of such families. In addition, a mutation at amino acid 693 of the 770-amino acid isoform of APP has been identified as the cause of the  $\beta$ -amyloid peptide deposition disease, HCHWA-D, and a change from alanine to glycine at amino acid 692 appears to cause a phenotype that resembles AD is some patients but HCHWA-D in others. The discovery of these and other mutations in APP in genetically based cases of AD prove that alteration of APP and subsequent deposition of its  $\beta$ -amyloid peptide fragment can cause AD.

Despite the progress which has been made in understanding the underlying mechanisms of AD and other  $\beta$ -amyloid peptide related diseases, there remains a need to develop methods and compositions for treatment of the

disease(s). Ideally, the treatment methods would advantageously be based on drugs which are capable of inhibiting  $\beta$ -amyloid peptide release and/or its synthesis in vivo.

5

#### SUMMARY OF THE INVENTION

This invention is directed to the discovery of a class of compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis and, therefore, are useful in the prevention of AD in patients susceptable to AD and/or in the treatment of patients with AD in order to inhibit further deterioration in their condition. The class of compounds having the described properties are defined by formula I below:

15

10

25

20

wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

30

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

each R<sup>3</sup> is independently selected from the group consisting of hydrogen and methyl and R<sup>3</sup> together with R<sup>4</sup> can be fused to form a cyclic structure of from 3 to 8 atoms which is optionally fused with an aryl or heteroaryl group;

5

10

20

25

30

each R<sup>4</sup> is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclic, substituted alkyl, substituted alkenyl and substituted alkynyl;

each R<sup>5</sup> is selected from hydrogen and methyl or together with R<sup>4</sup> forms a cycloalkyl group of from 3 to 6 carbon atoms;

X is selected from the group consisting of -C(O)Y and -C(S)Y where Y is selected from the group consisting of

- (a) alkyl or cycloalkyl,
- (b) substituted alkyl with the proviso that the substitution on said substituted alkyl do not include  $\alpha$ -haloalkyl,  $\alpha$ -diazoalkyl,  $\alpha$ -OC(O)alkyl, or  $\alpha$ -OC(O)aryl groups,
  - (c) alkoxy or thioalkoxy,
  - (d) substituted alkoxy or substituted thioalkoxy,
  - (e) hydroxy,
- 15 (f) aryl,
  - (g) heteroaryl,
  - (h) heterocyclic,
  - (i) -NR'R" where R' and R" are independently selected from hydrogen, alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, where one of R' or R" is hydroxy or alkoxy, and where R' and R" are joined to form a cyclic group having from 2 to 8 carbon atoms optionally containing 1 to 2 additional heteroatoms selected from oxygen, sulfur and nitrogen and optionally substituted with one or more alkyl, alkoxy or carboxylalkyl groups,
  - (j) -NHSO $_2$ -R $^8$  where R $^8$  is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic,
  - (k) -NR<sup>9</sup>NR<sup>10</sup>R<sup>10</sup> where R<sup>9</sup> is hydrogen or alkyl, and each R<sup>10</sup> is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, and
  - (1)  $-ONR^9[C(O)O]_zR^{10}$  where z is zero or one,  $R^9$  and  $R^{10}$  are as defined above;

5

10

15

20

25

30

X can also be -CR<sup>6</sup>R<sup>6</sup>Y' where each R<sup>6</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic and Y' is selected from the group consisting of hydroxyl, amino, thiol, alkoxy, substituted alkoxy, thioalkoxy, substituted thioalkoxy, -OC(O)R<sup>7</sup>, -SSR<sup>7</sup>, -SSC(O)R<sup>7</sup> where R<sup>7</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

X' is hydrogen, hydroxy, or fluoro;

X" is hydrogen, hydroxy or fluoro, or X' and X" together form an oxo group,

Z is selected from the group consisting of a bond covalently linking  $R^1$  to -CX'X"-, oxygen and sulfur;

n is an integer equal to 1 or 2; and pharmaceutically acceptable salts thereof with the provisos that:

A. when  $R^1$  is phenyl or 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH;

B. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is  $-CH(OH)CH_3$  derived from D-threonine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH or  $-C(O)OCH_3$ ;

C. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^4$  is benzyl,  $R^5$  is hydrogen, X is methoxycarbonyl, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$  is not methyl;

D. when  $R^1$  is iso-propyl,  $R^2$  is  $-CH_2C(O)NH_2$ ,  $R^3$  is hydrogen,  $R^4$  is iso-butyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

E. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^5$  is hydrogen, X is -C(O)OCH<sub>3</sub>, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$ , the nitrogen atom attached to  $R^3$ , and  $R^4$  do not form 1,2,3,4-tetrahydroiso-quinolin-2-yl or pyrrolidin-2-yl;

5

15

20

30

F. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not 4-amino-n-butyl;

G. when  $R^1$  is 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH<sub>2</sub> or -CH<sub>2</sub>OH;

H. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-CH_2OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not benzyl or ethyl;

I. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is methyl,  $R^4$  is methyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$ ;

J. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$  or -CH<sub>2</sub>OH;

K. when  $R_1$  is N-(2-pyrrolidinonyl),  $R_2$  is methyl,  $R_3$  is hydrogen,  $R_4$  is benzyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OCH<sub>3</sub>;

L. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl derived from D-alanine,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH-benzyl;

M. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is hydrogen,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CH<sub>2</sub>OH;

N. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is 4-phenylphenyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHC(CH_3)_3$ ; and

O. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHCH(CH_3)\phi$ .

Preferably, the compounds of this invention are derived from L-amino acids and, accordingly, are represented by formula IA:

5

$$R^1Z$$
 $X'$ 
 $X''$ 
 $X''$ 
 $Y''$ 
 $Y''$ 

10

15

Accordingly, in one of its method aspects, this invention is directed to a method for inhibiting  $\beta$ -amyloid peptide release and/or its synthesis in a cell which method comprises administering to such a cell an amount of a compound or a mixture of compounds of formula I above effective in inhibiting the cellular release and/or synthesis of  $\beta$ -amyloid peptide.

20

Because the *in vivo* generation of  $\beta$ -amyloid peptide is associated with the pathogenesis of AD<sup>8,9</sup>, the compounds of formula I can also be employed in conjunction with a pharmaceutical composition to prophylactically and/or therapeutically prevent and/or treat AD. Accordingly, in another of its method aspects, this invention is directed to a prophylactic method for preventing the onset of AD in a patient at risk for developing AD which method comprises administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I above.

30

25

In yet another of its method aspects, this invention is directed to a therapeutic method for treating a patient with AD in order to inhibit further deterioration in the condition of that patient which method comprises

administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I above.

5	Compounds suitable for use in the claimed methods include, by way of
	example only, the following:
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-histidine methyl ester
	N-benzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide
15	N-2-(N,N-dimethylamino) ethyl- $N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl-(S)-2-aminohexanamide$
20	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-( $S$ )-2-aminohexanamide
	N-2- $(N,N$ -dimethylamino)ethyl- $N'$ - $[N$ - $(3,5$ -difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
25	N-(4-pyridyl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
30	N-(3-pyridyl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
	N-(4-pyridyl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-( $S$ )-2-aminohexanamide
35	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate tert-butyl ester
	N-[N-(pent-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(dec-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester
40	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[3-(N,N-dimethylamino)propoxy]phenylalanine methyl ester

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[(tert-butyloxycarbonyl)methoxy]phenylalanine methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4- (carboxymethoxy)phenylalanine methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-6-(N,N-dimethylamino)hexanoate methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-pyridyl)propionate methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(3-pyridyl)propionate methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-proline methyl ester
25	1-[N-(3,5-difluorophenylacetyl)-L-alaninyl]piperidine-2-carboxylate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4-pyridyl)propionate methyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-morpholinopropionate methyl ester
35	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -4-(2-morpholinoethoxy)phenylalaninamide
	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-2-amino-3-methoxypropionamide
0	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine methyl ester
5	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-2-amino-3-(4-pyridyl)propionamide

	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-2-amino-3-(2-pyridyl)propionamide
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-yl)propionate methyl ester
	2-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1,2,3,4-tetrahydroisoquinoline-3-carboxylate methyl ester
10	N-(3-methoxybenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(1-naphthyl)propionate methyl ester
13	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-naphthyl)propionate methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-thienyl)propionate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine benzyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-bromo-propyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-iodopropyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine tert-butyl ester
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetamide
35	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetamide
40	$N$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]- $N_{\epsilon}$ -(tent-butoxycarbonyl)-L-lysine methyl ester
40	methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-4-phenylbutanoate
45	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 2-phenylethyl ester
<del>4</del> J	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 3-phenylpropyl ester

	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetamide
-	N-[N-(phenylacetyl)-L-alaninyl]-L-threonine methyl ester
5	N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide
	N'-[N-(phenylacetyl)-L-alaninyl]-L-alaninamide
10	N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide
	N'-[ $N$ -(phenylacetyl)-L-alaninyl)-L-valinamide
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetate ethyl ester
	N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide
20	N,N-dimethyl- $N'$ - $[N$ -(phenylacetyl)-L-alaninyl]-L-phenylalaninamide
20	N,N-dimethyl- $N'$ -[ $N$ -(phenylacetyl)- $L$ -alaninyl]- $L$ -leucinamide
	N,N-dimethyl- $N'$ -[ $N$ -(phenylacetyl)-L-alaninyl]-L-valinamide
25	N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide
	N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide
30	N-methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-( $S$ )-2-aminohexanamide
	N,N-dimethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-(S)-2-aminohexanamide
35	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-methoxyphenyl) acetate methyl ester
40	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-methoxyphenyl) acetate methyl ester
45	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetate ethyl ester

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetate ethyl ester
5	N-[N-(cyclohexylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
5	N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(cyclohex-1-enylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1-aminocyclopropane-1-carboxylate methyl ester
15	N-2-(N,N-dimethylamino) ethyl- $N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide$
15	N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine benzyl ester
20	N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine ethyl ester
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
25	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-alanine ethyl ester
23	N-[N-(3-nitrophenylacetyl)-L-alaninyl]glycine ethyl ester
	N-hydroxy-N'-[N-(3-nitrophenylacetyl)-L-alaninyl]-D,L-threoninamide
30	N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-2-amino-3-(3-hydroxyphenyl)propionate methyl ester
35	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-tyrosine ethyl ester
	N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester
40	N-[N-[N-(isovaleryl)-L-valinyl]-L-phenylglycinyl]-L-alanine iso-butyl ester
	N-[N-(isovaleryl)-L-phenylalaninyl]-L-alanine iso-butyl ester
45	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester

	1-[N-(3-nitrophenylacetyl)-L-alaninyl]-indoline-(S)-2-carboxylate ethyl ester
5	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide
3	N-methoxy-N-methyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide
	N-iso-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
10	N,N-di- $n$ -propyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide
	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valinamide
15	N-(4-nitrophenyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N'-[N-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninyl]-L-phenylalaninamide
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide
25	N-iso-butyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide
	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
30	N-(4-nitrobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N-(4-nitrophenyl)- $N'$ -[ $N$ -(isovaleryl)- $L$ -phenylglycinyl]- $L$ -alaninyl]- $L$ -alaninamide
35	N-(4-nitrophenyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
10	N-benzyl- $N$ -methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
<b>.</b>	N-(3,5-difluorobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
15	N-(3-nitrobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide

	N-benzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
5	N-(4-nitrobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophan methyl ester
10	N-(4-methoxybenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
10	N-[N-(phenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninyl]-L-phenylglycine methyl ester
15	N-[N-(cyclohexylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine methyl ester
20	N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]-L-phenylglycine methyl ester
0.5	N-(2-phenylethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
25	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophanamide
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-cyclohexylpropionate methyl ester
30	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-(S)-2-amino-3-(4-nitrophenyl)propionamide
25	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-serine ethyl ester
35	$N$ -[(R)- $\alpha$ -methylbenzyl]- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide
40	$N$ -[(S)- $\alpha$ -methylbenzyl]- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide
	N-(4-fluorobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
45	N-(4-pyridylmethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide

	N-(4-trifluoromethylbenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-phenylpropionate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-methylpropionate methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-cyclohexylacetate ethyl ester
13	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
20	N-[N-(isovaleryl)-2-amino-2-cyclohexylacetyl]-L-alanine ethyl ester
20	N-2-(N,N-dimethylamino) ethyl- $N'-[N-(3,5-difluorophenylacetyl)-L-$ alaninyl]-L-phenylglycinamide
25	N-(2-pyridylmethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N-[N-(3-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
30	N-[N-(2-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
30	N-[N-(4-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
35	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-fluorophenyl)acetate ethyl ester
33	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-fluorophenyl)acetate ethyl ester
40	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
70	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-phthalimidopropionate ethyl ester
45	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine neopentyl ester

	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine <i>tert</i> -butyl ester
•	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
10	4-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valinyl]morpholine
10	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine ethyl ester
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threonine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester
20	4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-amino-3-tert-butoxybutyryl]morpholine
20	4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucinyl]morpholine
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine methyl ester
25	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine
	N-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threoninyl]-L-valine ethyl ester
30	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-leucine methyl ester
35	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine methyl ester
	N-2-methoxyethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
40	N-2-(N,N-dimethylamino) ethyl- $N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide$
	N-cyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
45	N-neopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide

N-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 alaninamide 3-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]thiazolidine N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl 10 ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester N-(R)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-15 alaninamide 1-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]pyrrolidine N-(S)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 alaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valine methyl ester N-2-fluoroethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-25 alaninamide N-[(S)-6-methyl-3-oxohept-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-4-nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-30 aminobutyramide N-4-nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminopentanamide 35 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3fluorophenyl)acetate methyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-40 thienyl)acetamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(5chlorobenzothiophen-2-yl)acetate methyl ester 45 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-2yl)acetate ethyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-3yl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-thienyl)acetate 5 methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-5yl)acetate ethyl ester 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2thienyl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2thienyl)acetate tert-butyl ester 15 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2thienyl)acetic acid N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(1H-tetrazol-5-20 yl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(6-methoxy-2naphthyl)acetate methyl ester 25 N-IN-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3trifluoromethylphenyl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4,5,6,7tetrahydrobenzothiophen-2-yl)acetate methyl ester 30 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(thieno[2,3blthiophen-2-yl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-methylthiazol-35 4-yl)acetate methyl ester (3S,4S)-N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-4-amino-3-hydroxy-5-phenylpentanoate methyl ester 40 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohex-4-enoate methyl ester N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester 45 N-tert-butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(4-phenylphenyl)acetamide

	N-[N-(3,5-difluorophenylacetyl)-(S)-2-aminobutanoyl]-L-phenylglycine ten-Butyl Ester
5	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine tert-butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycine methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine methyl ester
·	N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-leucinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylalaninyl]-L-phenylglycine methyl ester
20	N-[N-(3,5-difluorophenylacetyl)glycinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine methyl ester
25	N-[N-(phenylacetyl)-L-alaninyl]-L-alanine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-leucine methyl ester
30	N-[N-(phenylacetyl)-L-alaninyl]-L-isoleucine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-proline methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
35	N-[ $N$ -(phenylacetyl)-L-alaninyl]- $N$ -( $tert$ -butoxycarbonyl)-L-lysine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-glycine methyl ester
10	N-[N-(phenylacetyl)-L-alaninyl]-L-valine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester
<b>.</b> 5	N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester
J	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine

	N-[N-(phenylacetyl)-L-alaninyl]-L-N-methylalanine methyl ester
	N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester
5	N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester
· .	N-Cyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-hydroxyproline ethylester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-lysine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-glutamide
	1-[N-(3,5-difluorophenylacetyl)-L-alaninyl]piperidine-2-carboxylate methyl ester
20	N-[(S)-3-hydroxy-6-methylhept-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
25	N-[(S)-2-hydroxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
23	$N$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ -fluoroacetyl)-L-alaniny]-L-phenylglycine <i>tert</i> -butyl ester
30	N-[N-(3,5-difluorophenylacetyl)-2-(S)-aminocyclohexylacetyl]-L-phenylglycine methyl ester
	N-[(1R,2S)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
35	N-[(1R,2S)-1-hydroxy-1,2-diphenyleth-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(1S,2R)-1-hydroxy-1-phenylprop-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
40	N-2-methoxyethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-glycinamide
<b>4</b> 5	$N$ -[(S)- $\alpha$ -hydroxy- $\alpha$ -phenyl- $iso$ -propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

	N-[(S)-2-hydroxy-1,2-diphenylethyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
5	N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	$N$ -[ $\alpha$ -hydroxy- $\alpha'$ -(4-hydroxyphenyl)-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
10	N-2-pyridylmethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
	$N$ -[ $\alpha$ -hydroxy- $\alpha'$ -pyrid-2-yl- $iso$ -propyl]-N'-(3,5-difluorophenylacetyl)-L alaninamide
15	$N$ -[ $\alpha$ -hydroxy- $\alpha'$ -pyrid-4-yl- $iso$ -propyl]-N'-(3,5-difluorophenylacetyl)-L alaninamide
	N-[(S)-1-hydroxy-4-methylpent-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
20	$N$ -[ $\alpha$ -methoxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[1-hydroxy-3-methyl-but-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(6-aminopyrid-2-yl)acetate methyl ester
30	N-[1-hydroxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
30	N-[(S)-2-methoxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
35	N-[(S)-1-methoxy-2-phenyl-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(S)-1-acetoxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
40	N-[(S)-1-( $tert$ -butylcarbonyloxy)-hex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[2-hydroxy-1-(thien-2-yl)ethyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
45	N-[(S)-2-hydroxy-2-methyl-1-phenylprop-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

	N-[N-(3,5-difluorophenylacetyl)-L-(thien-2-yl)glycinyl]-L-phenylalanine tert-butyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinol
	N-[N-(cyclopropaneacetyl)-L-phenylglycinyl]-L-phenylglycinol
	N-[N-(cyclopentaneacetyl)-L-phenylglycinyl]-L-phenylglycinol
10	N-[N-(3,5-difluorophenylacetyl)-D,L-phenylglycinyl]-D,L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-D,L-valinyl]-D,L-phenylglycinamide
15	N-[N-(2-thienylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-[N-(n-caprotyl)-L-alaninyl]-L-phenylglycinamide
20	N-[ $N$ -(3,5-difluorophenylacetyl)-L-norleucinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-norvalinyl]-L-phenylglycine methyl ester
25	$N$ -[ $N$ -(3,5-difluorophenylacetyl)- $\dot{\mathbf{L}}$ -tert-leucinyl]- $\mathbf{L}$ -phenylglycine methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-isoleucinyl]-L-phenylglycine methyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-cyclohexylalaninyl]-L-phenylglycine methyl ester
35	N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(cyclopropyl)acetyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-3-yl)acetyl]-L-phenylglycine methyl ester
40	N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-2-yl)acetyl]-L-phenylglycine methyl ester
45	N-[N-(3,5-difluorophenylacetyl)-L-(4-fluorophenyl)glycinyl]-L-phenylglycine methyl ester

	N-[N-(3,5-difluorophenylacetyl)-D-(4-fluorophenyl)glycinyl]-L-phenylglycine methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-(4-methoxyphenyl)glycinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester
10	N-[N-(cyclopropylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester
15	N-[N-(cyclopentylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester
15	N-[N-(tert-butylacetyl)-L-alaninyl]-L-phenylglycinamide
20	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -(5-bromothien-2-yl)glycinamide
20	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(5-bromothien-2-yl)glycinamide
25	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-bromothien-2-yl)glycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-2-yl)glycinamide
30	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ -(thien-2-yl)glycinamide
35	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-3-yl)glycinamide
33	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ -(thien-2-yl)glycinamide
40	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
45	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(5-chlorothien-2-yl)glycinamide

	N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-4-(phenyl)phenylglycinamide
5	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenoxy)phenylglycinamide
	$N$ -(S)-(-)- $\alpha$ -methylbenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
10	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenyl)phenylglycinamide
15	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(ethyl)phenylglycinamide
13	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-(phenyl)phenylglycinamide
20	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -2-(benzyl)phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -4-bromophenylglycinamide
25	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -4-(cyclohexyl)phenylglycinamide
30	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -4-(4-ethylphenyl)phenylglycinamide
30	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-(tert-butyl)phenylglycinamide
35	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -3-(4-chlorophenoxy)phenylglycinamide
	N-cyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(phenyl)phenylglycinamide
40	$N$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ -hydroxyacetyl)-L-alaninyl]-L-phenylglycine <i>tert</i> -butyl ester
45	$N$ -tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetyl)-L-alaninyl]-L-phenylglycinamide

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine tert-butyl ester
5	N-[(S)-1-oxo-1-phenylprop-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(pyrid-3-yl)glycine tert butyl ester
10	[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]morpholine
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(2-methoxy)phenylglycine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-butoxycarbonyl(hydroxyl amine) ester
20	N-neopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-tetrahydrofurfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
25	N-methoxy-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
30	[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]azetidine
30	N-iso-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
35	N-cyclopropanemethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-methoxy- $N$ -methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
40	N-2-methylprop-2-enyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L phenylglycinamide
45	N-(pyrid-3-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide

	N-(pyrid-4-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
5	N-furfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-cyclopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
10	N-1-benzylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
15	N,N-dimethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
15	N-2,2,6,6-tetramethylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
20	N-2-methylcyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-4-methylcyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
25	N-1-ethoxycarbonylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
20	N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
30	N-tert-butoxy- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
35	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-butyl(hydroxylamine) ester
,	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine hydrazide
40	N-(1-ethoxyethen-1-yl)-[ $N'$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine hydrazide
	N-[N-(phenylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
45	N-4-(phenyl)butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide

	N-3-(4-iodophenoxy)propyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
5	N-6-(amino)hexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide Hydrochloride
	N-1-(phthalimido)pent-2-yl-N'-(3,5-difluorophenylacetyl)-L-alaninamide
10	N-[N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenyl)glycinyl]-L-(3,5-difluorophenyl)glycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-norleucine
15	N-[N-(cyclopentaneacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
*3	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine isopropyl ester
20	N-(isopropyl) $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester
25	N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester
2.5	N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine iso-butyl ester
30	N-[ $N$ -(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycine methyl ester
	$N$ -[ $N$ -(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(3- $\alpha$ -phenyl)proline methyl ester
35	N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-azetidine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(5-chlorobenzothiophen-2-yl)acetate methyl ester
40	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-yl)propionate <i>tert</i> -butyl ester
45	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide tert-butyl ester
45	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide

	N-[N-(3,4-dichlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
5	N-[N-(3-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3-bromophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
10	N-[N-(4-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide
15	N-[N-(4-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide
15	N-[N-(3-trifluoromethylphenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3-methoxyphenylacetyl)-L-alaninyl]-D-phenylglycinamide
20	N-[N-(2-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(1-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide
25	N-[N-(2-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide
23	N-[N-(phenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-furanyl)acetamide
35	N'-[ $N$ -(3,5-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide
	N'-[ $N$ -(3,4-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide
40	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanin-N-methylsulfonamide
	N''-methyl- $N''$ -phenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-glycinamide
45	N''-methyl- $N''$ -phenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide

	N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycinamide
5	N''-methyl- $N''$ -benzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-glycinamide
3	N''-4-fluorobenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
10	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-fluoro)phenylglycine neopentyl ester
	N-[N-(2,3,4,5,6-pentafluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine tert-butyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)serinyl]-L-phenylglycine methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)threoninyl]-L-phenylglycine methyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-threoninyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-serinyl]-L-phenylglycine methyl ester
30	N''-4-methylphenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	N''-tetrahydrofurfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
35	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenyl-glycinamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)- $L$ -methionyl]- $L$ -phenylglycinamide
10	N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycinamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinamide
45	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycinamide

	$N$ -[(R)- $\alpha$ -methylbenzyl]- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
5	N-[1-phenyl-2-oxo-3-methylbutan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-propan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
10	N-[1-phenyl-2-oxo-pentan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
15	N-[1-phenyl-2-oxo-2-phenyl-ethan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-Lalaninamide
	N-[1-phenyl-2-oxo-butan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)- $L$ -alaninamide
20	N-[1-phenyl-2-oxo-4-methylpentan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide
	$N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L- $\alpha$ -hydroxyphenylalanine methyl ester
25	N''-[4-((2-hydroxy-4-azido)-phenyl)-NHC(O)-)butyl] $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
20	N-[(S)-1-phenyl-2-oxo-2-phenyl-ethan-1-yl]-N'-(3,5-difluorophenyl-acetyl)-L-alaninamide
30	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine tert-butyl ester
35	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-phenylphenylglycine <i>tert</i> -butyl ester
	[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(2,3-benzo[b]proline) methyl ester
10	N''-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4- $n$ -butylphenylglycinamide
_	N''-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-(phenylacetenyl)phenylglycinamide
15	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide

	N-[1,3-diphenyl-2-oxo-propan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)- $L$ -alaninamide
<b>. 5</b>	N-[1-phenyl-2-oxo-2-cyclopentylethan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-hexan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
10	N-[1-phenyl-2-oxo-3-methylpentan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
15	N''- $n$ -hexyl-6-biotinamidyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinthioamide
13	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-methioninyl]-L-methionine
20	N'-[N-(2-tert-BOC-amino)propionyl)-L-alaninyl]-L-phenylglycine methyl ester
20	N''-tert-butyl $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-fluorophenylglycinamide
25	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-2-phenylglycine methyl ester
	N-[(S)-1-phenyl-2-oxo-3-phenylpropan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
30	N'-[ $N$ -(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-phenylglycine
35	N'-[ $N$ -(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-phenylglycine $tert$ -butyl ester
33	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine
40	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine tert-butyl ester
	N-[2-hydroxy-1-(S)phenyleth-1-yl]- $N'$ -[(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-alaninamide
45	$N-[2-l_1ydroxyeth-1-yl]-N'-[(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide$

N'-[N-(3,5-difluorophenyl-2-oxo-acetyl)-L-alaninyl]-L-2-phenylglycine tert-butyl ester

[N-(2,5-dichlorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3,5-difluorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3,4-dichlorothiophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3-aminoproprionyl)-L-alaninyl]-L-phenylglycine tert-butyl ester

[N-(3-tert-butoxycarbonylamino)propionyl)-L-alaninyl]-L-phenylglycine

15

tert-butyl ester

10

5

The pharmaceutical compositions described above comprise a pharmaceutically inert carrier and a compound of the formula I above.

20

In formula I above, X'' is preferably hydrogen and X' is preferably hydrogen or fluoro.

In formula I above, Z is preferably a covalent bond linking  $R^1$  to -CX'X''-.

In formula I above, preferred R<sup>1</sup> unsubstituted aryl groups include, for example, phenyl, 1-naphthyl, 2-naphthyl, and the like.

Preferred R<sup>1</sup> substituted aryl groups include, for example, monosubstituted phenyls (preferably 3 or 5 substituents); disubstituted phenyls (preferably 3,5 substituents); and trisubstituted phenyls (preferably 3,4,5 substituents). Preferably, the substituted phenyl groups do not include more than 3 substituents.

Examples of substituted phenyls include, for instance, 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 3-methoxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2-hydroxyphenyl, 2-methylphenyl, 2-fluorophenyl, 2-chlorophenyl, 3,4-difluorophenyl, 2,3,4,5,6-pentafluorophenyl, 3,4-dibromophenyl, 3,4-dichlorophenyl, 3,4-methylene-dioxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 2,4-dichlorophenyl, and 2,5-difluorophenyl.

Preferred R<sup>1</sup> alkaryl groups include, by way of example, benzyl, 2-phenylethyl, 3-phenyl-*n*-propyl, and the like.

Preferred R¹ alkyl, substituted alkyl, alkenyl, cycloalkyl and cycloalkenyl groups include, by way of example, *iso*-propyl, *n*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, aminomethyl, N-*tert*-butoxycarbonylaminomethyl, and the like.

20

5

10

15

Preferred R<sup>1</sup> heteroaryls and substituted heteroaryls include, by way of example, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, fluoropyridyls (including 5-fluoropyrid-3-yl), chloropyridyls (including 5-chloropyrid-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-(thiophenyl)thiophen-5-yl, 6-methoxythionaphthen-2-yl, 3-phenyl-1,2,4-thiooxadiazol-5-yl, 2-phenyloxazol-4-yl, and the like.

30

25

Preferably  $R^2$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic. Particularly preferred  $R^2$  substituents include, by way of example, methyl, ethyl, n-propyl,

WO 98/22494 PCT/US97/20804

iso-propyl, n-butyl, iso-butyl, sec-butyl, phenyl, 4-fluorophenyl, 3,5-difluorophenyl, 4-methoxyphenyl, benzyl, cyclopropyl, cyclohexyl, cyclopentyl, cycloheptyl, thien-2-yl, thien-3-yl,  $-CH_2CH_2SCH_3$ ,  $-CH_2OCH_2\phi$ ,  $-CH(CH_3)OCH_2\phi$ ,  $-CH(OH)CH_3$ ,  $-CH_2OH$  and the like. As noted below,  $R^2$  (as well as  $R^4$ ) is preferably the side chain of an L-amino acid.

Preferably, R<sup>3</sup> is hydrogen, methyl or together with R<sup>4</sup> and the nitrogen to which R<sup>3</sup> is attached forms pyrrolidin-2-yl, 2,3-dihydroindol-2-yl, piperidin-2-yl, 4-hydroxy-pyrrolidin-2-yl, 1,2,3,4-tetrahydroisoquinolin-3-yl, and the like.

10

5

Preferred R<sup>4</sup> substituents include, for example, hydrogen, methyl, ethyl, iso-propyl, n-propyl, n-butyl, sec-butyl, iso-butyl, cyclopentyl, cyclohexyl, allyl, iso-but-2-enyl, 3-methylpentyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-indol-3-yl, phenyl, p-(phenyl)phenyl, m-(phenyl)phenyl o-fluorophenyl, m-fluorophenyl, p-fluorophenyl, p-bromophenyl, m-methoxyphenyl, 15 p-methoxyphenyl, phenethyl, benzyl, m-hydroxybenzyl, p-hydroxybenzyl, p-nitrobenzyl, m-trifluoromethylphenyl, p-(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, p-(CH<sub>3</sub>)<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, p-phenylphenyl, 3,5-difluorophenyl, p-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyrid-6-yl, 4-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)benzyl, -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>, -CH<sub>2</sub>-imidazol-4-yl, -CH<sub>2</sub>-(3-tetrahydrofuranyl), 20 - $CH_2$ -thien-2-yl, - $CH_2$ -thiazol-4-yl, - $CH_2$ (1-methyl)cyclopropyl, - $CH_2$ -thien-3-yl, thien-3-yl, thien-2-yl,  $-CH_2-C(O)O-t$ -butyl,  $-CH_2-C(CH_3)_3$ ,  $-CH_2CH(CH_2CH_3)_2$ , 2-methylcyclopentyl, -cyclohex-2-enyl, -CH[CH(CH<sub>3</sub>)<sub>2</sub>]COOCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>SCH<sub>3</sub>,  $-CH_2CH_2N(CH_3)_2$ ,  $-CH_2C(CH_3)=CH_2$ ,  $-CH_2CH=CHCH_3$  (cis and trans), -CH<sub>2</sub>OH, -CH(OH)CH<sub>3</sub>, -CH(O-t-butyl)CH<sub>3</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>NH-Boc, 25 -(CH<sub>2</sub>)<sub>4</sub>NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>4</sub>N(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-pyridyl (e.g., 2-pyridyl, 3-pyridyl and 4-pyridyl), pyridyl (2-pyridyl, 3-pyridyl and 4-pyridyl), -CH2-naphthyl (e.g., 1-naphthyl and 2-naphthyl), -CH<sub>2</sub>-(N-morpholino), p-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5chlorobenzo[b]thiophen-2-yl, 4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl, 30

benzo[b]thiophen-3-yl, tetrazol-5-yl, 5-chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl, -CH<sub>2</sub>-N-phthalimidyl, 2-methylthiazol-4-yl, and thieno[2,3-b]thiophen-2-yl, 5-bromothien-2-yl, 4-bromothien-2-yl, 5-chlorothien-2-yl, 3-phenoxyphenyl, 2-phenoxyphenyl, 4-ethylphenyl, 2-benzylphenyl, (4-ethylphenyl)phenyl, 4-tert-butylphenyl, 4-n-butylphenyl, o-(4-chlorophenoxy)phenyl, furan-2-yl, 4-phenylacetylenylphenyl and the like.

Preferably, R<sup>5</sup> is hydrogen. However, in another embodiment, R<sup>4</sup> and R<sup>5</sup> are fused to form a cycloalkyl group including, for example, cyclopropyl, cyclobutyl, and the like.

One preferred X substituent is -C(O)Y. Preferably Y is hydroxy, alkoxy or substituted alkoxy such as methoxy, ethoxy, n-propoxy, iso-propoxy, 15 n-butoxy, iso-butoxy, tert-butoxy, neo-pentoxy, benzyloxy, 2-phenylethoxy, 3phenyl-n-propoxy, 3-iodo-n-propoxy, 4-bromo-n-butoxy, -ONHC(O)OC(CH<sub>3</sub>)<sub>3</sub>, -ONHC(CH<sub>3</sub>)<sub>3</sub> and the like. Another preferred Y group is -NR'R" where R' and R" are as defined above. Such preferred Y groups include, by way of example, amino (-NH<sub>2</sub>), -NH(iso-butyl), -NH(sec-butyl), N-methylamino, N,Ndimethylamino, N-benzylamino, N-morpholino, azetidino, N-thiomorpholino, 20 N-piperidinyl, N-hexamethyleneimino, N-heptamethylene-imino, N-pyrrolidinyl, -NH-methallyl, -NHCH<sub>2</sub>-(furan-2-yl), -NHCH<sub>2</sub>-cyclopropyl, -NH(tert-butyl), -NH(p-methylphenyl), -NHOCH<sub>3</sub>, -NHCH<sub>2</sub>(p-fluorophenyl), -NHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, -NH-cyclopentyl, -NH-cyclohexyl, -NHCH2CH2N(CH3)2, -NHCH2C(CH3)3, -NHCH<sub>2</sub>-(pyrid-2-yl), -NHCH<sub>2</sub>-(pyrid-3-yl), -NHCH<sub>2</sub>-(pyrid-4-yl), 25 N-thiazolindinyl, -N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -N[CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -NHOH,  $-NH(p-NO_2-\phi)$ ,  $-NHCH_2(p-NO_2-\phi)$ ,  $-NHCH_2(m-NO_2-\phi)$ ,  $-N(CH_3)OCH_3$ , -N(CH<sub>3</sub>)CH<sub>2</sub>- $\phi$ , -NHCH<sub>2</sub>-(3,5-di-fluorophenyl), -NHCH<sub>2</sub>CH<sub>2</sub>F, -NHCH<sub>2</sub>(p- $CH_3O-\phi)$ ,  $-NHCH_2(m-CH_3O-\phi)$ ,  $-NHCH_2(p-CF_3-\phi)$ ,  $-N(CH_3)CH_2CH_2OCH_3$ , -NHCH<sub>2</sub>CH<sub>2</sub> $\phi$ , -NHCH(CH<sub>3</sub>) $\phi$ , -NHCH<sub>2</sub>-(p-F- $\phi$ ), -N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, 30

-NHCH<sub>2</sub>-(tetrahydrofuran-2-yl), -NHCH<sub>2</sub>(p-trifluoromethylphenyl),

(biotinamidyl), and the like.

5

- -NHCH<sub>2</sub>C(CH<sub>3</sub>) = CH<sub>2</sub>, -NH-[(p-benzyl)pyrid-4-yl], -NH-[(2,6-dimethyl)pyrid-4-yl], -NH-(2-methylcyclohexyl), -NH-(4-methylcyclohexyl), -NH-[N-ethoxycarbonyl]-piperidin-4-yl, -NHOC(CH<sub>3</sub>)<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- $\phi$ , -C(O)NH(CH<sub>2</sub>)<sub>3</sub>O-(p-CH<sub>3</sub>) $\phi$ , -C(O)NH(CH<sub>2</sub>)<sub>6</sub>NH<sub>2</sub>, -NH-(tetrahydrofuran-2-yl), -N(CH<sub>3</sub>) $\phi$ , -NH(CH<sub>2</sub>)<sub>4</sub>NHC(O)-(2-hydroxy-4-azido)-phenyl, -NH(CH<sub>2</sub>)<sub>6</sub>-
- Another preferred Y group is an alkyl group such as methyl, ethyl, iso-propyl, n-propyl, iso-butyl, n-butyl, sec-butyl, tert-butyl,

  -CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-pyridy-2-yl, -CH<sub>2</sub>-pyridy-3-yl, -CH<sub>2</sub>-pyridy-4-yl,
  -CH<sub>2</sub>-fur-2-yl, and the like; a substituted alkyl group such as benzyl; a cycloalkyl group such as cyclopentyl; and an aryl group such as phenyl.
- Still another preferred Y group is -NHSO<sub>2</sub>-R where R is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic. Such groups are exemplified by NH-SO<sub>2</sub>-CH<sub>3</sub>.
- Preferred Y' groups include a substituted alkyl group such as
  -CH<sub>2</sub>OH, -CH(OH)CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(OH)φ, -CH(OH)CH<sub>2</sub>C(O)OCH<sub>3</sub>,
  -C(OH)(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>OC(O)OCH<sub>3</sub>, -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>, and the like.

Preferred compounds for use in the methods of this invention include those set forth in the tables below:

ξg	×
o==	Z— &
I	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~
	~ ~

	7	_	1	T	_	<del>-</del>	_	_	_	<del></del>	-		<del>-</del>	_	<del></del>	_	<del></del> -	_
×		-C(0)NII,	TIN(O)	(0)2-	-C(O)NII,	110/00/0/2	(いつつつ(ここう)	ווייטוטיטי	(1120/0/2	ייייייייייייייייייייייייייייייייייייייי	(1)26(1)	-CONNITCIL-A	A (115.11.1/6)	-C(O)NIICII'CII'NCII')	7/5	-C(0)NHCH,CH,OCH,		-C(O)NHCH,CH,N(CH,)
₩		=	=		-	11	:	=======================================		=		=		=			:	=
*≃		φ	\$-	*	À	•		-CII,CII,CII,		-CII3-imidazol-4-yl		-CII,CII,CII,CII,		-(CII <sub>2</sub> ),CII <sub>3</sub>		الکرردالی)۔	7 110	
K3	11	=	=			I		Port Port		Post -		=		11		=	Ξ	•
R	IIJ	(II)	-CII)	-CII,		ĊII,		-CII,		-CII,		-CII,	110	(11)	,CII.		-Cill	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
R	-Purivi	l (non )	thien-2-yl	n-butyl		cyclopentyl		-φ-,1-In-C'C		-\$1-ID-C'F		-φI-ID-C'F	3 S. (15. 15. 16.	A 1-10-0'0	3.5-di-F-o-		3,5-di-F-φ-	

R¹	K <sup>2</sup>	R³	R*	1 R3	X
3,5-di-ŀ-ф-	-CII,	=	-CII <sub>2</sub> -ф	н	-C(0)NIICII <sub>2</sub> -(pyrid-4-yl)
3,5-di-F-ф-	·CII,	=	-CII <sub>1</sub> -ф	==	-C(O)NIICII <sub>1</sub> -(pyrid-3-y1)
3,5-di-F-φ-	-CII,	11	-(CH <sub>J</sub> ),CH <sub>J</sub>	=	-C(0)NIICII <sub>2</sub> -pyrid-4-yl
3,5-di-F-φ-	-CII,	-	-CII,CII,CII,	1	-C(0)0C(CiI,),
CII,=CIICII1,-	-CII,	11	-CII <sub>2</sub> φ	=	-C(0)0CII,
CH,(CII,),CII=CIICII,-	-CII,	l-vil dens	-CII <sub>2</sub> $\phi$	1	-C(0)0CII,
3, 5-di-F-φ-	·CII]	==	p-(CII <sub>3</sub> ) <sub>2</sub> NCII <sub>2</sub> CII <sub>2</sub> CII <sub>2</sub> O- benzyl-	Ξ	-C(0)0CH,
3, 5-di-F-φ.	·CII,	11	p-(CH <sub>3</sub> ),COC(O)CH <sub>2</sub> O- benzyl-	=	-C(0)0CH,
3,5-di-F-φ-	-CII,	Ξ	p-hydroxybenzyl-	=	-C(0)0CII,
3, 5-di-F- <i>φ</i> -	-CII,	11	p-IIOOCCII <sub>2</sub> O-benzyl	=	-C(0)0CII,
3, 5-di-F∙¢-	-CII,	=	p-(N-morpholino-CII <sub>2</sub> CII <sub>2</sub> O-benzyl-	=	-C(0)0C11,
3,5-di-F-4-	-CII,	=	-(CII <sub>2</sub> ) <sub>4</sub> -N(CII <sub>3</sub> ) <sub>2</sub>	=	-C(0)0CII,
3,5-di-F-ф-	-CII,	11	-CII <sub>2</sub> -(pyrid-2-yl)	Ξ	-C(0)0CII,
3,5-di-F-φ-	-CII,	11	-CII <sub>2</sub> -(pyrid-3-yl)	Н	-C(0)0CH,
3,5-di-F-ф-	-CII,	$R_{\rm J}/R_{\rm J}$ and N = L-pyrrolidinyl		II	-С(0)ОСН,
					Å

RI	R²	R³	R	- IR3	X
φ.	-CIII,	$R_{\rm J}/R_{\rm 4}$ and N = piperidin-2-y!		=	-C(0)0CII,
3,5-di-F-ф-	-CII,	11	-CII <sub>2</sub> -(pyrid-4-yl)	=	-C(0)0CII,
3,5-Ji-F-ф-	-CII,	1	-CII,OCII,	=	-C(0)0CII,
3,5-di-F-ф-	-CII,	-	-CII <sub>2</sub> -(N-morpholino)	=	-C(0)0CII,
3,5-di-I <sup>2</sup> -φ-	-CII,	=	p-(N-morpholino-CII <sub>2</sub> CII <sub>1</sub> -O)-benzyl-		-C(O)NIICII,CII,OCII,
3, 5-di-F-φ-	-CII,	=	·CH,0CH,	H	-C(0)NIICHICHIOCH
3,5-di-F-ф-	-CII,	=	H	11	-C(0)0CH,
3,5-di-F-φ-	-CII,	11	-CII <sub>2</sub> -pyrid-4-yl	Ξ	-C(0)NIICII,CII,OCII,
3,5-di-F-4-	-CII,	П	-CII <sub>2</sub> -pyrid-2-yl	H	-C(0)NIICH,CH,OCH,
3,5-di-F-ф-	-CII,	-	-CH <sub>2</sub> -thiazol-4-yl	=	-C(0)0CII,
3,5-di-F-ф-	·CII)	$R_3/R_4$ and $N = 1,2,3,4$ -tetrahydro-isoquinolin-3-yl		=	-C(0)0CII,
3,5-di-F-φ-	·CII,	11	-CII <sub>2</sub> -φ	11	-C(0)NIICII <sub>1</sub> -(m-CII <sub>3</sub> 0-¢)
3,5-di-P-ф-	-CII,	11	-CII <sub>2</sub> -1-naphthyt	=	-C(0)0CII,
3,5-di-F-φ-	-CII,	==	-CII <sub>2</sub> -2-naphthyi	H	-C(0)0CII,
3,5-di-F-ф-	-CII,	II	-CII <sub>2</sub> -thien-2-yl	H	-C(0)0CII,
3,5-di-F-ф-	-CH,	П	-CII <sub>1</sub> -φ	11	-C(0)0CII <sub>1</sub> -¢

R2 R3 R4 R5 V			-CII, II -CII,CII(CII,), II -C(0)OC(CII,),	-CII, II Pyrid-2-yl II -C(O)NII,	-CII, II pyrid-3-yl II -C(O)NII,	-CII, II -(CII <sub>2</sub> ),-NHC(O)OC(CII <sub>3</sub> ), II -C(O)OCII <sub>3</sub>	-CII, II -CII <sub>1</sub> CII <sub>1</sub> - $\phi$ II -C(0)OCII,	-CII, II II -C(0)OCII,CII,\$\phi\$	-CII, II -C(0)0CII,CII,¢	-CII <sub>3</sub> II pyrid-4-yl II -C(O)NII <sub>1</sub>	-CH, II -CII(OH)CH, II -C(0)OCH,	-CII, II -CII,CII(CII,), II -C(0)NII,	-CII, II -CIII, II -C(O)NII,	-CII, $II$ -CII <sub>2</sub> - $\phi$ II -C(0)NII <sub>1</sub>	. 11 -CH <sub>2</sub> -φ II 11 -CH(CH <sub>3</sub> ) <sub>2</sub> II	. II -CH <sub>2</sub> -φ H II -CH(CH <sub>3</sub> ), II II pyrid-3-yl II	. II -CH <sub>2</sub> -φ H II -CH(CH <sub>3</sub> ), II -CI II pyrid-3-yl II -CI II -CH <sub>2</sub> CH(CH <sub>3</sub> ), II -C	1         -CH <sub>2</sub> -φ         H           11         -CH(CII <sub>3</sub> )         H           11         pyrid-3-yl         H           11         -CH <sub>2</sub> CH(CH <sub>3</sub> )         H           H         -CH <sub>2</sub> CH(CH <sub>3</sub> )         H
								-CII,			-CII, II	-CII, II	-CII,	-CII, , II	-CII, , II	-CII, , II -CII, II	-	-
2 5.16.8.4.	- m- 1-1n-c'c	3,5-di-F-ф-	3,5-di-F-4-	3,5-di-F-4-	3, 5-di-F-¢-	3,5-di-F-ф-	3, 5-di-F-ф-	3,5-di-F-4-	3,5-di-F-4-	3,5-di-F-φ-	- <del>-</del> -	φ-	÷	φ-	φ φ	φ- φ- 3,5-di-F-φ-	φ- φ- 3,5-di- F-φ- φ-	φ- φ- 3,5-di-F-φ- φ-

R¹	R <sup>2</sup>	R³	R	R5	X
-φ	-CII,	11	-C11 <sub>2</sub> C11(C11 <sub>3</sub> ) <sub>2</sub>	п	-C(0)N(CII))
-φ	-CII,		-CII(CII <sub>3</sub> ) <sub>2</sub>	11	-C(0)N(CII <sub>3</sub> ) <sub>2</sub>
φ.	-CII,	11	-CII <sub>2</sub> -φ	н	·-C(O)NIICII
φ-	-CII,	11	-CII(CII <sub>3</sub> ) <sub>1</sub>	11	-C(O)NIICII,
3,5-di-F-4-	-CII,	=	-CII,CII,CII,	11	-C(0)NIICII,
3,5-di-P-4·	-CII,	11	-CII,CII,CII,CII,	11	-C(0)N(CII)1
3, 5-di-F-ф-	-CII,	=	-ຕາ <sub>ເ</sub> ຕາກູຕາກູຕາເ	Н	-C(0)NII <sub>2</sub>
3,5-di-F-ф-	-CII,	11	m-CII <sub>3</sub> Ο-φ-		-C(0)0C1I,
3,5-di-F-φ-	-CII,	Ξ	$p$ -CII $_{\rm J}$ O- $\phi$ -	II	-C(0)0CII,
3, 5-di-F-ф-	-CII,	11	pyrid-2-yl	11	-C(0)0CII,CII,
3,5-di-F-φ-	-CII,	1	pyrid-4-yl	Н	-C(0)0CH,CH,
cyclohexyl	-CII,	11	-C11 <sub>2</sub> -φ	11	٠٢(٥)٥٠:
cyclopentyl	-CII,	11	-CII <sub>2</sub> -φ	H	-C(0)0CII,
cyclohex-1-enyl	-CH,	=	φ-1II2-	П	-C(0)0CII,
3,5-di-F-φ-	-CII,	11	R4/R5 = -cyclopropyl		·C(0)0CH3
3,5-di-F-ф-	-CII,	II	-CII <sub>3</sub>	11	-C(0)N(CII <sup>3</sup> )CII <sup>5</sup> CII <sup>1</sup> N(CII <sup>3</sup> ) <sup>3</sup>
cyclopropyl	-CII <sub>3</sub>		-C11 <sub>2</sub> φ	=	·C(0)0CH,
3,5-di-F-φ-	-CH,	11	11	Н	-C(0)0CII <sub>1</sub> -\$

	T	T	<del>'</del>	T	T	1	Ī	Τ-	7	T		7		<del>T-</del>	7	<del></del>	7
×	-C(0)0CH,CH,	-C(0)0CII,	-C(O)OCII,CII,	-C(0)0CH,CH,	-C(0)NIIOII	-C(0)0CH,CH(CH,),	-C(0)0CII,	-C(0)0CII.CII.	-C(0)0CHCH3:	-C(O)OCH-CH(CH.)	-C(O)OCH CH	-C(0)0C(I,C)I,	-C(O)NH.	-C(O)N(CH)-OCH	-C(O)NICH,CII(CH,).	-C(O)N(CH,CH,CH,),	THE PARTY OF THE P
ž	=	=	=	=	=	=	=	=	Ξ	=	H	=	H	=	=	=	=
R*	-CII,	-CII <sub>2</sub> -ф	-CII,		-CII(OII)CII,	-CII,	m-hydroxybenzyl	p-liydroxybenzyl	·CII,	ĊII,	-CIII,		·CII	-CIII3	-CII,	-CII,	-CII(CIII)
R³	11	11	11	ш	11	-	II	Н	11	=	Ξ	R <sub>3</sub> /R <sub>4</sub> and N = 2,3-dilydro-indol-2-yl	H	=	11	H	11
R²	φ-	-CII)	-CII,	-CII)	-CII,	φ-	-CII,	-CII)	-CII(CII,)CII,CII,	-CII <sub>1</sub> -ф	-CII,	-CII,	-CII,	φ-	-CII,	-CH <sub>3</sub>	-CII,
R¹	(CII,),CII-	3-NO₂-φ-	3-NO <sub>2</sub> -φ-	3-NO <sub>1</sub> -φ-	3·NO <sub>2</sub> -φ-	(CII,),CII.	3-NO <sub>2</sub> -φ-	3-NO <sub>2</sub> -φ-	(CH <sub>1</sub> ) <sub>1</sub> CH-	(CH <sub>J</sub> ),CH-	3,5-di-F-ф-	3-NO₂-φ-	3,5-di-F-4-	(C11,),C11.	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-6-

	The state of the s		The state of the s		
	R <sup>2</sup>	R³	<b>K</b> 4	R³	×
3,5-di-F	-CII,	П	-CII,	. 11	-C(0)NII-(p-N0 <sub>1</sub> -¢)
3, 5-di-I:-ф-	-CII,	11	-CI1 <sub>2</sub> -\$	=	-C(0)0CII,
3, S-di-l:-φ-	-CII,	11	-CII₂-ф	11	, -C(0)NII,
(CII,),CII-	ф-	=	-CII,	11	-C(O)NIICH2CII(CH3)2
3,5-di-I <sup>-</sup> -φ-	-CII,	11	-CII₂-¢	=	-C(0)NIICII,CII,OCII,
3, 5-di-F-φ-	-CII,	=	-CII <sub>3</sub>	=	-C(0)NIICII <sub>2</sub> -(p-NO <sub>2</sub> -¢)
3,5-di-F-φ-	-CII,	11	-CII <sub>2</sub> -ф	H	-C(0)NII-(p-NO <sub>1</sub> -¢)
3,5-di-l <sup>-</sup> -φ-	-CII,	=	-CII <sub>3</sub>	Н	-C(0)N(CII,)CII,-ф
3,5-di-F	-CII,	=	-CII,	11	-C(0)NIICH <sub>2</sub> -(3,5-di-F¢)
3,5-di-l <sup>2</sup> -φ-	-CII,	=	-CII,	11	-C(0)NHCH <sub>1</sub> -(m-NO <sub>1</sub> -¢)
3,5-di-1:-4-	-CII,		-CII,	Н	-C(0)NIICH-4
3,5-di-F-¢-	-CII,	11	-CII <sub>2</sub> -φ	11	-C(0)NIICII <sub>1</sub> -(p-NO <sub>1</sub> -φ)
3,5-di-F-ф-	-CII,	11	-CII <sub>2</sub> -indol-3-yl	11	·C(0)0CII,
3,5-di-F-¢-	-CH,	-	-CII,	11	-C(0)NIICII1-(p-0CII1-4)
	φ-		-CII,	11	·C(0)0CH,CH,
cyclohexyl-	φ-	=	-CII,	П	-C(0)0CII,CII,
3,5-di-F-¢-	-CII,	11	φ-	11	-C(0)0CII,
3,5-di-F-4-	-CII,	H	-cII <sub>3</sub>	н	-C(0)NHCH,CH,-ф

R¹	R²	R³	אי	' R³	X
3, 5-di-F-4-	-CII,	Е	-CH <sub>2</sub> -indol-3-yl	=	-C(O)NII,
3, 5-di-F-φ-	-CII,	11	-CI12-cyclohexyl	=	C(0)0CII,
3, 5-di-F-ф-	-CII,	=	p-NO <sub>2</sub> -benzyl-	Ξ	-C(0)NIICH,CH,OCH,
3-NO <sub>2</sub> -φ-	-CII,	11	-CII <sub>2</sub> OII	П	-C(0)0CII,CII,
3, S-di-F-ф-	-CII,	11	-CII)	=	-C(0)NIICII(CII))
3, 5-di-F-ф-	-CH,	11	·CII,	11	-C(0)NIICII(CII,)\$
3, S-di-F-ф-	-CII,	-	·CII,	Ξ	-C(0)NIICII <sub>1</sub> -(p-F-¢)
3, 5-di-F-ф-	-CH,	II	-CII,	H	-C(O)NIICII <sub>1</sub> -(pyrid-4-yl)
3,5-di-F-ф-	-CII,	11	-CII,	11	-C(0)NIICII <sub>2</sub> -(p-F <sub>3</sub> C-4)
3,5-di-F-ф-	-CII,	Н	φ.	-CII,	-C(0)OCH,CH,
3,5-di-F-φ-	-CII,	=	-C11 <sub>2</sub> -φ	II	-C(0)0C(CII))
3,5-di-F-φ-	-CII)	11	·CII,	-CII,	·UO(0)O-
3,5-di-F-φ-	-CII,	11	cyclohexyl	H	-C(0)0CII,CII,
3,5-di-F-ф-	-CII,	II	φ-	11	-C(O)NIICII,CII,OCII,
(CII <sub>3</sub> ) <sub>2</sub> CII-	cyclohexyl	11	-CII,	11	-C(0)0CII,CII,
3,5-di-F-ф-	-CII,	Н	φ-	11	-C(O)NIICII,CII,N(CII,),
3,5-di-F-φ-	-CII,	11	φ-	11	-C(0)NIICII <sub>2</sub> -(pyrid-2-yl)
pyrid-3-yl	-СН,	Н	-CII <sub>2</sub> -φ	Н	-с(о)осн,

R¹	R²	Kì	R.	' R³	×
pyrid-2-yl	-CII,	11	-CII₁-ф	Ξ	-C(0)0CII,
pyrid-4-yl	-CII,	==	φ-CI1 <sub>2</sub> -φ	11	-C(0)0CII,
3,5-di-F-4-	-CII,	=	p-F-ф	11	-c(0)0C11,C11,
3, 5-di-F- <b></b>	-CII,	Ш	o-F-¢	=	-C(0)0CII <sub>1</sub> CII <sub>3</sub>
3,5-di-F-φ-	φ-	11	-CII <sub>3</sub>	11	-C(0)0CII,CII,
3,5-di-F-ф-	-CII,	Ξ	-CII <sub>2</sub> -N-phthalimidyl	Ξ	-C(0)0CII,CII,
3,5-di-F- <b>6</b> -	-CIIJ	11	p-Γ-φ	11	-C(0)0CII,C(CII,),
3,5-di-F-ф-	-CH <sub>3</sub>	Ξ	φ-	=	-C(0)0CII,C(CII,),
3,5-di-F-4-	-CII,	II	φ-	b-ct	-C(0)NIIC(CII,)
3,5-di-F-4-	-CII,	11	φ-	11	-C(0)0C(CH <sub>1</sub> ),
3, 5-di-F-ф-	-CII,	Н	φ-	II	-C(0)0C(CII,),
3,5-di-F-ф-	-CII,	П	φ-	Н	-C(0)NH <sub>1</sub>
3-NO <sub>2</sub> -φ-	-CII,	11	-CII(CII <sub>3</sub> ) <sub>1</sub>	11	-C(O)-N-morpholino
3-NO <sub>2</sub> -φ-	-CII,	=	·CII(CII,)	Η	-C(0)0CII,CII,
3-NO <sub>2</sub> -φ-	-CII,	-	-CII(011)CII,	11	-C(0)0CH <sub>1</sub>
3-NO <sub>2</sub> -4-	·CII)	=	-CIICH,	=	-C(O)-N-morpholino
			   O-1-butyl		
3-NO <sub>2</sub> -φ-	-СН,	Н	-СП(СН,)СП,СН,	П	-C(O)-N-morpholino

	I .	T	ī		<u> </u>	1	<u> </u>	ī	Ī		<del></del>	<del></del>	T	1		T	<del></del>	
X	-C(0)0CII,	-C(0)0II	.C(0)0CII,	,-c(0)0CH,	-C(0)0CII,	-C(0)NHCH,CH,OCH,	-C(O)NIICII,CII,N(CII,)	-C(O)NII-cyclohexyl	-C(0)NIICII,C(CII,),	-C(O)NHCH <sub>1</sub> -(tetra- hydrofuran-2-yl)	-C(O)NIICII <sub>2</sub> -(pyrid-2-yl)	-C(0)-(N-thiazolidinyl)	·C(0)0CII,	-C(0)0CII,	-C(0)OCII,	·C(0)NIICII(CII,)CII,CII,	-C(0)-(N-pyrrolidinyl)	-C(0)NHCH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>
1. R3	H	11	11	11	Н	11	11	П	11	11	11	И	11	Н	П	11	Н	Н
R4	-CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	-CH(CH,)CH,CH,	-CII,CII,CII,	·CII,CII(CII,),	-CII,CII(CII,) <sub>1</sub>	-CII,	-CII,	-CII,	-CII,	-CIÍ,	-CII,	-CII,	-CII,CII,	-CII,CII,CH,	-CII,CII,	-CII,	-CII,	-СН,
R³	II	11				hur 1		П	11	=	П	=	=	11 -			Ξ	Н
R1	-CII <sub>3</sub>	-CII,	-CII,	-CII,	-CII,	-CII,	-C11,	-CH,	-CII,	-CII,	-CII,	-CII <sub>3</sub>	-CII,	-CII,	-CII,	-CII,	-CII,	-CH,
R¹	3-NO <sub>1</sub> -φ-	3-NO <sub>2</sub> -φ-	3-NO <sub>1</sub> -ф-	3-NO <sub>2</sub> -φ-	3,5-di-F-4-	3,5-di-F-ф-	3,5-di-F-ф-	3, 5-di-F-φ·	3, 5-di-F-4-	· 3,5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-φ-	3-NO₁-φ-	3,5-di-1∶-φ-	3,5-di-F-¢-	3,5-di-F-φ-

	11,	CII,-F	11(CH <sub>3)1</sub>	λ-NO <sub>2</sub> -φ)	۰-NO <sub>2</sub> -ه)	11,	1,	[]	11,	CII,	CII,	CII,	1,	11,	CII,	I,	11,),	
×	C(0)0CH	-C(Q)NIICII,CII,-F	-C(0)CII,CII,CII(CII,)	-C(0)NIICII,-(p-N0,-4)	-C(0)NIICII <sub>1</sub> -(p-NO <sub>1</sub> -¢)	-C(0)0CII,	-C(0)NII,	-C(O)NII	-C(0)0CII,	-C(0)0CH,CH	-C(0)0CII,CII,	-C(0)0CH2CH3	-C(0)0CII,	-C(0)0CII,	-C(0)0CII,CII,	-C(0)0CII,	-C(0)0C(CII)	
R <sup>5</sup>	11	=	11	11	=	11	1	=	<u>-</u>	11	11	11	=	11	=	11	1	:
R	·CII(CII <sub>1</sub> ) <sub>2</sub>	-CII,	-CII,	-CII,CII,	-CII,CII,CII,	m-F-φ-	thien-2-yl	thien-2-yl	5-chlorobenzo[b]thiophen- 2-yl	benzo[b]thiophen-6-yl	benzo[b]thiophen-2-yl	benzo[b]thiophen-2-yl	beuzo[b]thiophen-3-yl	thien-2-yl	benzo[b]thiophen-5-yl	thien-2-ÿl	thien-2-yl	
R³	11	11	11	Н	11	11	П	11	=	H	П	11	<b>II</b>	11	=	11	11	1
R	-CII,	-CII,	-CIII,	·CII,	-CH3	-CII,	-CII,	-CII,	-CII,	-Cıİ,	·CII)	-CII,	-CII,	-CII,	-CII,	-CII,	-CII,	
R¹	3, 5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-4-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-¢-	3,5-di-F-ф-	3, 5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-ф-	

	1	1	1	ī	<del></del>	_	T	Ţ	<del>-</del>		<del></del>	<del></del>	<del></del>	<del>,</del>		<del></del>	<del>,</del>
×	-C(0)0CII,	-C(0)0CII,	C(0)0CII,	-C(0)0CII,	-C(0)0CH,	-C(0)0CII,	-C(0)0CH,	-C(0)0CII,	-C(0)0C(CII,),	-C(O)NIIC(CII,),	-C(0)0C(CII <sub>1</sub> ) <sub>1</sub>	-C(0)0C(CII,)	-C(0)OCII,	-C(0)0CII,	-C(0)0CII,	-C(0)0CH,	-С(0)ОСН,
' R <sup>5</sup>	=	=	=	=	=	П	П	11	=	H	=	Н	=	=	=	=	11
π,	tetrazof-5-yl	2-aminopyrid-6-yl	6-methoxynaphth-2-yl	<i>m</i> -CF <sub>3</sub> -φ-	4,5,6,7-tetrahydro- benzo{b]thiophen-2-yl	thieno[2,3-b]thiophen-2-yl	2-methylthiazol-4-yl	-CH <sub>2</sub> CH = CHCH <sub>3</sub> (Irans)	φ-	φ-(φ-d)	φ-	φ-	φ-	φ-	φ-	ф.	φ-
R³	II	=	1	П		11	11	=	11	11	11	=	111	=	П	Н	M
R1	-CII,	-CH,	-CII,	-CII)	-CII,	-CII,	-CII,	-CII,	-CII,	-CH,	-CII,CII,	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub> CH <sub>2</sub> SCH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>1</sub>	-CII,CII,	-CII,CII(CII,),	-CH <sub>2</sub> $\phi$
R¹	3,5-di-F:-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-4-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-ф-	cyclopropyl	3,5-di-F-φ-	3,5-di-F-4-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-φ-

				1 1	
R¹	R <sup>2</sup>	R³	. Κ⁴	R³	Х
\$ 5.di-Ε-ά-	II.	11	φ-	П	-C(0)0CII,
3 5-di-Fi-d-	9-	11	φ-	11	.C(0)OC(CII,),
3.5-di-F-o-	φ-	11	<b>\$</b> -	H	-,c(0)0CH
φ-	-CII,	1	-CII,	11	-C(0)0CII <sub>1</sub>
φ	-CII,	11	-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	П	-C(0)0CII,
· •	-CII,	=	-CII(CII,)CII,CII,	П	-C(0)0CII,
φ.	-CII,	R <sub>3</sub> /R <sub>4</sub> and N = L-pyrrolidinyl		H	-C(0)0CII,
÷	·CII,	=	-CII <sub>2</sub> -\$	II	-C(0)0CII,
-φ	.CII,	=	-(CII <sub>3</sub> ),NIIC(O)O- r-butyl	=	-C(0)0C11,
φ-	-CII,	=	11	=	-C(0)0C1I,
φ-	-CII,	H	-CII(CII,),	=	-C(0)0CH,
φ-	-CII,	П	-CII <sub>1</sub> CII <sub>1</sub>	11	-C(0)0C11,
φ-	-CII)	11 ,	-CII,CII,CII,	=	-C(0)0CII,
3-NO <sub>2</sub> -φ-	-CII,	П	-CII(CII <sub>1</sub> ) <sub>2</sub>	Ξ	-C(0)0II
φ.	-CII)	·CII,	·CII,	H	-C(0)0C11,
(CH <sub>1</sub> ),CH-	<del>p</del>	11	-CII,	11	-C(0)0CH,CH(CH,),
(CH <sub>3</sub> ),CH-	-CH(CH3)CH2CH3	Н	-CII,	H	-C(0)0CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>

R¹	R²	R¹	R4	R³	X
3,5-di-F-ф-	-CII <sub>3</sub>	11	φ-	11	-C(O)NII-cyclohexyl
3,5-di-F-6-	·CII,	$R^3/R^4$ and $N = 4 \cdot \beta \cdot lydroxy-$ pyrrolidin-2-yl		=	-С(0)ОСП,СН,
3,5-di-F-4-	-CII,	11	-(CII <sub>1</sub> ),NII <sub>2</sub>	11	-C(0)0CII,
3,5-di-F-4-	-CII,	И	-CII,CII,C(0)NII,	И	-C(0)NII <sub>2</sub>
3, 5-di-F- <i>φ</i> -	-CII,	R <sub>3</sub> /R <sub>4</sub> and N = piperidin-2-yl		н	-C(0)0CH,
3, S-di-F-4-	-CII,	П	11	H	-C(0)NHCII,CII,0CII,
3,5-di-F-ф-	-cyclohexyl	11	φ-	11	-C(0)0CII,
3,5-di-F-4-	-CH,	Н	-CI1 <sub>2</sub> -φ	11	-C(O)NIICII <sub>2</sub> -pyrid-2-yl
3,5-di-F-ф-	-thien-2-yl	11	φ-	11	-C(0)0C(CH))
3,5-di-F-4-	φ	11	φ-	11	-C(O)NH,
3,5-di-F-ф-	-CII(CII <sub>3</sub> ) <sub>2</sub>	11	φ-	11	-C(0)NII <sub>1</sub>
-thienyl	-CII,	11	φ,	И	-C(O)NII,
CII,(CII,) <sub>1</sub> -	-CII,	11	φ-	11	-C(O)NII,
3,5-di-F-φ-	-(CII <sub>2</sub> ),CII <sub>3</sub>	=	φ-	=	-C(0)0CII,
3,5-di-F-¢-	-(CII <sub>1</sub> ) <sub>1</sub> CII <sub>3</sub>		φ-	=	-C(0)0C11,
3,5-di-F-ф-	-C(CII <sub>1</sub> ),	ш	φ-	H	-C(O)OCH,

						T .						T	T		Ī	T	T	T	T
	×	-C(0)0CII'	-C(0)0CII,	-C(0)0CII,	-¢(0)0CII,	-C(0)0CII,	-C(0)0CH,	-C(0)0CII,	-C(0)OC(CH')	-C(0)OC(CII,),	(11)))(()))	-CONNIT	CIONINIO)	-C(O)NIIC(CII).	-CONNICCITY	-C(O)NHC(CH),	-C(O)NIIC(CII.)	-C(O)NIICICILI).	(01)
	1,4%	п	=	=	=	Е	H	H	н	=	=			=	=	=	=	=	-
n.4	Κ.	Φ-	φ-	<i>p</i> -	φ-	ф	φ-	φ-	φ-	φ-	φ-	φ-	5-bromothien-2-yl	4-bromothien-2-yl	thien-2-yl	thien-3-yl	φ.	5-chlorothien-2-yl	
103	4	11	П	11	11	11	11	11	II	Н	11	11	11	11	11	11		11	1
P2		-C(CII,)CII,CII,	-CII <sub>2</sub> $\phi$	-cyclopropyl	-thien-3-yl	-thien-2-yl	p-F-φ	$p$ -OCH <sub>3</sub> - $\phi$	φ-	φ-	9	-CH,	-CII)	-CII,	-CII,	-CII,	-CII,	-CII,	
~~		3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3, 5-di-F-φ-	3,5-di-F-¢-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	-cyclopropyl	-cyclopentyl	-C(CH <sub>3</sub> ),	3,5-di-F-φ-	3,5-di-F-ø-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-¢-	2 5 4: E.A.

R	R²	R³	R*	R <sup>5</sup>	×
3.5-di-F-d-	-CII <sub>1</sub>	11	(m-phenoxy)-φ-	И	-C(O)NIIC(CII,)
3 5-(li-F-φ-	-CIII,		φ.	11	C(O)NII-cyclohexyl
3,5-di-1 <sup>2</sup> -φ-	-CII)	11	-φ-(φ- <i>ιιι</i> )	1	-C(O)NIIC(CII,),
3, 5-di-F-φ-	-CH,	1	(ρ-CII <sub>3</sub> CII <sub>2</sub> )-φ-	11	-C(O)NHC(CH <sub>3</sub> ),
3,5-di-F-ф-	-CII,	П	$-\phi$ - $(\phi$ - $o)$	П	-C(0)NIIC(CII)3
3,5-di-F-ф-	-CII,	11	(ο-henzyl)-φ-	П	-C(O)NIIC(CII))
3,5-di-F-4-	-CII,	П	(ρ-Βr)-φ-	П	-C(0)NIIC(CII <sub>j</sub> ),
3,5-di-F-ф-	-CH <sub>3</sub>	11	-φ-(φ-d)	=	-C(O)NIIC(CII <sub>3</sub> )3
3,5-di-F-ф-	-CII,	11	$(\rho\text{-CII}_3\text{CII}_2\phi)$ - $\phi$ -	=	-C(0)NIIC(CII <sub>3</sub> ),
3.5-di-F	·CII,	1	(p-tert-butyl)-4-	11	-C(0)NIIC(CII,)
3, S-di-F-φ-	-CH)		o-(4-Cl-phenoxy)-φ-	=	-C(0)NIIC(CII <sub>3</sub> ),
3,5-di-1 <sup>-</sup> -ф-	-CII,	П	-φ-(φ-d)	11	-C(O)NII-cyclohexyl
3,5-di-F-ф-	-CII,	П	φ-	=	-C(0)OC(CH <sub>3</sub> ),
3,5-di-F-ф-	-CII <sub>3</sub>	н	-CII,	11	φ-(O)-φ
3,5-di-F-ф-	-CII,	11	pyrid-3-yl	11	-C(0)OC(CII))
3, 5-di-F-4-	-CII,	H	φ-	11	.C(O)-morpholino
3,5-di-F-ф-	-CII,	И	(m-methoxy)-φ	Н	-C(0)0CII,
3,5-di-F-ф-	-CH,	Н	φ-	н	-C(0)ONHC(0)OC(CH)),

				-	
R¹	R²	R³	×.	~	X
3,5-di-1:-ф-	-CII)	П	ф-	П	-C(O)NIIC(CII.),
3,5-di-F-¢-	·CII,	II	φ-	=	-C(O)NIICII, furan-2-vi
3,5-di-F-ф-	-C11,	11	<del>9</del> -	=	-C(0)NHOCH,
3,5-di-F-4-	-CII,	11	φ.	=	-C(O)-cyclobutylamide
3,5-di-F-φ-	-CII,	11	φ-	=	-C(0)NIICII,CII(CII,),
3,5-di-F-φ-	-CII,	Н	φ-	=	-C(0)NHCII <sub>2</sub> -cyclopropyl
3,5-di-F-6-	-CII,	11	φ-	=	-C(O)N(CH,)OCH,
3,5-di-F-φ-	-CII,	11	φ-	=	-C(0)NIICII,C(CII,)=CII,
3,5-di-F-¢-	-CII)	=	φ-	=	-C(0)CH,-pyrid-3-yl
3,5-di-F-ф-	-CII,	==	φ-	=	-C(0)CH,-pyrid-4-yl
3,5-di-F-¢-	-CII)	11	φ.	=	-C(0)CII <sub>3</sub> -fur-2-yl
3,5-di-F-4-	-CII,	Н	φ.	=	C(0)NII-cyclonentyl
3,5-di-F-4-	-C11,	II	φ.	=	-C(O)NII-[(p-benzyl)-pyrid-4-yl]
3,5-di-F-¢-	-CII,	11	φ.	=	-C(0)N(CII,),
3,5-di-F-ф-	-CII,	1	φ.	=	-C(O)NH-[2,6-dimethylpyrid-4-yl]
3,5-di-F-ф-	-CH <sub>3</sub>	-	φ-	H	-C(O)NII-(2-nethylcyclollexyl)
3,5-di-F-φ-	-CH,	in in its second	<del>p</del>	H	-C(O)NH-(4-methylcyclohexyl)


X	-C(O)NH-[N-ethoxycarbonyl]- piperidin-4-yl	-Ċ(O)NIICII,	-C(0)NIIOC(CII))	·C(O)ONIIC(CII)	-C(0)NIINH1	$-C(O)NIIN = C(CII_1)OCII_1CII_1$	-C(0)OC(CII,),	-C(0)ΝΙΙ(CΙΙ <sup>)</sup> ),·φ	-C(O)NH(CH <sub>2</sub> ),O-(p-CH <sub>3</sub> )¢	-C(O)NII(CII,),NII,	-C(0)0CII,	-C(0)0II	-C(0)OC(CH,),	-C(0)OC(CII,),	-С(О)ОСН,	-C(0)OCH3
R3	11	=	=	11	11	11	II	Ξ	Н	=	=	11	=	11	=	H
R*	ф-	φ-	ф-	φ-	φ-	φ-	φ-	φ-	φ-	φ-	3,5-di-F-ф-	-ຕາ <sub>ເ</sub> ຕາ <sub>ເ</sub> ຕາ <sub>ເ</sub> ຕາເ,	φ-	φ.	,	
R³	Ξ	11	II	н	11	П	П	11	П	11	=	11	Ξ	11	$R_J/R_c$ and $N \approx L-(3-\alpha-phenyl)$ . pyrrolidinyl	$R_{\rm J}/R_{\rm d}$ and N = L-azetidin-2-yl
R²	-CIII,	-CII,	-CII,	·CII,	-CII,	-CII,	-CII,	-CII,	-CII,	-CII,	3,5-di-F-ф-	-CII,	-CII,	thien-2-yl	-CII,	·CII]
R¹	3,5-di-F-ф-	3,5-di-F	3,5-di-F-ф-	3,5-di-F-4-	3,5-di-F-ф-	3,5-di-F- <b></b>	φ-	3,5-di-F-4-	3,5-di-F-4-	3,5-di-Γ-φ-	3, 5-di-I <sup>2</sup> -φ-	3,5-di-F-4-	-cyclopentyl	3,5-di-I <sup>2</sup> -ф-	3,5-di-F-φ-	3,5-di-F-φ-

3.5 di.F.φ.         -CII,         II         5-chlorobenzolbliniophen-3-yI         II         -COOCCII,           3.5 di.F.φ.         -CII,         II         -CHI-thiazol 4-yI         II         -COOCCII,           3.5 di.F.φ.         -CII,         II         +φ         II         -COONII,           3.5 di.F.φ.         -CII,         II         +φ         II         -COONII,           3.6 di.C.φ.         -CII,         II         +φ         II         -COONII,           3.6 di.C.φ.         -CII,         II         +φ         II         -COONII,           3.6 di.C.φ.         -CII,         II         +φ         II         -COONII,           4.F.φ.         -CII,         II         +φ         II         -COONII,           4.F.φ.         -CII,         II         +φ         II         -COONII,           3.C.q.φ.         -CII,         II         +φ         II         -COONII,           4.C.II.φ.         -CII,         II         +φ         II         -COONII,           3.C.q.φ.         -CII,         II         +φ         II         -COONII,           3.C.q.φ.         -CII,         II         +φ         -COONII,	R¹	R²	R³	κ⁴	R³	X
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3,5.di-F-¢-	-CII,	11	5-chlorobenzo[b]thiophen- 3-yl	11	·IIDO(0)D
CH,         II         -φ         II           -CH,         II         thien-2-yI         II           -CH,         II         -φ         II         -φ           -CH,         II         -φ         II         -φ <t< td=""><td>3,5-di-F-ф-</td><td>-CII,</td><td>11</td><td>-CH<sub>2</sub>-thiazol-4-yl</td><td>11</td><td>-C(0)OC(CH))</td></t<>	3,5-di-F-ф-	-CII,	11	-CH <sub>2</sub> -thiazol-4-yl	11	-C(0)OC(CH))
CH, H         thien-2-yl         II           -CH, H         +φ         II           -CH, H         +φ         II           -CH, H         H         +φ         II           -CH, H         II         +φ         II           -CH, H         H         +φ         II           -CH, H         H         +φ         II           -CH, H         H         +φ         II           -CH, H         +φ         II         H	3,5-di-F-ф-	-CII <sub>1</sub>	П	φ-	=	-C(O)NIIC(CII))
CH,         H         -φ         H           -CH,         H         -φ	3,5-di-F-φ-	-CH3	Н	thien-2-yl	=	-C(0)NII,
-CH <sub>3</sub> H1 -φ H1 H2 H1 H2 H2 H1 H2	3,4-di-CI-ф-	-CH <sub>3</sub>	Н	φ.	н	-C(O)NH <sub>1</sub>
-CII, II -φ II II - η - η II	3-CI-φ-	-CII <sub>3</sub>	111,	φ-	И	-C(0)NII,
-CII,5         II         -φ         II           -CII,5         II         φ         II           -CII,5         II         φ         II           -CII,5         II         -φ         II           -CII,7         II         -φ         II           -CII,5         II         -φ         II           -CII,7         II         -φ         III	3-Br-ф-	-CIII3	П	φ-	=	-C(0)NII,
-CH <sub>3</sub> H -C	3.F-ф-	-CII,	11	φ-	=	-C(O)N11 <sup>1</sup>
-CII,         II         -φ         II	4-F-¢-	-CIII3	п	φ-	=	-C(O)NII
-CH <sub>3</sub>	3-CII <sub>3</sub> -φ-	-CII,	11	φ-	=	-C(O)NII,
-CII, ' II -φ II II -φ III	4-CII <sub>3</sub> -φ-	-CIIs	H	φ-	=	-C(0)NII,
-CH, II -φ II   II   II   II   II   II   II	3-CF3-4-	-CII,	, 11	φ-	×	-C(O)NII,
-CII, II -φ II -CII, II -φ II -CII, II -φ II   II   II   II   II   II   II	3-CII₃O-φ-	-CII,	11	φ-	=	-C(O)NII,
-CII, II -φ II -CII, II -φ II -CII, II -φ II	2-CI-φ-	-CII,	11	φ-	=	-C(O)NH,
-CII, II -\$p\$ III -\$p	1-naphthyl	-CII <sub>3</sub>	Н	φ-	11	-C(0)NII,
-CII, II -ф II -CH, H -ф II	2-naphthyl	-CII,	11	φ-	=	-C(0)NH,
-Сп, н	-ф	-CII,	Ш	φ-	11	-C(0)NII,
	3, 5-di-F-ф-	-CII,	Н	φ-	=	Н00Э-

		T	7	7		T	T	T		T	<u>.</u> T	_	T	7	T		_	_	7		<del></del>		<del>~~~</del>
	×	CONNIC		-C(O)NII,	-C(O)NII-SO <sub>2</sub> -CII,	-C(O)N(CII')®	ACHON(O)D-	A(III)	I I I I I I I I I I I I I I I I I I I	COMMICTION TO	(φ-1-φ)	-C(0)0CII,C(CII,),	11000070	-(1)0(11)	-C(O)OCH,	-C(0)0CII,	-C(0)0CII,	1150005-	COMM	-(C)trit-(4-memyiphenyi)	-C(O)NII-2-tetrahydrofurfuryl	-C(0)NII,	-C(O)NH,
	RS	=	: :	= -	Ξ	=	=	=	=	:  =	:	=	=	: =		=	Ξ	H	=	: :	=	=	Ξ
		furan-2-y]	*	<b>*</b>	-CII <sub>2</sub> -\$	11	-CII,	φ-		φ-		4-fluorophenyl	φ	-6		Φ-	φ-	φ.	φ-	4		φ-010π0-φ	φ
	R³	11	11	:	1	=		II	=	=		=		=	11	:	=	н	11				H
	R <sup>2</sup>	-CII,	-CII,	115	الن-	-CH <sub>3</sub>	-CII,	-CII,CII,SCII,	-CII,	-CII,	110-	(17)	-CII,	-CII,OCII <sub>2</sub> -\$	-CHCHOOCH-4	47	-CII(OH)CII,	-CII,0II	-CII,	-CII,	,CII,		-CH,CH,SCH,
-	R	3,5-di-F-4-	3,4-di-F-ф-	2 S.di. F.A.	- A-1-10-C'C	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-ф-	3.5-di-F-A-		2,3,4,5,6-penta-F-φ-	3, 5-di-F-4-	3,5-di-F-ф-	- L : 0 6	-\$-,I-ID-C'C	3, 5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-6-		3, J-dI-F-φ-

R³		R4	R\$	×
-CH <sub>1</sub> CH,	11	φ-	11	-C(0)NII <sub>3</sub>
φ	П	φ-	=	-C(0)NII,
-CII(CII <sub>1</sub> ) <sub>2</sub>	11	φ-	i	-C(0)NH <sub>1</sub>
-CII,	11	φ-	-=	-C(0)NHCH(CH3)4
-CII,	П	φ-	=	C(0)CII(CII,),
-CII,	=	ф-	H	-C(0)CII,
-CH,	11	φ-	11	-C(0)CII,CII,CII,
-CII,	П	φ-	н	φ-(O)Ͻ-
-CII,	11	φ-	н	-C(O)CH <sub>1</sub> CH <sub>1</sub>
-CII,	II	φ-	=	-C(0)CII,CII(CII,),
-CII,	11	φ-	H	-C(0)0CII,
-CII,	=	φ-	Ш	-C(O)NH(CH <sub>2</sub> ),NHC(O)- (2-hydroxy-4-azido)-
				phenyl
-CII,	11	4-Nuoro-φ	И	-C(0)0C(CII,),
-CII,	=	4-φ-φ-	Н	-C(0)0C(CII,),
-CII,	R <sub>y</sub> /R <sub>4</sub> and N = 3,3-diliydro-2-isobenzazolyl	•	11	-C(0)OCH,
-CII,	H	4-n-butylphenyl	Н	-C(O)NHC(CH,),

	T	<del></del>	T	T	Т	_	T -	T		1	1	7	T	7	Τ-	7
X	-C(O)NHC(CH <sub>1</sub> ),	-C(S)NH,	-C(0)CH <sub>2</sub> -\$	-C(0)-cyclopentyl	-C(O)-n-butyl	-с(о)сн(сн;)сн;сн;	-C(O)NH(CH <sub>1</sub> ) <sub>8</sub> -(biotinamidyl)	-C(0)0CH	-C(0)OCH	-C(O)NHC(CH;),	-с(0)осн	но(о)о-	-C(0)OC(CH <sub>1</sub> ),	-C(0)0CH,	-C(0)0CH,	-С(0)ОСН,
R3	Ħ	H	H	H	Ħ	н	Ξ	Ħ	×	н	H	Н	Н	Н	Н	Н
R*	4-phenylacetylenylphenyl	φ-	φ-	φ-	φ-	φ-	φ-	-CH,CH,SCH,	ф	2-F-4-	φ-	φ-	φ-	φ-	φ-	φ-
R³	Н	Н	Н	Н	Н	Н	H	Н	11	Н	I	Н	H	Н	H	H
R <sup>1</sup>	-СН,	-СН,	-СН,	-сн,	-CH3	-СН,	-CH,	-СН,СН,ЅСН,	-CH,	-СН,	-СН,	thien-3-yl	thien-3-yl	-CH,	-СН,	-сн,
R¹	3,5-di-F-4-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-φ-	3,5-di-F-4-	3,5-di-F-ф-	1-BOC-NII-CII <sub>1</sub> -	3,5-di-F-ф-	3,5-di-F-ф-	3,5-di-F-φ-	. 3,5-di-F-φ-	(2,5-di-Cl-φ)-O-	(3,5-di-F-φ)-O-	(3,4-di-Cl-4)-S-



	T	7	T	T	<del>T .</del>	T	T.	Т.	<del></del>	7	<del>1-</del>
<b>"</b> \	H		H	H	H	H	Ħ	H	H	H	H
λ,	-CH <sub>2</sub> C(0)0CH <sub>1</sub>	-CH <sub>2</sub> CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	Н	H	H	φ-	φ-	Н	H	H	H
R <sup>4</sup>	$\text{-CH}_2\phi$	-CH <sub>3</sub>	φ-	-CH <sub>2</sub> $\phi$	-CII <sub>3</sub>	-CH <sub>3</sub>	φ-	-(CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	-CH <sub>2</sub> -( $p$ - hydroxyphenyl)	2-pyridyl	4-pyridyl
R³	H	Н	Н	H	Н	Н	Н	H	Н	H	Н
R <sup>2</sup>	-CII,	-CH,	-CH3	-CH3	-CH3	-CH3	-СН,	-CH3	-СН,	-СН,	-CH3
R¹	3,5-di-F-4-	3,5-di-F-4-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3,5-di-F-φ-	3, 5-di-F-φ-	3,5-di-F-φ-

R¹	R <sup>2</sup>	R³	$ m R^4$	Υ'	,'Å
3,5-di-F-φ-	-CH3	Н	-CII <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	H
3,5-di-F-4-	-CH3	Н	-CH(CH <sub>3</sub> ) <sub>2</sub>	H	Н
3,5-di-F-6-	-CH3	Н	-CH <sub>3</sub>	H	Н
3,5-di-F-¢-	-CH3	Н	thien-2-yI	H	H
3,5-di-F-φ-	-СН3	Н	φ-	-CH <sub>3</sub>	-CH3
3,5-di-F-φ-	φ-	Н	φ-	Н	Н
-cyclopropyl	φ-	Η	φ-	Н	Н
-cyclopentyl	φ-	H	φ-	H	Н

3
_
<u>-</u>

R¹	R²	R³	R4	ð	Υ'
3,5-di-F-φ-	-СН,	Н	-CH3	-OCH3	H
3,5-di-F- <i>φ</i> -	-CH3	H	φ-	-OCH <sub>3</sub>	Н
3,5-di-F-4-	-CH3	Н	$\text{-CII}_2\phi$	-OCH <sub>3</sub>	Н
3,5-di-F-4-	-CH3	H	-CH2CH2CH2CH3	-OC(0)CII3	П
3,5-di-F-ф-	-CH3	II	-CH,CH,CH,CH,	-0C(0)C(CH <sub>3</sub> )3	11
3,5-di-F-ф-	-CH3	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	-phthalimido	Н

>	~_~~~
IZ	
<u>~</u>	ŹΙ
0=	~ ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~
IZ	= =0
•	<u>r</u>

	<del></del>	<del></del>	<del></del>	<del></del>				_
×	-C(0)0CII2CH(CH1),	-C(0)NH,	-C(0)NH(p-NO,-φ)	-C(0)0CH	-C(0)0CH <sub>1</sub>	-C(0)0CH,CH,	-СН2ОН	-СН,ОН
R4	-CH <sub>3</sub>	-CII <sub>2</sub> -\$	-CH <sub>3</sub>	φ-	φ	-CH(CH <sub>3</sub> ) <sub>2</sub>	φ-	Н
]¥⁴′	φ-	-CH <sub>3</sub>	-CH <sub>3</sub>	-CII <sub>2</sub> -φ	-CH3	-CH(OH)CH,	-CH3	φ-
R <sup>2</sup>	(CH <sub>3</sub> ) <sub>2</sub> CH-	φ-	φ-	-CH <sub>3</sub>	-CH <sub>3</sub>	-CH3	φ-	-CH3
R¹	(CH <sub>3</sub> ) <sub>2</sub> CH-	(CH <sub>3</sub> ),CH-	(CH <sub>3</sub> )2CH	3, 5-di-F-φ-	3,5-di-F-φ-	3-NO <sub>2</sub> -φ-	3,5-di-F-φ-	3, 5-di-F-φ-

	× /
₩—	/
°=(	ŽI
\ ئوسەسىيە	~ ~ ~
×××	<b>&gt;</b>
×	<u>~</u>

R¹	X,	,,X	R <sup>2</sup>	R⁴	X
3,5-di-F-φ-	но-	Н	-CH <sub>3</sub>	φ-	-C(O)NHC(CH,),
3,5-di-F-4-	4-	Ħ	-CH <sub>3</sub>	φ-	-C(O)NHC(CH,),
3,5-di-F-ф-	= "X//X"		-CH3	φ-	-C(0)OC(CH <sub>3</sub> ) <sub>3</sub>
	0=				

## DETAILED DESCRIPTION OF THE INVENTION

As above, this invention relates to methods for inhibiting  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. However, prior to describing this invention in further detail, the following terms will first be defined.

## **Definitions**

5

10

The term " $\beta$ -amyloid peptide" refers to a 39-43 amino acid peptide having a molecular weight of about 4.2 kD, which peptide is substantially homologous to the form of the protein described by Glenner, et al. including mutations and post-translational modifications of the normal  $\beta$ -amyloid peptide. In whatever form, the  $\beta$ -amyloid peptide is an approximate 39-43 amino acid fragment of a large membrane-spanning glycoprotein, referred to as the  $\beta$ -amyloid precursor protein (APP). Its 43-amino acid sequence is:

- 15 Asp Ala Glu Phe Arg His Asp Ser Gly Tyr
  - 11 Glu Val His His Gln Lys Leu Val Phe Phe
  - 21 Ala Glu Asp Val Gly Ser Asn Lys Gly Ala
- 20 31
  Ile Ile Gly Leu Met Val Gly Gly Val Val
  - 41 Ile Ala Thr (SEQ ID NO: 1)

or a sequence which is substantially homologous thereto.

"Alkyl" refers to monovalent alkyl groups preferably having from 1 to 10 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, n-hexyl, and the like.

5

10

15

25

"Substituted alkyl" refers to an alkyl group, preferably of from 1 to 10 carbon atoms, having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted alkoxy, aryloxy, heteroaryloxy, heterocyclyloxy, acyl, acylamino, amino, aminoacyl, aminocarboxy esters, cyano, cycloalkyl, halogen, hydroxyl, carboxyl, carboxylalkyl, oxyacyl, oxyacylamino, thiol, thioalkoxy, substituted thioalkoxy, aryl, heteroaryl, heterocyclic, nitro, and mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic.

"Alkylene" refers to divalent alkylene groups preferably having from 1 to 10 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methylene (-CH<sub>2</sub>-), ethylene (-CH<sub>2</sub>CH<sub>2</sub>-), the propylene isomers (e.g., -CH<sub>2</sub>CH<sub>2</sub>-CH<sub>2</sub>- and -CH(CH<sub>3</sub>)CH<sub>2</sub>-), and the like.

"Alkaryl" refers to -alkylene-aryl groups preferably having from 1 to 10 carbon atoms in the alkylene moiety and from 6 to 10 carbon atoms in the aryl moiety. Such alkaryl groups are exemplified by benzyl, phenethyl and the like.

"Alkoxy" refers to the group "alkyl-O-". Preferred alkoxy groups include, by way of example, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, ten-butoxy, sec-butoxy, n-pentoxy, n-hexoxy, 1,2-dimethylbutoxy, and the like.

"Substituted alkoxy" refers to the group "substituted alkyl-O-" where substituted alkyl is as defined above.

5

10

15

20

25

"Alkylalkoxy" refers to the group "-alkylene-O-alkyl" where alkylene and alkyl are as defined above. Such groups include, by way of example, methylenemethoxy (-CH<sub>2</sub>OCH<sub>3</sub>), ethylenemethoxy (-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>), n-propylene-iso-propoxy (-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH(CH<sub>3</sub>)<sub>2</sub>), methylene-t-butoxy (-CH<sub>2</sub>O-C(CH<sub>3</sub>)<sub>3</sub>) and the like.

"Alkylthioalkoxy" refers to the group "-alkylene-S-alkyl" wherein alkylene and alkyl are as defined above. Such groups include, by way of example, methylthiomethoxy (-CH<sub>2</sub>SCH<sub>3</sub>), ethylthiomethoxy (-CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>), n-propyl-iso-thiopropoxy (-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>SCH(CH<sub>3</sub>)<sub>2</sub>), methylthio-t-butoxy (-CH<sub>2</sub>SC(CH<sub>3</sub>)<sub>3</sub>) and the like.

"Alkenyl" refers to alkenyl groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation. Preferred alkenyl groups include ethenyl (-CH=CH<sub>2</sub>), n-propenyl (-CH<sub>2</sub>CH=CH<sub>2</sub>), iso-propenyl (-C(CH<sub>3</sub>)=CH<sub>2</sub>), but-2-enyl (-CH<sub>2</sub>CH=CHCH<sub>3</sub>), and the like.

"Substituted alkenyl" refers to an alkenyl group as defined above having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, amino, aminoacyl, aminocarboxy esters, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, cycloalkyl, oxyacyl, oxyacylamino, thiol, thioalkoxy, substituted thioalkoxy, aryl, heteroaryl, heterocyclic, nitro, and mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic.

"Alkynyl" refers to alkynyl groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1

5

10

15

20

and preferably from 1-2 sites of alkynyl unsaturation. Preferred alkynyl groups include ethynyl (-CH  $\equiv$  CH<sub>2</sub>), propargyl (-CH<sub>2</sub>C  $\equiv$  CH) and the like.

"Substituted alkynyl" refers to an alkynyl group as defined above having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, amino, aminoacyl, aminocarboxy esters, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, cycloalkyl, oxyacyl, oxyacylamino, thiol, thioalkoxy, substituted thioalkyoxy, aryl, heteroaryl, heterocyclic, nitro, and mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic.

"Acyl" refers to the groups alkyl-C(O)-, substituted alkyl-C(O)-, cycloalkyl-C(O)-, aryl-C(O)-, heteroaryl-C(O)- and heterocyclic-C(O)- where alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Acylamino" refers to the group -C(O)NRR where each R is independently hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclic and where each of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Aminoacyl" refers to the group -NRC(O)R where each R is independently hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclic and where each of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

10

20

25

"Oxyacyl" refers to the groups -OC(O)-alkyl, -OC(O)-aryl, -C(O)O-heteroaryl-, and -C(O)O-heterocyclic where alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Oxyacylamino" refers to the groups -OC(O)NR-alkyl, -OC(O)NR-substituted alkyl, -OC(O)NR-aryl, -OC(O)NR-heteroaryl-, and -OC(O)NR-heterocyclic where R is hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclic and where each of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Aminocarboxy esters" refers to the groups -NRC(O)O-alkyl,
-NRC(O)O-substituted alkyl, -NRC(O)O-aryl, -NRC(O)O-heteroaryl, and
-NRC(O)O-heterocyclic where R is hydrogen, alkyl, substituted alkyl,
cycloalkyl, aryl, heteroaryl, and heterocyclic and where each of alkyl,
substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic are as defined
herein.

"Aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl). Preferred aryls include phenyl, naphthyl and the like.

Unless otherwise constrained by the definition for the aryl substituent, such aryl groups can optionally be substituted with from 1 to 5 and preferably 1 to 3 substituents selected from the group consisting of hydroxy, biotinamidyl, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, aminoacyl, aminocarboxy esters, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, acylamino, cyano, halo, nitro, heteroaryl, heterocyclic, oxyacyl, oxyacylamino, thioalkoxy, substituted thioalkoxy, trihalomethyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-

15

20

heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric disubstituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic, and the like. Preferred substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy.

"Aryloxy" refers to the group aryl-O- wherein the aryl group is as defined above including optionally substituted aryl groups as also defined above.

The term "carboxy terminal  $R^4$  group" refers to that  $R^4$  group in compounds of formula I which, when n is two, is closest to the X group.

"Carboxyalkyl" refers to the groups -C(O)O-alkyl and -C(O)O-substituted alkyl where alkyl and substituted alkyl are as defined above.

"Cycloalkyl" refers to cyclic alkyl groups of from 3 to 10 carbon atoms having a single cyclic ring or multiple condensed rings which can be optionally substituted with from 1 to 3 alkyl groups. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, 1-methylcyclopropyl, 2-methylcyclopentyl, 2-methylcyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

"Cycloalkenyl" refers to cyclic alkenyl groups of from 4 to 8 carbon atoms having a single cyclic ring and at least one point of internal unsaturation which can be optionally substituted with from 1 to 3 alkyl groups. Examples of suitable cycloalkenyl groups include, for instance, cyclobut-2-enyl, cyclopent-3-enyl, cyclooct-3-enyl and the like.

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo and preferably is either chloro or bromo.

10

15

25

"Heteroaryl" refers to a monovalent aromatic group of from 2 to 10 carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and sulfur within the ring.

Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with 1 to 3 substituents selected from the group consisting of hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, aminoacyl, aminocarboxy esters, alkaryl, aryl, aryloxy, carboxyl, carboxylalkyl, aminoacyl, cyano, halo, nitro, heteroaryl, heterocyclic, oxyacyl, oxyacylamino, thioalkoxy, substituted thioalkoxy, trihalomethyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic, and the like. Preferred substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy.

Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indolizinyl or benzothienyl). Preferred heteroaryls include pyridyl, pyrrolyl and furyl.

20 "Heteroaryloxy" refers to the group heteroaryl-O- wherein the heteroaryl group is as defined above including optionally substituted heteroaryl groups as also defined above.

"Heterocycle" or "heterocyclic" refers to a monovalent (i.e., one point of attachment) saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 8 carbon atoms and from 1 to 4 hetero atoms selected from nitrogen, sulfur or oxygen within the ring.

Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 4 substituents selected from the group consisting of hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, aminoacyl, aminocarboxy esters, alkaryl, aryl, aryloxy, carboxyl, carboxylalkyl, aminoacyl, cyano, halo, nitro, heteroaryl, heterocyclic, oxyacyl, oxyacylamino, thioalkoxy, substituted thioalkoxy, trihalomethyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic, and the like. Such heterocyclic groups can have a single ring or multiple condensed rings. Preferred heteroaryls include morpholino, piperidinyl, and the like.

15

20

5

10

Examples of heterocycles and heteroaryls include, but are not limited to, furan, thiophene, thiazole, oxazole, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, phthalimide, 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholino, piperidinyl, pyrrolidine, tetrahydrofuranyl, and the like.

25

"Heterocyclyloxy" refers to the group heterocyclyl-O- wherein the heterocyclic group is as defined above including optionally substituted heterocyclic groups as also defined above.

20

"Oxyacyl" refers to the groups -OC(O)-alkyl, -OC(O)-aryl, -C(O)O-heteroaryl-, and -C(O)O-heterocyclic where alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Oxyacylamino" refers to the groups -OC(O)NH-alkyl, -OC(O)NH-5 substituted alkyl, -OC(O)NH-aryl, -OC(O)NH-heteroaryl-, and -OC(O)NH-heterocyclic where alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Thiol" refers to the group -SH.

"Thioalkoxy" refers to the group -S-alkyl.

"Substituted thioalkoxy" refers to the group -S-substituted alkyl.

"Thioaryloxy" refers to the group aryl-S- wherein the aryl group is as defined above including optionally substituted aryl groups also defined above.

"Thioheteroaryloxy" refers to the group heteroaryl-S- wherein the heteroaryl group is as defined above including optionally substituted aryl groups as also defined above.

"Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound of Formula I which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like.

## Compound Preparation

The compounds of formula I are readily prepared via several divergent synthetic routes with the particular route selected relative to the ease of compound preparation, commercial availability of starting materials, etc.

A first synthetic method involves conventional coupling of an acetic acid derivative with a primary amine of an esterified amino acid as shown in reaction (1) below:

10
$$X' \longrightarrow OH + H_2N \longrightarrow XR^3$$

$$1 \qquad 2$$

$$X' \longrightarrow X'' \longrightarrow XR^3$$

$$R^1 \longrightarrow XR^3$$

$$R^1 \longrightarrow R^2 \longrightarrow XR^3$$

$$20 \qquad 3$$

wherein  $R^1$ ,  $R^2$ ,  $R^3$ , X' and X'' are as defined above, and X is either oxygen or -NH-.

Reaction (1) merely involves coupling of a suitable acid derivative 1 with the primary amine of amino acid ester 2 under conditions which provide for the N-acetyl derivative 3. This reaction is conventionally conducted for

WO 98/22494

5

10

15

20

25

-- 77 --

PCT/US97/20804

peptide synthesis and synthetic methods used therein can also be employed to prepare the N-acetyl amino acid esters 3 of this invention. For example, well known coupling reagents such as carbodiimides with or without the use of well known additives such as N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. can be used to facilitate coupling. The reaction is conventionally conducted in an inert aprotic diluent such as dimethylformamide, dichloromethane, chloroform, acetonitrile, tetrahydrofuran and the like. Alternatively, the acid halide of compound 1 can be employed in reaction (1) and, when so employed, it is typically employed in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, triethylamine, diisopropylethylamine, N-methylmorpholine and the like.

Reaction (1) is preferably conducted at from about  $0^{\circ}$ C to about  $60^{\circ}$ C until reaction completion which typically occurs within 1 to about 24 hours. Upon reaction completion, N-acetyl amino acid ester 3 is recovered by conventional methods including precipitation, chromatography, filtration and the like or alternatively is hydrolyzed to the corresponding acid without purification and/or isolation other than conventional work-up (e.g., aqueous extraction, etc.). Alternatively, the synthesis described above in reaction (1) can be conducted on the amino acid (XR<sup>3</sup> = OH) and subsequent to N-acetyl formation as described above.

In any event, if an N-acetyl amino acid ester is formed, it is converted to the corresponding acid prior to the coupling step with another amino acid ester/amide, HNR<sup>3</sup>CR<sup>4</sup>R<sup>5</sup>C(O)Y. Coupling is accomplished using well known peptide coupling chemistry with well known coupling reagents such as carbodiimides with or without the use of well known additives such as N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. which can be used to facilitate coupling. The reaction is conventionally conducted in an inert aprotic polar diluent such as dimethylformamide, dichloromethane, chloroform, acetonitrile, tetrahydrofuran and the like.

10

15

20

25

Such coupling yields compounds of formula I where n is 1. The synthesis of compounds of formula I where n is 2 is accomplished via a second coupling reaction. Specifically, in the first coupling reaction, HNR<sup>3</sup>CR<sup>4</sup>R<sup>5</sup>C(O)Y is selected to be an amino acid ester. That is to say that Y is -O-alkyl. After coupling, the ester is hydrolyzed via conventional conditions well known in the art to provide for the corresponding carboxylic acid which can now be used to couple a second amino acid ester/amide.

In reaction (1), each of the reagents (compound 1 and amino acid ester 2) are well known in the art with a plurality of each being commercially available.

Alternatively, the compounds of formula I can be prepared by first forming the dipeptide ester and then N-acylating these esters. That is to say that the amino acid ester or amide HNR<sup>3</sup>CR<sup>4</sup>R<sup>5</sup>C(O)Y is coupled to the N-blocked amino acid BlockNHCHR<sup>2</sup>COOH via conventional coupling conditions to provide for the dipeptide BlockNHCHR<sup>2</sup>C(O)N(R<sup>3</sup>)CR<sup>4</sup>R<sup>5</sup>C(O)Y. The blocking group is then removed via conventional conditions to provide for the free amine which is then N-acylated in the manner described above to provide for the compounds of formula I.

After coupling and N-acylation (in whatever order) is complete, the resulting esters and amides can be derivatized via conventional chemistry to provide for derivatives of the synthesized compounds. For example, conventional reduction of a terminal ester group with lithium borohydride leads to the terminal -CH<sub>2</sub>OH group. Alternatively, an ester group can be converted to a primary amide [-C(O)NH<sub>2</sub>] by reaction with ammonia in methanol with a catalytic amount of sodium cyanide while heating.

Similarly, reactive functionality which is blocked on either R<sup>2</sup> and/or R<sup>3</sup> groups can be deblocked and then derivatized. For example, the a BOC

10

15

20

25

protected amino group on R<sup>3</sup> (e.g., lysine side chain) can be deblocked after synthesis and the amino group acylated or otherwised derivatized.

Additionally, a terminal ester can be subjected to transesterification techniques to provide for other esters. Numerous techniques are known in the art to effect transesterification and each technique merely replaces one ester group with a different ester group derived from the corresponding alcohol or thioalcohol and, in some cases, a catalyst such as titanium (IV) iso-propoxide is used to facilitate reaction completion. In one technique, the alcohol or thioalcohol is first treated with sodium hydride in a suitable diluent such as toluene to form the corresponding sodium alkoxide or thioalkoxide which is then employed to effect transesterification. The efficiency of this technique makes it particularly useful with high boiling and/or expensive alcohols or thioalcohols.

In another transesterification technique, the ester to be transesterified is placed in a large excess of the alcohol or thioalcohol which effects transesterification. A catalytic amount of sodium hydride is then added and the reaction proceeds quickly under conventional conditions to provide the desired transesterified product. Because this protocol requires the use of a large excess of alcohol or thioalcohol, this procedure is particularly useful when the alcohol or thioalcohol is inexpensive.

Transesterification provides a facile means to provide for a multiplicity of different ester substituents on the compounds of formula I above. In all cases, the alcohols and thioalcohols employed to effect transesterification are well known in the art with a significant number being commercially available.

Other methods for preparing the esters of this invention include, by way of example, first hydrolyzing the ester to the free acid followed by O-alkylation with a halo-R<sup>3</sup> group in the presence of a base such as potassium carbonate.

Alternatively, for esterification procedures for alcohols containing an ester group can be achieved by using the methods of Losse, et al.<sup>11</sup>

The compounds described herein can also be prepared by use of polymer supported forms of carbodiimide peptide coupling reagents. A polymer supported form of EDC, for example, has been described (*Tetrahedron Letters*, 34(48), 7685 (1993))<sup>10</sup>. Additionally, a new carbodiimide coupling reagent, PEPC, and its corresponding polymer supported forms have been discovered and are very useful for the preparation of the compounds of the present invention.

10

15

5

WO 98/22494

Polymers suitable for use in making a polymer supported coupling reagent are either commercially available or may be prepared by methods well known to the artisan skilled in the polymer arts. A suitable polymer must possess pendant sidechains bearing moieties reactive with the terminal amine of the carbodiimide. Such reactive moieties include chloro, bromo, iodo and methanesulfonyl. Preferably, the reactive moiety is a chloromethyl group. Additionally, the polymer's backbone must be inert to both the carbodiimide and reaction conditions under which the ultimate polymer bound coupling reagents will be used.

20

25

Certain hydroxymethylated resins may be converted into chloromethylated resins useful for the preparation of polymer supported coupling reagents. Examples of these hydroxylated resins include the 4-hydroxymethyl-phenylacetamidomethyl resin (Pam Resin) and 4-benzyloxybenzyl alcohol resin (Wang Resin) available from Advanced Chemtech of Louisville, Kentucky, USA (see Advanced Chemtech 1993-1994 catalog, page 115). The hydroxymethyl groups of these resins may be converted into the desired chloromethyl groups by any of a number of methods well known to the skilled artisan.

10

Preferred resins are the chloromethylated styrene/divinylbenzene resins because of their ready commercial availability. As the name suggests, these resins are already chloromethylated and require no chemical modification prior to use. These resins are commercially known as Merrifield's resins and are available from Aldrich Chemical Company of Milwaukee, Wisconsin, USA (see Aldrich 1994-1995 catalog, page 899). Methods for the preparation of PEPC and its polymer supported forms are outlined in the following scheme.

Such methods are described more fully in U.S. Patent Application Serial No. 60/019,790 filed June 14, 1996 which application is incorporated herein by reference in its entirety. Briefly, PEPC is prepared by first reacting ethyl isocyanate with 1-(3-aminopropyl)pyrrolidine. The resulting urea is treated with 4-toluenesulfonyl chloride to provide PEPC. The polymer supported form is prepared by reaction of PEPC with an appropriate resin under standard conditions to give the desired reagent.

The carboxylic acid coupling reactions employing these reagents are performed at about ambient temperature to about 45°C, for from about 3 to 120 hours. Typically, the product may be isolated by washing the reaction with CHCl<sub>3</sub> and concentrating the remaining organics under reduced pressure. As discussed *supra*, isolation of products from reactions where a polymer bound reagent has been used is greatly simplified, requiring only filtration of the reaction mixture and then concentration of the filtrate under reduced pressure.

Still other methods for the preparation of esters are provided in the examples below.

10

15

5

Compounds where X is -CR<sup>6</sup>R<sup>6</sup>Y' are readily prepared by coupling, e.g., an amino alcohol H<sub>2</sub>NCR<sup>4</sup>R<sup>5</sup>CR<sup>6</sup>R<sup>6</sup>OH, to the carboxyl group of R<sup>1</sup>ZCX'X"C(O)NHCHR<sup>2</sup>C(O)OH under standard coupling conditions well known in peptide coupling chemistry which can use well known coupling reagents such as carbodiimides with or without the use of well known additives such as N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. If necessary, well known blocking groups on Y' can be employed to protect the group during coupling. Such blocking groups are particularly desirable when Y' is an amino group.

20

The reaction is conventionally conducted in an inert aprotic polar diluent such as dimethylformamide, dichloromethane, chloroform, acetonitrile, tetrahydrofuran and the like. Upon reaction completion, any blocking groups on Y' are selectively removed to provide for the desired compound.

25

When Y' is -OH or -SH, post-synthetic conversion of these groups to the corresponding esters (i.e., -OC(O)R<sup>7</sup>), disulfides (i.e., -SSR<sup>7</sup>) and -SSC(O)R<sup>7</sup> groups is accomplished using well known chemistry. For example, ester synthesis requires only reaction with a suitable acid such as acetic acid ( $R^7$ )

10

15

25

= methyl), acid halide (e.g., acid chloride) or acid anhydride under suitable esterification conditions.

When one of R<sup>6</sup> is hydrogen, post-synthetic oxidation of the -CHR<sup>6</sup>OH group leads to the ketone derivatives. Alternatively, such ketones can be prepared by coupling the suitable aminoketone HCl salt with the terminal carboxyl group of the amino acid as illustrated in Example 168 below.

In these synthetic methods, the starting materials can contain a chiral center (e.g., alanine) and, when a racemic starting material is employed, the resulting product is a mixture of R,S enatiomers. Alternatively, a chiral isomer of the starting material can be employed and, if the reaction protocol employed does not racemize this starting material, a chiral product is obtained. Such reaction protocols can involve inversion of the chiral center during synthesis.

Accordingly, unless otherwise indicated, the products of this invention are a mixture of R,S enatiomers or diasteriomers. Preferably, however, when a chiral product is desired, the chiral product corresponds to the L-amino acid derivative. Alternatively, chiral products can be obtained via purification techniques which separate enatiomers from a R,S mixture to provide for one or the other stereoisomer. Such techniques are well known in the art.

#### 20 Pharmaceutical Formulations

When employed as pharmaceuticals, the compounds of formula I are usually administered in the form of pharmaceutical compositions. These compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

10

15

20

25

This invention also includes pharmaceutical compositions which contain, as the active ingredient, one or more of the compounds of formula I above associated with pharmaceutically acceptable carriers. In making the compositions of this invention, the active ingredient is usually mixed with an excipient, diluted by an excipient or enclosed within such a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release

10

15

of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 to about 100 mg, more usually about 10 to about 30 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient. Preferably, the compound of formula I above is employed at no more than about 20 weight percent of the pharmaceutical composition, more preferably no more than about 15 weight percent, with the balance being pharmaceutically inert carrier(s).

The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

20

25

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type

WO 98/22494 PCT/US97/20804

described above containing from, for example, 0.1 to about 500 mg of the active ingredient of the present invention.

5

10

15

20

25

The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can separated by enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face masks tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered,

preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The following formulation examples illustrate the pharmaceutical compositions of the present invention.

5

## Formulation Example 1

Hard gelatin capsules containing the following ingredients are prepared:

	Ingredient	Quantity (mg/capsule)
10	Active Ingredient Starch Magnesium stearate	- 30.0 305.0 5.0

The above ingredients are mixed and filled into hard gelatin capsules in 340 mg quantities.

## Formulation Example 2

A tablet formula is prepared using the ingredients below:

÷	Ingredient	Quantity (mg/tablet)
	Active Ingredient Cellulose, microcrystalline	25.0 200.0
20	Colloidal silicon dioxide Stearic acid	10.0 5.0
		5.0

The components are blended and compressed to form tablets, each weighing 240 mg.

## Formulation Example 3

A dry powder inhaler formulation is prepared containing the following components:

WO 98/22494 PCT/US97/20804

-- 88 --

Ingredient	Weight %
Active Ingredient	5
Lactose	. 05

The active ingredient is mixed with the lactose and the mixture is added to a dry powder inhaling appliance.

## Formulation Example 4

Tablets, each containing 30 mg of active ingredient, are prepared as follows:

10	Ingredient	Quantity (mg/tablet)
	Active Ingredient Starch Microcrystalline cellulose Polyvinylpyrrolidone	30.0 mg 45.0 mg 35.0 mg
15	(as 10% solution in sterile water) Sodium carboxymethyl starch Magnesium stearate Talc	4.0 mg 4.5 mg 0.5 mg 1.0 mg
	Total	120 mg

The active ingredient, starch and cellulose are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The solution of polyvinyl-pyrrolidone is mixed with the resultant powders, which are then passed through a 16 mesh U.S. sieve. The granules so produced are dried at 50° to 60°C and passed through a 16 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 30 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

## Formulation Example 5

Capsules, each containing 40 mg of medicament are made as follows:

	Ingredient	Quantity (mg/capsule)	
-	Active Ingredient	40.0 mg	
	Starch	109.0 mg	
5	Magnesium stearate	1.0 mg	
	Total	150.0 mg	

The active ingredient, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 150 mg quantities.

## 10

## Formulation Example 6

Suppositories, each containing 25 mg of active ingredient are made as follows:

	Ingredient	Amount	
	Active Ingredient	25 mg	
15	Saturated fatty acid glycerides to	2,000 mg	

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2.0 g capacity and allowed to cool.

## 20

## Formulation Example 7

Suspensions, each containing 50 mg of medicament per 5.0 ml dose are made as follows:

	<u>Ingredient</u>	Amount
	Active Ingredient	50.0 mg
	Xanthan gum	4.0 mg
- +-	Sodium carboxymethyl cellulose (11%)	
5	Microcrystalline cellulose (89%)	50.0 mg
	Sucrose	1.75 g
	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.
٠	Purified water to	5.0 ml

The active ingredient, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water. The sodium benzoate, flavor, and color are diluted with some of the water and added with stirring. Sufficient water is then added to produce the required volume.

## Formulation Example 8

	Ingredient	Quantity (mg/capsule)
	Active Ingredient	15.0 mg
20	Starch	407.0 mg
	Magnesium stearate	_3.0 mg
	Total	425.0 mg

The active ingredient, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 560 mg quantities.

## Formulation Example 9

A subcutaneous formulation may be prepared as follows:

	Ingredient	Quantity	
30	Active Ingredient corn oil	5.0 mg 1 ml	

25

15

20

25

#### Formulation Example 10

## A topical formulation may be prepared as follows:

	Ingredient	Quantity	
	Active Ingredient	1-10 g	
5	Emulsifying Wax	30 g	
	Liquid Paraffin	20 g	
	White Soft Paraffin	to 100 g	

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991, herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system used for the transport of biological factors to specific anatomical regions of the body is described in U.S. Patent 5,011,472 which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion

of hydrophilic drugs into lipid-soluble drugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

Other suitable formulations for use in the present invention can be found in *Remington's Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985).

## 10 <u>Utility</u>

15

20

25

5

WO 98/22494

The compounds and pharmaceutical compositions of the invention are useful in inhibiting  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease in mammals including humans.

As noted above, the compounds described herein are suitable for use in a variety of drug delivery systems described above. Additionally, in order to enhance the *in vivo* serum half-life of the administered compound, the compounds may be encapsulated, introduced into the lumen of liposomes, prepared as a colloid, or other conventional techniques may be employed which provide an extended serum half-life of the compounds. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka, et al., U.S. Patent Nos. 4,235,871, 4,501,728 and 4,837,028 each of which is incorporated herein by reference.

The amount of compound administered to the patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions are administered to a

10

15

20

25

patient already suffering from AD in an amount sufficient to at least partially arrest further onset of the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on the judgment of the attending clinician depending upon factors such as the degree or severity of AD in the patient, the age, weight and general condition of the patient, and the like. Preferably, for use as therapeutics, the compounds described herein are administered at dosages ranging from about 1 to about 500 mg/kg/day.

In prophylactic applications, compositions are administered to a patient at risk of developing AD (determined for example by genetic screening or familial trait) in an amount sufficient to inhibit the onset of symptoms of the disease. An amount adequate to accomplish this is defined as "prophylactically effective dose." Amounts effective for this use will depend on the judgment of the attending clinician depending upon factors such as the age, weight and general condition of the patient, and the like. Preferably, for use as prophylactics, the compounds described herein are administered at dosages ranging from about 1 to about 500 mg/kg/day.

As noted above, the compounds administered to a patient are in the form of pharmaceutical compositions described above. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 and 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of WO 98/22494 PCT/US97/20804

-- 94 --

this invention. Unless otherwise stated, all temperatures are in degrees Celsius.

## **EXAMPLES**

In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted

5	meaning.		
	BOC	=	tert-butoxycarbonyl
,	BOP	=	benzotriazol-1-yloxy-tris(dimethylamino)phosphonium
			hexafluorophosphate
	bd	=	broad doublet
10	bs	=	broad singlet
	С	=	concentration (g/mL)
	CDI	==	1,1'-carbonyldiimidazole
	d	=	doublet
	dd	=	doublet of doublets
15	DCM	=	dichloromethane
	DEAD	=	diethyl azodicarboxylate
	DMF	=	dimethylformamide
	DMSO	=	dimethylsulfoxide
	EDC	=	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
20		1	hydrochloride
	EEDQ	=	2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline
	eq.	, =	equivalents
	EtOAc	=	ethyl acetate
	EtOH	=	ethanol
25	g	=	grams
	L	==	liter
	m	=	multiplet
	max	=	maximum
	MeOH	=	methanol
30	meq	=	milliequivalent
	mg	=	milligram
	mL	=	milliliter
	mm	=	millimeter
	mmol	==	millimole
35	N/A	=	not available
	N	=	normal
	ng	=	nanogram
	nm	=	nanometers
	OD	=	optical density
40	φ	==	phenyl
	PEPC	=	1-(3-(1-pyrrolidinyl)propyl)-3-ethylcarbodiimide
	psi	=	pounds per square inch
	q	=	quartet

WO 98/22494 PCT/US97/20804

-- 95 --

quintet quint. rotations per minute rpm singlet triplet trifluoroacetic acid 5 TFA tetrahydrofuran THF thin layer chromatography tlc microliter μL UV ultraviolet

15

20

25

30

In the examples below, all temperatures are in degrees Celcius (unless otherwise indicated) and each of the compounds set forth in these examples was prepared by one of the following general procedures, unless otherwise indicated.

Additionally, the term "Aldrich" indicates that the compound or reagent used in the following procedures is commercially available from Aldrich Chemical Company, Inc., 1001 West Saint Paul Avenue, Milwaukee, WI 53233 USA; the term "Fluka" indicates that the compound or reagent is commercially available from Fluka Chemical Corp., 980 South 2nd Street, Ronkonkoma NY 11779 USA; the term "Lancaster" indicates that the compound or reagent is commercially available from Lancaster Synthesis, Inc., P.O. Box 100 Windham, NH 03087 USA; the term "Sigma" indicates that the compound or reagent is commercially available from Sigma, P.O. Box 14508, St. Louis MO 63178 USA; the term "Chemservice" indicates that the compound or reagent is commercially available from Chemservice Inc., Westchester, PA; the term "Bachem" indicates that the compound or reagent is commercially available from Bachem Biosciences Inc., 3700 Horizon Drive, Renaissance at Gulph Mills, King of Prussia, PA 19406 USA; the term "Maybridge" indicates that the compound or reagent is commercially available from Maybridge Chemical Co. Trevillett, Tintagel, Cornwall PL34 OHW United Kingdom; and the term "TCI" indicates that the compound or reagent is commercially available from TCI America, 9211 North Harborgate Street, Portland OR 97203; the term "Alfa" indicates that the compound or reagent is commercially available from

10

15

20

25

Johnson Matthey Catalog Company, Inc. 30 Bond Street, Ward Hill, MA 01835-0747; the term "Novabiochem" indicates that the compound or reagent is commercially available from Calbiochem-Novabiochem Corp. 10933 North Torrey Pines Road, P.O. Box 12087, La Jolla CA 92039-2087; the term "Oakwood" indicates that the compound or reagent is commercially available from Oakwood, Columbia, South Carolina; the term "Advanced Chemtech" indicates that the compound or reagent is commercially available from Advanced Chemtech, Louisville, KY; and the term "Pfaltz & Bauer" indicates that the compound or reagent is commercially available from Pfaltz & Bauer, Waterbury, CT, USA.

The following General Procedures A'-P' and Examples A1-A74 illustrate the synthesis of N-(aryl/heteroarylacetyl)amino acid esters which can be hydrolyzed to provide for N-(aryl/heteroarylacetyl)amino acid starting materials of this invention. Other N-(aryl/heteroarylacetyl)amino acid esters can be prepared using these procedures from commercially available or known starting materials.

# GENERAL PROCEDURE A' Coupling of R<sup>1</sup>C(X')(X")C(O)Cl with H<sub>2</sub>NCH(R<sup>2</sup>)C(O)XR<sup>3</sup>

To a stirred solution of (D,L)-alanine iso-butyl ester hydrochloride (from Example B below) (4.6 mmol) in 5 mL of pyridine was added 4.6 mmol of an acid chloride. Precipitation occurred immediately. The mixture was stirred for 3.5 h, diluted with 100 mL of diethyl ether, washed with 10% HCl three times, brine once, 20% potassium carbonate once and brine once. The solution was dried over magnesium sulfate, filtered, and evaporated at reduced pressure to yield the product. Other amino acid esters may also be employed in this procedure.

10

15

20

25

## GENERAL PROCEDURE B' Coupling of R1C(X')(X")C(O)OH with H,NCH(R2)C(O)XR3

A solution of the acid (3.3 mmol) and CDI in 20 mL THF was stirred for 2 h. L-alanine iso-butyl ester hydrochloride (from Example B below) (3.6 mmol) was added, followed by 1.5 mL (10.8 mmol) of triethylamine. The reaction mixture was stirred overnight. The reaction mixture was diluted with 100 mL of diethyl ether, washed with 10% HCl three times, brine once, 20% potassium carbonate once and brine once. The solution was dried over magnesium sulfate, filtered, and evaporated at reduced pressure to yield the product. Other amino acid esters may also be employed in this procedure.

## GENERAL PROCEDURE C'

## Esterification of R¹C(X')(X")C(O)NHCH(R²)C(O)OH With HOR³

To a stirred solution of phenylacetylvaline (1.6470 g, 7.0 mmol) in 20 mL THF was added CDI (1.05 g, 6.5 mmol) and the mixture was stirred for 1.5 h. 2-Methylbutanol (0.53 g, 6 mmol) was added the mixture, followed by addition of NaH (0.16 g, 6.5 mmol). Bubbling occurred immediately. The reaction mixture was stirred overnight. The reaction mixture was diluted with 100 mL of diethyl ether, washed with 10% HCl three times, brine once, 20% potassium carbonate once and brine once. The solution was dried over magnesium sulfate, filtered, and evaporated at reduced pressure to yield the product. Other N-acyl amino acids and alcohols may also be employed in this procedure.

## GENERAL PROCEDURE D' Ester Hydrolysis to the Free Acid

Ester hydrolysis to the free acid was conducted by conventional methods. Below are two examples of such conventional de-esterification methods.

To the ester in a 1:1 mixture of CH<sub>2</sub>OH/H<sub>2</sub>O was added 2-5 equivalents of K<sub>2</sub>CO<sub>3</sub>. The mixture was heated to about 50°C for about 0.5 to 1.5 hours until tle showed complete reaction. The reaction was cooled to room temperature

10

15

20

25

and the methanol was removed at reduced pressure. The pH of the remaining aqueous solution was adjusted to about 2, and ethyl acetate was added to extract the product. The organic phase was then washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent at reduced pressure to yield the product.

The amino acid ester was dissolved in dioxane/water (4:1) to which was added LiOH ( $\sim$ 2 eq.) that was dissolved in water such that the total solvent after addition was about 2:1 dioxane:water. The reaction mixture was stirred until reaction completion and the dioxane was removed under reduced pressure. The residue was diluted with EtOAc, the layers were separated and the aqueous layer acidified to pH 2. The aqueous layer was back extracted with EtOAc, the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed under reduced pressure after filtration. The residue was purified by conventional methods (e.g., recrystallization).

The following exemplifies this later example. The methyl ester of 3-NO<sub>2</sub> phenylacetyl alanine 9.27 g (0.0348 mols) was dissolved in 60 mL dioxane and 15 mL of H<sub>2</sub>O and adding LiOH (3.06 g, 0.0731 mol) that has been dissolved in 15 mL of H<sub>2</sub>O. After stirring for 4 hours, the dioxane was removed under reduced pressure and the residue diluted with EtOAc, the layers were separated and the aqueous layer acidified to pH 2. The aqueous layer was back extracted with EtOAc (4 X 100 mL), the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed under reduced pressure after filtration. The residue was recrystallized from EtOAc/isooctane giving 7.5 g (85%) of 3-nitrophenylacetyl alanine.  $C_{11}H_{12}N_2O_5$  requires C = 52.38, H = 4.80, and N = 11.11. Analysis found C = 52.54, H = 4.85, and N = 11.08.  $[\alpha]_{23} = -29.9$  @ 589 nm.

#### GENERAL PROCEDURE E'

## Low Temperature BOP Coupling of Acid and Alcohol

A solution of methylene chloride containing the carboxylic acid (100M%) and N-methyl morpholine (150 M%) was cooled to -20°C under nitrogen. BOP (105 M%) was added in one portion and the reaction mixture was maintained at -20°C for 15 minutes. The corresponding alcohol (120 M%) was added and the reaction mixture was allowed to warm to room temperature and stirred for 12 hours. The reaction mixture was then poured into water and extracted with ethyl acetate (3x). The combined ethyl acetate portions were backwashed with saturated aqueous citric acid (2x), saturated aqueous sodium bicarbonate (2x), brine (1x), dried over anhydrous magnesium sulfate or sodium sulfate and the solvent removed under reduced pressure to yield the crude product.

## GENERAL PROCEDURE F'

#### 15

20

5

10

# EDC Coupling of Acid and Amine

The acid derivative was dissolved in methylene chloride. The amine (1 eq.), N-methylmorpholine (5 eq.), and hydroxybenzotriazole monohydrate (1.2 eq.) were added in sequence. The reaction was cooled to about 0°C and then 1.2 eq. of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride was added. The solution was allowed to stir overnight and come to room temperature under N, pressure. The reaction mix was worked up by washing the solution with saturated, aqueous Na<sub>2</sub>CO<sub>3</sub>, 0.1M citric acid, and brine before drying with Na<sub>2</sub>SO<sub>4</sub> and removal of solvents to yield crude product. Pure products were obtained by flash chromatography in an appropriate solvent.

#### 25

## GENERAL PROCEDURE G'

## EDC Coupling of Acid and Amine

A round bottom flask was charged with carboxylic acid (1.0 eq.), hydroxybenzotriazole hydrate (1.1 eq.) and amine (1.0 eq.) in THF under nitrogen atmosphere. An appropriate amount (1.1 eq. for free amines and 2.2 eq. for

hydrochloride amine salts) of base, such as Hunig's base was added to the well stirred mixture followed by EDC (1.1 eq.). After stirring from 4 to 17 hours at room temperature the solvent was removed at reduced pressure, the residue taken up in EtOAc (or similar solvent)/water. The organic layer was washed with saturated aqueous sodium bicarbonate solution, 1N HCl, brine and dried over anhydrous sodium sulfate. In some cases, the isolated product was analytically pure at this stage while, in other cases, purification via chromatography and/or recrystallization was required prior to biological evaluation.

10

15

20

5

#### GENERAL PROCEDURE H'

## Coupling of R<sup>1</sup>C(X')(X")C(O)Cl with H<sub>2</sub>NCH(R<sup>2</sup>)C(O)XR<sup>3</sup>

An excess of oxalyl chloride in dichloromethane was added to the acid derivative together with one drop of DMF. The resulting mixture was stirred for about 2 hours or until bubbling ceases. The solvent was then removed under reduced pressure and rediluted with dry methylene chloride. To the resulting solution was added about 1.1 eq. of the appropriate amino acid ester and triethylamine (1.1 eq. in methylene chloride). The system was stirred at room temperature for 2 hours and then the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate, washed with 1N HCl followed by 1N NaOH. The organic layer was dried over anhydrous soldium sulfate, filtered and the solvent removed under reduced pressure to provide for the desired product.

#### GENERAL PROCEDURE I'

## P-EPC coupling

25

P-EPC coupling employs an amino acid ester and a substituted acetic acid compound. The acetic acid derivative is well known in the art and is typically commercially available. The amino acid ester is prepared by conventional methods from the known and typically commercially available N-BOC amino acid as described in GENERAL PROCEDURE J' below.

15

20

25

Specifically, the appropriate amino ester free base (0.0346 mmols) and substituted phenylacetic acid (0.069 mmols) were dissolved in 2.0 mL CHCl<sub>3</sub> (EtOH free), treated with 150 mg of P-EPC (0.87 meq./g) and the reaction was mixed for 4 days at 23°C. The reaction was filtered through a plug of cotton, rinsed with 2.0 mL of CHCl<sub>3</sub> and the filtrate evaporated under a stream of nitrogen. The purity of each sample was determined by <sup>1</sup>H NMR and ranged from 50% to >95%. Between 8.0 and 15.0 mg of final product was obtained from each reaction and was tested without additional purification.

## GENERAL PROCEDURE J'

Synthesis of Amino Acid Esters From the Corresponding N-BOC Amino Acid

A. Esterification of the Acid.

The N-BOC amino acid was dissolved in dioxane and treated with an excess of alcohol (~1.5 eq.) and catalytic DMAP (100 mg) at 0°C. Stirring was continued until reaction completion whereupon the product was recovered by conventional methods.

## B. Removal of N-BOC Group.

The N-BOC protected amino acid was dissolved in methylene chloride (0.05M) and treated with 10 eq. of TFA at room temperature under a nitrogen atmosphere. The reaction was monitored by tlc until starting material was consumed usually within 1-5 hours. An additional 10 eq. of TFA was added to the reaction if the starting material was still present after 5 hours. The reaction was carefully neutralized with Na<sub>2</sub>CO<sub>3</sub>, separated, the organic layer washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The crude amine was then used without purification.

Specific exemplification of these procedures are as follows:

1. Racemic (+/-)-N-BOC- $\alpha$ -amino butyric acid (Aldrich) (9.29 g, 0.0457 mol) was dissolved in 100 mL of dioxane and treated with *iso*-butyl

20

alcohol (6.26 mL, 0.0686 mol), EDC (8.72 g, 0.0457) and catalytic DMAP (100 mg) at 0°C. After stirring for 17 hours, the organics were evaporated at reduced pressure, the residue diluted with EtOAc washed with NaHCO<sub>3</sub>, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporation yields 8.42 g (71%) of an oil.  $C_{13}H_{25}NO_4$  requires: C = 60.21, H = 9.72, and N = 5.40. Anal found: C = 59.91, C = 9.89, and C = 59.91, and C = 59.91, C = 9.89, and C = 59.91,

The above N-BOC amino acid ester (8.00 g, 0.032 mol) was deprotected as above giving 3.12 g (61%) of the free base as a colorless oil which solidifies upon standing.

L-N-BOC-alanine (Aldrich) (8.97 g, 0.047 mol) was dissolved in 100 mL of CH<sub>2</sub>Cl<sub>2</sub>, iso-butyl alcohol (21.9 mL, 0.238 mol) and treated with DMAP (100 mg) and EDC (10.0 g, 0.52 mol) at O°C. The mixture was stirred for 17 hours, diluted with H<sub>2</sub>O, washed with 1.0 N HCl, NaHCO<sub>3</sub>, then brine and the organics were dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and evaporation
 yields 11.8 g (quantitative) of L-N-BOC alanine iso-butyl ester which is contaminated with a small amount of solvent. A sample was vacuum dried for analytical analysis. C<sub>12</sub>H<sub>23</sub>NO<sub>4</sub> requires: C = 58.79, H = 9.38, and N = 5.71. Anal found: C = 58.73, H = 9.55, and N = 5.96.

The above N-BOC amino acid ester (11.8 g, 0.0481 mol) was deprotected as above. The free base was converted to the corresponding HCl salt using saturated HCl (g)/EtOAc to give L-N-alanine *iso*-butyl ester hydrochloride. Obtained 4.2 g (48%) of a colorless solid.  $C_7H_{15}NO_2$ . HCl requires: C = 46.28, H = 8.88, and N = 7.71. Anal found: C = 46.01, C = 46.01, C = 46.01, and C = 46.01.

#### GENERAL PROCEDURE K'

#### Methyl ester formation from amino acids

The amino acid (amino acid or amino acid hydrochloride) is suspended in methanol and chilled to 0°C. HCl gas is bubbled through this solution for 5 minutes. The reaction is allowed to warm to room temperature then stirred for 4 hours. The solvents are then removed at reduced pressure to afford the desired amino acid methyl ester hydrochloride. This product is usually used without further purification.

## Example A'

10

15

20

25

5

## Synthesis of free and polymer bound PEPC

#### N-ethyl-N'-3-(1-pyrrolidinyl)propylurea

To a solution of 27.7 g (0.39 mol) ethyl isocyanate in 250 mL chloroform was added 50 g (0.39 mol) 3-(1-pyrrolidinyl)propylamine dropwise with cooling. Once the addition was complete, the cooling bath was removed and the reaction mixture stirred at room temperature for 4 hours. The reaction mixture was then concentrated under reduced pressure to give 74.5 g (96.4%) of the desired urea as a clear oil.

## 1-(3-(1-pyrrolidinyl)propyl)-3-ethylcarbodiimide (P-EPC)

To a solution of 31.0 g (0.156 mol) N-ethyl-N'-3-(1-pyrrolidinyl)propylurea in 500 mL dichloromethane was added 62.6 g (0.62 mol) triethylamine and the solution was cooled to 0°C. To this solution were then added 59.17 g (0.31 mol) 4-toluenesulfonyl chloride in 400 mL dichloromethane dropwise at such a rate as to maintain the reaction at 0-5°C. After the addition was complete, the reaction mixture was warmed to room temperature and then heated to reflux for 4 hours. After cooling to room temperature, the reaction mixture was washed with saturated aqueous potassium carbonate (3 x 150 mL). The aqueous phases were combined and extracted with dichloromethane. All organic phases were combined and concentrated under reduced pressure. The resultant orange slurry was suspended in 250 mL diethyl ether and the solution

decanted off from the solid. The slurry/decantation process was repeated 3 more times. The ether solutions were combined and concentrated under reduced pressure to give 18.9 g (67%) of the desired product as a crude orange oil. A portion of the oil was distilled under vacuum to give a colorless oil distilling at 78-82°C (0.4 mm Hg).

# Preparation of a polymer supported form of 1-(3-(1-pyrrolidinyl)propyl)-3-ethylcarbodiimide (P-EPC)

A suspension of 8.75 g (48.3 mmol) 1-(3-(1-pyrrolidin-yl)propyl)-3-ethylcarbodiimide and 24.17 g (24.17 mmol) Merrifield's resin (2% crosslinked, 200-400 mesh, chloromethylated styrene/divinylbenzene copolymer, 1 meq. Cl/g) in dimethylformamide was heated at 100°C for 2 days. The reaction was cooled and filtered and the resulting resin washed sequentially with 1L DMF, 1L THF and 1L diethyl ether. The remaining resin was then dried under vacuum for 18 hours.

15

20

10

5

#### Example B'

#### Preparation of alanine iso-butyl ester hydrochloride

A mixture of 35.64 g (0.4 mol) of (D,L)-alanine (Aldrich) (or L-alanine (Aldrich)); 44 mL (0.6 mol) of thionyl chloride (Aldrich) and 200 mL of isobutanol was refluxed for 1.5 hours and the volatiles were removed completely on a rotavapor of 90°C under reduced pressure to give (D,L)-alanine *iso*-butyl ester hydrochloride (or L-alanine *iso*-butyl ester hydrochloride), which was pure enough to be used for further transformations.

#### Example C'

## Preparation of 3,5-dichlorophenylacetic acid

25

To a solution of 3.5 g of 3,5-dichlorobenzyl alcohol (Aldrich) in 75 mL of dichloromethane at 0°C was added 1.8 mL of methane sulfonylchloride followed by 3.5 mL of triethylamine added dropwise. After 2 hours the solution was diluted to 150 mL with dichloromethane, washed with 3N HCl,

10

15

20

saturated aqueous NaHCO<sub>3</sub> dried with Na<sub>2</sub>SO<sub>4</sub> and the solvents removed to yield the desired 3,5-dichlorobenzyl methanesulfonate as a yellow oil that was used without purification.

The crude sulfonate was dissolved in 50 mL of DMF at 0°C and then 3 g of KCN was added. After 2 hours an additional 50 mL of DMF was added and the solution was stirred for 16 hours. The red solution was diluted with 1 L of H<sub>2</sub>O and acidified to pH 3 with 3N HCl. The aqueous solution was extracted with dichloromethane. The combined organics were washed with 3N HCl, dried with Na<sub>2</sub>SO<sub>4</sub> and the solvents removed at reduced pressure to yield crude 3,5-dichlorophenylacetonitrile which was used without purification.

The nitrile was added to a mixture of 40 mL of concentrated sulfuric acid and 50 mL H<sub>2</sub>O and heated to reflux for 48 hours, cooled to room temperature and stirred for 48 hours. The reaction was diluted into 1 L of crushed ice, warmed to toom temperature and extracted with 2 x 200 mL of dichloromethane and 2 x 200 mL of ethylacetate. Both sets of organics were combined and washed with saturated aqueous NaHCO<sub>3</sub>. The NaHCO<sub>3</sub> fractions were combined and acidified to pH 1 with 3N HCl. The white solid was too fine to filter and was extracted out with 2 X 200 mL of dichloromethane. The combined organics were dried with Na<sub>2</sub>SO<sub>4</sub> and the solvents removed at reduced presure to yield crude 3,5-dichlorophenylacetic acid as a white solid. The solid was slurried with hexane and filtered to get 1.75g of white solid.

NMR (CDCl<sub>3</sub>): (in ppm) 3.61 (s, 2H), 7.19 (s,1H), 7.30 (s, 1H)

## Example D'

# Synthesis of N-(3-chlorophenylacetyl)alanine

The title compound was prepared using L-alanine (Nova Biochem) and 3-chlorophenyl acetic acid (Aldrich) by following General Procedures F' or G', followed by hydrolysis using General Procedure D'.

## Example A1

## Synthesis of N-(phenylacetyl)-D,L-alanine iso-butyl ester

Following General Procedure A' above and using phenylacetyl chloride (Aldrich) and D,L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.23-7.36$  (m, 5H), 6.18 (d, 1H), 4.58 (t, J = 7.3 Hz, 1H), 3.87 (m, 2H), 3.57 (s, 2H), 1.90 (m, 1H), 1.34 (d, J = 7.2 Hz, 3H), 0.89 (d, J = 6.8 Hz, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.7$ , 170.3, 134.5, 129.2, 128.8, 127.2, 71.3, 48.1, 43.4, 27.5, 18.8, 18.3.

 $C_{15}H_{21}NO_3$  (MW = 263.34; Mass Spectroscopy (MH<sup>+</sup> = 264))

15

20

5

10

#### Example A2

## Synthesis of N-(3-phenylpropionyl)-D, L-alanine iso-butyl ester

Following General Procedure A' above and using 3-phenylpropionyl chloride (Aldrich) and D,L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of from 51°-54°C. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.25$  (m, 2H), 7.19 (m, 3H), 6.28 (d, J = 7.2 Hz, 1H), 4.58 (quint., J = 7.2 Hz, 1H), 3.89 (m, 2H), 2.95 (t, J = 7.7 Hz, 2H), 2.50 (m, 2H), 1.92 (m, 1H), 1.33 (d, J = 7.1 Hz, 3H), 0.91 (d, J = 6.7 Hz, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.0$ , 171.5, 140.6, 128.3, 128.1, 126.0, 71.2, 47.8, 37.9, 31.4, 27.5, 18.79, 18.77, 18.3.

30  $C_{16}H_{23}NO_3$  (MW = 277.37, Mass Spectroscopy (MH<sup>+</sup> 278))

#### Example A3

#### Synthesis of N-(3-methylpentanoyl)-L-alanine iso-butyl ester

Following General Procedure B' and using 3-methylpentanoic acid (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as an oil. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.08$  (d, J = 5.9 Hz, 1H), 4.62 (quint., J = 7.3 Hz, 1H), 3.92 (m, 2H), 2.22 (m, 1H), 1.84-2.00 (m, 3H), 1.40 (d, J = 7.2 Hz, 3H), 1.35 (m, 1H), 1.20 (m, 1H), 0.85-0.96 (m, 12H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.3, 172.1, 71.4, 47.9, 43.9, 32.3, 29.38, 29.35, 27.6, 19.10, 19.06, 18.93, 18.91, 18.72, 18.67, 11.3.

 $C_{13}H_{25}NO_3$  (MW = 243.35, Mass Spectroscopy (MH<sup>+</sup> 244))

15

20

5

10

#### Example A4

#### Synthesis of N-[(4-chlorophenyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure B' and using 4-chlorophenylacetic acid (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 111°-

113°C. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.30$  (d, J = 8.2 Hz, 2H), 7.21 (d, J = 8.3 Hz, 2H), 6.18 (d, J = 5.5 Hz, 1H), 4.57 (quint., J = 7.2 Hz, 1H), 3.88 (m, 2H), 3.53 (s, 2H), 1.91 (m, 1H), 1.36 (d, J = 7.1 Hz, 3H), 0.90 (d, J = 6.8 Hz, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.8, 169.8, 133.1, 133.0, 130.6, 128.9, 71.4, 48.2, 42.6, 27.6, 18.85, 18.82, 18.4.

30  $C_{15}H_{20}NO_3Cl$  (MW = 297.78, Mass Spectroscopy (MH<sup>+</sup> 298))

15

20

#### Example A5

#### Synthesis of N-[(3,4-dichlorophenyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure B' and using 3,4-dichlorophenylacetic acid (Aldrich) and L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 81°-83°C. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

10 <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 0.90 (d, J = 6.8 Hz, 6H), 1.38 (d, J = 7.1 Hz, 3H), 1.91 (m, 1H), 3.50 (s, 2H), 3.90 (m, 2H), 4.57 (quint., J = 7.1 Hz, 1H), 6.31 (d, J = 4.9 Hz, 1H),7.12 (m, 1H), 7.38 (m, 2H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.4$ , 18.8, 18.9, 27.6, 42.2, 48.3, 71.5, 128.6, 130.6, 131.2, 131.3, 132.6, 134.7, 169.2, 172.8.

 $C_{15}H_{19}NO_3Cl_2$  (MW = 332.23, Mass Spectroscopy (MH<sup>+</sup> 332))

#### Example A6

#### Synthesis of N-[(4-methylphenyl)acetyl]-D, L-alanine iso-butyl ester

Following General Procedure B' and using 4-methylphenylacetic acid (Aldrich) and D,L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of  $102^{\circ}-104^{\circ}$ C. The reaction was monitored by tlc on silica gel (Rf =0.6 in 33% ethyl acetate/hexanes) and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

25 <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.90$  (d, J = 6.7 Hz, 6H), 1.35 (d, J = 7.2 Hz, 3H), 1.91 (m, 1H), 2.34 (s, 3H), 3.55 (s, 2H), 3.88 (m, 2H), 4.58 (m, 1H), 6.05 (bd, 1H), 7.16 (s, 4H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 18.5, 18.85, 18.87, 21.0, 27.6, 43.1, 48.1, 71.3, 129.2, 129.6, 131.3, 136.9, 170.6, 172.8.

 $C_{16}H_{23}NO_3$  (MW = 277.37, Mass Spectroscopy (MH<sup>+</sup> 278))

15

20

#### Example A7

# Synthesis of N-[(3-pyridyl)acetyl]-D,L-alanine iso-butyl ester

Following General Procedure F' and using 3-pyridylacetic acid hydrochloride (Aldrich) and D,L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 62°-64°C. The reaction was monitored by tlc on silica gel (Rf = 0.48 10% methanol/dichloromethane) and purification was by silica gel chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 8.40 (d, J = 2.8, 2H); 7.6 (m, 1H): 7.16 (m, 2H); 4.5 (quint., J = 7.2, 7.2, 1H); 3.8 (m, 2H); 3.48 (s, 2H); 1.8 (m, 1H); 1.30 (d, J = 7.2, 3H); 0.81 (d, J = 6.7, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.4, 170.1, 150.6, 148.8, 137.4, 131.4, 124.1, 71.9, 48.9, 40.6, 28.1, 19.5, 19.4, 18.6.

 $C_{14}H_{20}N_2O_3$  (MW = 264, Mass Spectroscopy (MH<sup>+</sup> 265))

#### Example A8

## Synthesis of N-[(1-naphthyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure B' and using 1-naphthylacetic acid (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 69°-73°C. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.83$  (m, 6H), 1.25 (d, J = 7.1 Hz, 3H), 1.81 (m, 2H), 3.79 (m, 2H), 4.04 (2s, 2H), 4.57 (quint., J = 7.3 Hz, 1H), 5.99 (d, J = 7.1 Hz, 1H), 7.44 (m, 2H), 7.53 (m, 2H), 7.85 (m, 2H), 7.98 (m, 1H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.2$ , 18.81, 18.83, 27.5, 41.5, 48.2, 71.3, 123.7, 125.6, 126.1, 126.6, 128.2, 128.5, 128.7, 130.7, 132.0, 133.9, 170.3, 172.5. C<sub>19</sub>H<sub>23</sub>NO<sub>3</sub> (MW = 313.40, Mass Spectroscopy (MH<sup>+</sup> 314))

#### Example A9

#### Synthesis of N-[(2-naphthyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure B' and using 2-naphthylacetic acid (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 128°-129°C. The reaction was monitored by the on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.86$  (m, 6H), 1.35 (d, J = 7.1 Hz, 3H), 1.78 (m, 1H), 3.76 (s, 2H), 3.87 (m, 2H), 4.62 (quint., J = 7.2 Hz, 1H), 6.13 (d, J = 7.1 Hz, 1H), 7.41 (m, 1H), 7.48 (m, 2H), 7.74 (s, 1H), 7.83 (m, 3H). <sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.4$ , 18.82, 18.85, 27.6, 43.7, 48.2, 71.4, 125.9, 126.3, 127.2, 127.6, 127.7, 128.2, 128.7, 132.0, 132.5, 133.5, 170.3, 172.8.  $C_{19}H_{23}NO_3$  (MW = 313.40, Mass Spectroscopy (MH<sup>+</sup> 314)).

15

20

25

5

#### Example A10

#### Synthesis of N-(4-phenylbutanoyl)-L-alanine iso-butyl ester

Following General Procedure B' and using 4-phenylbutanoic acid (Aldrich) and L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared as an oil. The reaction was monitored by tlc on silica gel and purification was by extraction with  $Et_2O$  followed by washes with aqueous  $K_2CO_3$  and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.92$  (d, J = 6.7 Hz, 6H), 1.38 (d, J = 7.1 Hz, 3H), 1.96 (m, 3H), 2.21 (t, J = 7.1 Hz, 2H), 2.64 (t, J = 7.3 Hz, 2H), 3.90 (m, 2H), 4.59 (quint., J = 7.2 Hz, 1H), 6.31 (d, 1H), 7.16 (m, 3H), 7.24 (m, 2H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.3$ , 18.75, 18.78, 26.8, 27.5, 34.9, 35.3, 47.8, 71.2, 125.7, 128.2, 128.3, 141.3, 172.1, 173.0.

 $C_{17}H_{25}NO_3$  (MW = 291.39, Mass Spectroscopy (MH<sup>+</sup> 292)).

15

20

#### Example A11

### Synthesis of N-(5-phenylpentanoyl)-L-alanine iso-butyl ester

Following General Procedure B' and using 5-phenylpentanoic acid (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as an oil. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.23$  (m, 2H), 7.17 (m, 3H), 6.30 (d, 1H), 4.59 (quint., J = 7.3 Hz, 1H), 3.91 (m, 2H), 2.61 (t, J = 7.2 Hz, 2H), 2.22 (t, J = 7.2 Hz, 2H), 1.93 (m, 1H), 1.66 (m, 4H), 1.38 (d, J = 7.2 Hz, 3H), 0.92 (d, J = 6.7 Hz, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.1, 172.3, 142.0, 128.2, 128.1, 125.6, 71.2, 47.8, 36.1, 35.5, 30.8, 27.5, 25.0, 18.80, 18.77, 18.4.

 $C_{18}H_{27}NO_3$  (MW = 305.39, Mass Spectroscopy (MH<sup>+</sup> 306)).

#### Example A12

#### Synthesis of N-[(4-pyridyl)acetyl]-D,L-alanine iso-butyl ester

Following General Procedure F' and using 4-pyridylacetic acid hydrochloride (Aldrich) and (D,L)-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 64°-66°C. The reaction was monitored by tlc on silica gel (Rf = 0.43 10% methanol/dichloromethane) and purification was by silica gel chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.51$  (dd, J = 1.6, 2.8, 1.6, 2H); 7.23 (dd, J = 4.3, 1.6, 4.4, 2H); 6.71 (d, J = 6.8, 1H); 4.56 (quint., J = 7.3, 7.2, 1H); 3.88 (m, 2H); 3.53 (s, 2H); 1.89 (m, 1H); 1.36 (d, J = 7.2, 3H); 0.88 (d, J = 6.7, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.5$ , 169.3, 150.5, 144.4, 125.1, 72.1, 48.9, 43.0, 28.2, 19.5, 19.5, 18.9.

15

20

 $C_{14}H_{20}N_2O_3$  (MW = 264, Mass Spectroscopy (MH<sup>+</sup> 265))

#### Example A13

## Synthesis of N-(phenylacetyl)-L-alanine iso-butyl ester

Following General Procedure B' and using phenylacetyl chloride (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 45°-47°C. The reaction was monitored by tlc on silica gel and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous K<sub>2</sub>CO<sub>3</sub> and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.24$ -7.39 (m, 5H), 6.14 (d, 1H), 4.58 (t, J = 7.3 Hz, 1H), 3.88 (m, 2H), 3.58 (s, 2H), 1.90 (m, 1H), 1.35 (d, J = 7.2 Hz, 3H), 0.89 (d, J = 6.7 Hz, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.8, 170.4, 134.5, 129.3, 128.9, 127.2, 71.3, 48.1, 43.5, 27.5, 18.9, 18.8, 18.4.

 $C_{15}H_{21}NO_3$  (MW = 263.34, Mass Spectroscopy (MH<sup>+</sup> 264)).

#### Example A14

# Synthesis of 2-[(3,4-dichlorophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3,4-dichlorophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure I' above) the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.36$  (m, 3H), 6.03 (bd, 1H), 4.54 (m, 1H), 3.87 (m, 2H), 3.49 (s, 2H), 1.93 (m, 2H), 1.72 (m, 1H), 0.88 (d, 6H), 0.80 (t, 3H).

#### Example A15

# Synthesis of 2-[(3-methoxyphenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3-methoxyphenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared frollowing General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

5

30

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.75$  (m, 4H), 5.93 (bd, 1H), 4.51 (m, 1H), 3.83 (m, 2H), 3.75 (s, 2H), 3.52 (s, 2H), 1.82 (m, 2H), 1.60 (m, 1H), 0.84 (d, 6H), 0.74 (t, 3H).

 $C_{17}H_{25}NO_4$  (MW = 307.39, Mass Spectroscopy (MH<sup>+</sup> 309)).

#### Example A16

# Synthesis of 2-[(4-nitrophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 4-nitrophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.16$  (d, 2H), 7.44 (d, 2H), 6.04 (bd, 1H), 4.55 (m, 1H), 3.86 (m, 2H), 3.66 (s, 2H), 1.86 (m, 2H), 1.67 (m, 1H), 0.85 (d, 6H), 0.81 (t, 3H).

 $C_{16}H_{22}N_2O_5$  (MW = 322.36, Mass Spectroscopy (MH<sup>+</sup> 323)).

Example A17

# Synthesis of 2-[(3,4-methylenedioxyphenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3,4-(methylenedioxy)-phenyl acetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction

was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.72$  (m, 3H), 5.92 (bd, 1H), 4.54 (m, 1H), 3.86 (m, 2H), 3.66 (s, 2H), 1.86 (m, 2H), 1.66 (m, 1H), 0.89 (d, 6H), 0.79 (t, 3H).

#### Example A18

## Synthesis of 2-[(thien-3-yl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3-thiopheneacetic acid

(Aldrich) and iso-butyl 2-aminobutyrate (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

#### Example A19

# Synthesis of 2-[(4-chlorophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 4-chlorophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

25

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.22$  (m, 2H), 7.11 (m, 2H), 5.80 (m, 1H), 4.44 (m, 1H), 3.78 (m, 2H), 3.43 (s, 2H), 1.77 (m, 2H), 1.56 (m, 1H), 0.83 (d, 6H) 0.71 (t, 3H).

15

#### Example A20

# Synthesis of 2-[(3-nitrophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3-nitrophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.15$  (m, 2H), 7.65 (m, 1H), 6.08 (m, 1H), 4.46 (m, 1H), 3.92 (m, 2H), 3.68 (s, 2H), 1.91 (m, 2H), 1.75 (m, 1H), 0.98 (d, 6H) 0.71 (t, 3H).

#### Example A21

# Synthesis of 2-[(2-hydroxyphenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 2-hydroxyphenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.14$  (m, 1H), 7.01 (m, 1H), 6.93 (m, 1H), 6.79 (m, 1H), 6.46 (m, 1H), 4.51 (m, 1H), 3.87 (m, 2H), 3.57 (s, 2H), 2.01 (m, 2H), 1.75 (m, 1H), 0.89 (d, 6H), 0.85 (t, 3H).

#### Example A22

# Synthesis of 2-[(2-naphthyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 2-naphthylacetic acid (Aldrich) and iso-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.83$  (m, 7H), 5.95 (m, 1H), 4.58 (m, 1H), 3.84 (m, 2H), 3.75 (s, 2H), 1.89 (m, 2H), 1.63 (m, 1H), 0.91 (d, 6H), 0.81 (t, 3H).  $C_{20}H_{25}NO_3$  (MW = 327.42, Mass Spectroscopy (MH<sup>+</sup> 328)).

5

10

15

20

#### Example A23

## Synthesis of 2-[(2,4-dichlorophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 2,4-dichlorophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.49$  (m, 1H), 7.22 (m, 2H) 5.98 (m, 1H), 4.52 (m, 1H), 3.86 (m, 2H), 3.61 (s, 2H), 1.84 (m, 2H), 1.62 (m, 1H) 0.87 (d, 6H), 0.80 (t, 3H).

#### Example A24

## Synthesis of 2-[(4-bromophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 4-bromophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tle on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.43$  (d, 2H), 7.19 (d, 2H) 5.85 (m, 1H), 4.51 (m, 2H), 3.81 (m, 2H), 3.47 (s, 2H), 1.84 (m, 2H), 1.61 (m, 1H) 0.84 (d, 6H), 0.76 (t, 3H).

 $C_{16}H_{22}NO_3Br$  (MW = 356.26, Mass Spectroscopy (MH<sup>+</sup> 358)).

#### Example A25

# Synthesis of 2-[(3-chlorophenyl)acetamido])butyric acid iso-butyl ester

Following General Procedure I' above and using 3-chlorophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.25$  (m, 3H), 7.12 (m, 1H) 5.80 (m, 1H), 4.52 (m, 1H), 3.86 (m, 2H), 3.50 (s, 2H), 1.87 (m, 2H), 1.67 (m, 1H) 0.88 (d, 6H), 0.77 (t, 3H).

 $C_{16}H_{22}NO_3Cl$  (MW = 311.81 Mass Spectroscopy (MH+ 313)).

#### Example A26

# Synthesis of 2-[(3-fluorophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3-fluorophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.31$  (m, 1H), 7.01 (m, 3H) 5.95 (m, 1H), 4.54 (m, 1H), 3.84 (m, 2H), 3.54 (s, 2H), 1.88 (m, 2H), 1.65 (m, 1H) 0.87 (d, 6H), 0.81 (t, 3H).

 $C_{16}H_{22}NO_3F$  (MW = 295.35 Mass Spectroscopy (MH<sup>+</sup> 296)).

25

#### Example A27

Synthesis of 2-[(benzothiazol-4-yl)acetamido]butyric acid iso-butyl ester Following General Procedure I' above and using 4-benzothiazol-4-yl acetic acid (Chemservice) and iso-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was

monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.82$  (m, 1H), 7.51-7.21 (m, 4H) 5.84 (m, 1H), 4.51 (m, 1H), 3.90 (s, 2H), 3.79 (m, 2H), 1.78 (m, 2H), 1.58 (m, 1H) 0.80 (d, 6H), 0.66 (t, 3H).

#### Example A28

Synthesis of 2-[(2-methylphenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 2-methylphenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

15  $^{1}$ H-nmr (CDCl<sub>3</sub>):  $\delta = 7.18$  (m, 4H), 5.79 (m, 1H), 4.54 (m, 1H), 3.85 (m, 2H), 3.59 (s, 2H), 3.29 (s, 3H), 1.81 (m, 2H), 1.59 (m, 1H) 0.87 (d, 6H), 0.77 (t, 3H).

 $C_{17}H_{25}NO_3$  (MW = 291.39 Mass Spectroscopy (M<sup>+</sup> 291)).

### Example A29

Synthesis of 2-[(2-fluorophenyl)acetamido]butyric acid iso-butyl ester
Following General Procedure I' above and using 2-fluorophenylacetic acid
(Aldrich) and iso-butyl 2-aminobutyrate (prepared following General Procedure
J' above), the title compound was prepared. The reaction was monitored by tlc
on silica gel and purification was by filtration as described in the general
procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.28$  (m, 1H), 7.09 (m, 3H) 6.03 (m, 1H), 4.54 (m, 1H), 3.87 (m, 2H), 3.57 (s, 2H), 1.89 (m, 2H), 1.64 (m, 1H) 0.88 (d, 6H), 0.80 (t, 3H).

25

30

#### Example A30

## Synthesis of 2-[(4-fluorophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 4-fluorophenylacetic acid (Aldrich) and iso-butyl 2-aminobutyrate (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.20$  (m, 2H), 6.97 (m, 2H) 5.87 (m, 1H), 4.492 (m, 1H), 3.83 (m, 2H), 3.48 (s, 2H), 1.86 (m, 2H), 1.60 (m, 1H) 0.87 (d, 6H), 0.78 (t, 3H).

 $C_{16}H_{22}NO_3F$  (MW = 295.35 Mass Spectroscopy (MH<sup>+</sup> 296)).

#### Example A31

## Synthesis of 2-[(3-bromophenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3-bromophenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.45$  (m, 2H), 7.23 (m, 2H) 5.95 (m, 1H), 4.55 (m, 1H) 3.84 (m, 2H) 3.55 (s, 2H), 1.89 (m, 2H), 1.68 (m, 1H) 0.91 (d, 6H), 0.81 (t, 3H).

 $C_{16}H_{22}NO_3Br$  (MW = 356.26 Mass Spectroscopy (M<sup>+</sup> 357)).

#### Example A32

# Synthesis of 2-[(3-trifluoromethylphenyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 3-trifluoromethylphenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction

was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.52$  (m, 1H), 7.47 (m, 2H) 6.01 (m, 1H), 4.56 (m, 1H), 3.86 (m, 2H), 3.61 (s, 2H), 1.84 (m, 2H), 1.62 (m, 1H) 0.87 (d, 6H), 0.80 (t, 3H).

 $C_{17}H_{22}NO_3F_3$  (MW = 345.36 Mass Spectroscopy (MH<sup>+</sup> 345)).

#### Example A33

## Synthesis of 2-[(2-thienyl)acetamido]butyric acid iso-butyl ester

Following General Procedure I' above and using 2-thiopheneacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.89$  (m, 3H), 6.07 (bd, 1H), 4.50 (m, 1H), 3.82 (m, 2H), 3.71 (s, 2H), 1.85 (m, 2H), 1.62 (m, 1H), 0.81 (d, 6H), 0.75 (t, 3H).

 $C_{14}H_{21}NO_3S$  (MW = 283.39, Mass Spectroscopy (MH<sup>+</sup> 284)).

20

25

#### Example A34

# Synthesis of 2-(phenylacetamido)butyric acid iso-butyl ester

Following General Procedure H' above and using phenylacetic acid (Aldrich) and *iso*-butyl 2-aminobutyrate (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by chromatography on silica gel using 9:1 toluene:EtOAc as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.17-7.28$  (m, 5H), 6.23 (bd, 1H), 4.51 (m, 1H), 3.86 (m, 2H), 3.54 (s, 2H), 1.87 (m, 2H), 1.62 (m, 1H), 0.87 (d, 6H), 0.78 (t, 3H).

 $C_{16}H_{23}NO_3$  (MW = 277.36, Mass Spectroscopy (MH<sup>+</sup> 277)).

5

10

15

20

25

#### Example A35

# Synthesis of N-(phenylacetyl)valine 2-methylbutyl ester

Step A. Preparation of N-(phenylacetyl) valine

To a stirred solution of 5.15 g (44 mmol) of valine (Bachem) in 50 mL (100 mmol) of 2N NaOH cooled to 0°C was added dropwise 5.3 mL (40 mmol) of phenylacetyl chloride (Aldrich). A colorless oil precipitated. The reaction mixture was allowed to warm to room temperature and stirred for 18 hours, washed with 50 mL diethyl ether, acidified to pH 2-3 with aqueous HCl. The white precipitate formed was filtered off, washed thoroughly with water, followed by diethyl ether to give 7.1 g (30 mmol, 69% yield) of the title compound.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta$  = 12.63 (s, 1H), 8.25 (d, J = 8.6 Hz, 1H), 7.27 (m, 5H), 4.15 (m, 1H), 3.56 (d, J = 13.8 Hz, 1H), 3.47 (d, J = 13.8 Hz, 1H), 2.05 (m, 1H), 0.87 (d, J = 6.8, Hz, 3H), 0.84 (d, J = 6.8 Hz, 3) <sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta$  = 173.2, 170.4, 136.6, 129.0, 128.2, 126.3, 57.1, 41.9, 30.0, 19.2, 18.0 C<sub>13</sub>H<sub>17</sub>NO<sub>3</sub> (MW=235.29; Mass Spectroscopy (MH<sup>+</sup> = 236))

Step B. Synthesis of N-(phenylacetyl)valine 2-methylbutyl ester

Following General Procedure C' and using the N-(phenylacetyl) valine prepared in Step A above and 2-methylbutan-1-ol (Aldrich), the title compound was prepared as a diastereomeric mixture. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

20

<sup>1</sup>H-nmr (CDCI<sub>3</sub>):  $\delta = 7.25$ -7.40 (m, 5H), 5.95 (d, 1H), 4.56 (m, 1H), 3.84-4.00 (m, 2H), 3.61 (s, 2H), 2.10 (m, 1H), 1.68 (m, 1H), 1.38 (m, 1H), 1.15 (m 1H), 0.82-0.94 (m, 9H), 0.76 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 171.84, 171.81, 170.7, 134.6, 129.31, 129.27, 128.9, 127.3, 69.8, 57.0, 43.7, 33.9, 31.3, 25.9, 25.8, 18.9, 17.4, 16.34, 16.27, 11.12, 11.07.

 $C_{18}H_{27}NO_3$  (MW = 305.42, Mass Spectroscopy (MH 306)).

#### Example A36

#### Synthesis of N-(phenylacetyl)-L-methionine iso-butyl ester

L-Methionine (0.129g, 0.869 mmols) (Aldrich) was taken-up in dioxane (5.0 mL) and treated with a saturated solution of sodium bicarbonate (5.0 mL) followed by phenylacetyl chloride (Aldrich) (0.114 mL, 0.822 mmols). After stirring for 17 hours at room temperature the mixture was diluted with ethyl acetate, the layers separated and the aqueous layer acidified to pH 2 with 5N HCl. The crude product was extracted into ethyl acetate, dried over sodium sulfate, vacuum dried and used without further purification.

N-phenylacetyl-L-methionine (0.1285 g, 0.447 mmol) was dissolved in 3.0 mL dioxane and *iso*-butyl alcohol (0.2 mL) and treated with EDC (0.094 g, 0.492 mmol), and catalytic DMAP (0.015g). After stirring for 17 hours at 23°C, the mixture was evaporated at reduced pressure to an oil, the residue was diluted in EtOAc and washed with 0.1 N HCl and saturated sodium bicarbonate. Chromatography on silica gel using 98:2 CHCl<sub>3</sub>/MeOH as eluant provided the pure product.

NMR data was as follows:

 $C_{17}H_{25}NO_3S$  (MW = 323.17, Mass Spectroscopy (M<sup>+</sup> 323)

15

### Example A37

## Synthesis of N-(phenylacetyl)-L-leucine iso-butyl ester

L-Leucine (Aldrich) (0.114g, 0.869 mmols) was taken-up in dioxane (5.0 mL) and treated with a saturated solution of sodium bicarbonate (5.0 mL) followed by phenylacetyl chloride (Aldrich) (0.114 mL, 0.822 mmols). After stirring for 17 hours at room temperature the mixture was diluted with ethyl acetate, the layers separated and the aqueous layer acidified to pH 2 with 5N HCl. The crude product was extracted into ethyl acetate, dried over sodium sulfate, vacuum dried and used without further purification.

N-Phenylacetyl-L-leucine (0.0081 g, 0.038 mmol) was dissolved in 2.0 mL CHCl<sub>3</sub> (EtOH free) and *iso*-butyl alcohol (0.055 mL) and treated with P-EPC (100 mg, 0.87 milliequivalents). The mixture was rotated for 4 days, filtered through a plug of cotton and the filtrate evaporated at reduced pressure to an oil which was sufficiently pure for testing.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.22$  (m, 5H), 5.57 (d, 1H), 4.35 (m, 1H), 3.35 (m, 3H), 1.35 (m, 4H), 0.68 (m, 9H).

 $C_{18}H_{27}NO_3$  (MW = 305.40, Mass Spectroscopy (M<sup>+</sup> 305)).

#### Example A38

Synthesis of N-[(3-chlorophenyl)acetyl]alanine 3-methylbut-2-enyl ester

Following General Procedure C' above and using N-(3-chlorophenylacetyl
alanine (from Example D' above) and 3-methylbut-2-en-1-ol (Aldrich), the title
compound can be prepared. The reaction was monitored by tlc on silica gel
and purification was by liquid chromatography using 30% EtOAc/hexane as the
eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.39-7.16$  (m, 4H), 6.06 (bd, 1H), 5.38-5.29 (m, 1H), 4.63 (d, J = 9Hz, 2H), 3.56 (s, 2H), 1.79 (s, 3H), 1.7 (s, 3H), 1.39 (d, J = 9Hz, 3H).

15

#### Example A39

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine cyclopropylmethyl ester

Following General Procedure C' above, and using N-(3-chlorophenylacetyl alanine (from Example D' above) and cyclopropylmethanol (Aldrich), the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.2$ -7.1 (m, 4H), 6.09 (bs, 1H), 4.6 (dq, J = 9 Hz, 1H), 3.96 (dd, J = 9Hz, 2H), 3.59 (s, 2H), 1.2 (d, J = 9Hz, 3H), 1.2-1.0 (m, 1H), 0.603-0.503 (m, 2H), 0.300-0.203 (m, 2H).

#### Example A40

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine 2-thienylmethyl ester

Following General Procedure C' above, and using N-(3-chlorophenylacetyl alanine (from Example D' above) and 2-thiophenemethanol (Aldrich) the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.37$ -6.97 (m, 7H), 5.97 (q, J = 14 Hz, 2H), 4.6 (dq, J = 9 Hz, 1H), 3.76 (s, 2H), 1.38 (d, J = 9Hz, 3H).

#### Example A41

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine (1-methylcyclopropyl)methyl ester

25 Following General Procedure C' above, and using N-(3-chlorophenylacetyl alanine (from Example D' above) and (1-methylcyclopropyl)methanol (Aldrich) the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.6$  (bd, J = 9 Hz, 1H), 3.86 (q, J = 14 Hz, 2H), 3.4 (s, 2H), 2.29 (q, J = 9 Hz, 1H), 1.3 (d, J = 9Hz, 3H), 1.03 (s, 3H), 0.5-0.4 (m, 2H), 0.4-0.28 (m, 2H).

5

10

15

20

#### Example A42

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine 3-thienylmethyl ester

Following General Procedure C' above, and using N-(3-chlorophenylacetyl alanine (from Example D' above) and 3-thiophenemethanol (Aldrich) the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.03$  (bd, J = 9 Hz, 1H), 7.56-7.5 (m, 1H), 7.47 (bs, 1H), 7.4-7.17 (m, 4H), 7.06 (d, J = 9 Hz, 1H), 5.1 (s, 2H), 4.3 (dq, 1H), 1.3 (d, J = 9 Hz, 3H).

#### Example A43

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine 2-methylcyclopentyl ester

Following General Procedure C' above, and using N-(3-chlorophenylacetylalanine (from Example D' above) and 2-methylcyclopentanol (Aldrich) the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.39-7.16$  (m, 4H), 6.3 (bd, 1H), 4.79-4.7 (m, 1H), 4.6-4.25 (m, J = 9 Hz, 1H), 3.577 (s, 2H), 2.09-1.8 (m, 2H), 1.74-1.6 (m, 2H), 1.39 (dd, J = 9 Hz, 3H), 1.2 (dt, J = 9 Hz, 1H), 0.979 (dd, J = 9 Hz, 2H)

 $C_{17}H_{22}NO_3Cl$  (MW = 323.82, Mass Spectroscopy (MH<sup>+</sup> 323).

15

eluant.

#### Example A44

## Synthesis of N-[(3-chl rophenyl)acetyl] alanine 2-methylprop-2-enyl ester

Following General Procedure C' above, and using N-(3-chlorophenylacetyl alanine (from Example D' above) and 2-methylprop-2-en-1-ol (Aldrich) the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.39-7.16$  (m, 4H), 6.03 (bs, 1H), 4.77 (s, 2H), 4.7-10 4.29 (m, 3H), 2.59 (s, 2H), 1.73 (s, 3H), 1.43 (d, J = 9 Hz, 3H)  $C_{15}H_{18}NO_3Cl$  (MW = 295.76, Mass Spectroscopy (MH<sup>+</sup> 295)).

#### Example A45

Synthesis of N-[(3-chlorophenyl)acetyl]alanine cyclohex-2-enyl ester

Following General Procedure C' above, and using N-(3-chlorophenylacetyl
alanine (from Example D' above) and cyclohex-2-en-1-ol (Aldrich) the title
compound can be prepared. The reaction was monitored by tlc on silica gel
and purification was by liquid chromatography using 3:7 EtOAc:hexane as the

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.6$  (bd, J = 9 Hz, 1H), 7.4-7.2 (m, 4H), 6.0-5.8 (m, 1H), 5.7-5.5 (m, 1H), 5.1 (bs, 1H), 4.13-4.29 (m, 1H), 3.5 (s, 2H), 2.1-1.9 (m, 2H), 1.8-1.69 (m, 1H), 1.69-1.49 (m, 4H), 1.3 (dd, J = 9 Hz, 3H)  $C_{17}H_{20}NO_3Cl$  (MW = 321.8, Mass Spectroscopy (MH<sup>+</sup> 321.2)).

#### Example A46

Synthesis of N-[(2-phenylbenzoxazol-5-yl)acetyl]alanine iso-butyl ester
Following General Procedure I' above, and using 5-(2-phenylbenzoxazol)yl-acetic acid (CAS# 62143-69-5) and alanine iso-butyl ester (prepared
following General Procedure I' above), the title compound was prepared.

10

20

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.24$  (m, 3H), 7.68 (m, 1H), 7.51 (m, 5H), 6.04 (m, 1H), 4.58 (m, 1H), 3.85 (m, 2H), 3.68 (s, 2H), 1.9 (m, 1H), 1.35 (d, 3H), 0.87 (d, 6H).

 $C_{22}H_{24}N_2O_4$  (MW = 380, Mass Spectroscopy (MH<sup>+</sup> 381)).

#### Example 47

#### Synthesis of N-[(3-methylthiophenyl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using 3-methylthiophenylacetic acid (CAS# 18698-73-2) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.14$  (m, 2H), 7.01 (m, 1H), 4.56 (m, 1H), 3.88 (m, 2H), 3.54 (s, 2H), 2.46 (s, 3H), 1.89 (m, 1H), 1:35 (d, 3H) 0.85 (d, 6H).  $C_{16}H_{23}NO_{3}S$  (MW = 309, Mass Spectroscopy (MH<sup>+</sup> 310)).

#### Example A48

#### Synthesis of N-4-[(2-furyl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using 2-furylacetic acid (CAS# 2745-26-8) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.36$  (m, 1H), 6.34 (m, 1H), 6.21 (m, 1H), 4.56 (m, 1H), 3.91 (m, 2H), 3.61 (s, 2H), 1.92 (m, 1H), 1.38 (d, 3H) 0.89 (d, 6H).  $C_{13}H_{19}NO_4$  (MW = 253, Mass Spectroscopy (MH<sup>+</sup> 254)).

25

#### Example A49

Synthesis of N-[(benzofuran-2-yl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using benzofuran-2-ylacetic acid (Maybridge) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.51$  (m, 1H), 7.44 (m, 1H), 7.25 (m, 2H), 6.67 (s, 1H), 4.60 (m, 1H), 3.87 (m, 2H), 3.77 (s, 2H), 1.88 (m, 1H), 1.38 (d, 3H), 0.87 (d, 6H).

 $C_{17}H_{21}NO_4$  (MW = 303, Mass Spectroscopy (MH<sup>+</sup> 304)).

#### Example A50

# Synthesis of N-[(benzothiophen-3-yl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using thianaphthen-3-ylacetic acid (Lancaster) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.89$  (m, 1H), 7.76 (m, 1H), 7.38 (m, 3H), 6.07 (m, 1H), 4.57 (m, 1H), 3.92 (m, 2H), 3.82 (s, 4H), 1.84 (m, 1H), 1.32 (d, 3H) 0.85 (d, 6H).

 $C_{17}H_{21}NO_3S$  (MW = 319, Mass Spectroscopy (MH<sup>+</sup> 320)).

Example A51

Synthesis of N-[(2-chloro-5-thienyl)acetyl]alanine iso-butyl ester
Following General Procedure I' above, and using 5-chloro-2-thienyl)acetic
acid (CAS# 13669-19-7) and alanine iso-butyl ester (prepared following General

Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.77$  (m, 1H), 6.68 (d, 1H), 6.31 (bm, 1H), 4.59 (m, 1H), 3.91 (m, 2H), 3.38 (s, 2H), 1.90 (m, 1H), 1.39 (d, 3H) 0.89 (d, 6H).  $C_{13}H_{18}NO_3SCl$  (MW = 303, Mass Spectroscopy (MH<sup>+</sup> 303)).

#### Example A52

Synthesis of N-[(3-methylisoxazol-5-yl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using (3-methyl-isoxazol-5-yl)acetic acid (CAS# 19668-85-0) and alanine iso-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by the on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.07$  (s, 2H), 4.56 (m, 1H), 3.92 (m, 2H), 3.68 (s, 2H), 2.29 (s, 3H), 1.94 (m, 1H), 1.89 (d, 3H) 0.91 (d, 6H).  $C_{13}H_{20}N_2O_4$  (MW = 268, Mass Spectroscopy (MH<sup>+</sup> 269)).

#### Example A53

Synthesis of N-[(2-phenylthiothienyl)acetyl]alanine iso-butyl ester
Following General Procedure I' above, and using (2-phenylthiothienyl)acetic acid and alanine iso-butyl ester (prepared following General
Procedure J' above), the title compound was prepared. The reaction was
monitored by tlc on silica gel and purification was by filtration as described in
the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.21-7.11$  (m, 6H), 6.92 (d, 1H), 4.56(m, 1H), 3.87 (m, 2H), 3.72 (s, 2H), 1.94 (m, 1H), 1.38 (d, 3H) 0.89 (d, 6H).  $C_{19}H_{23}NO_3S_2$  (MW = 377, Mass Spectroscopy (MH<sup>+</sup> 378)).

#### Example A54

#### Synthesis of N-[(6-methoxybenzothiophen-2-yl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using (6-methoxythianaphthen-2-yl)acetic acid and alanine iso-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.59$  (d, 1H), 7.33 (d, 1H), 7.16 (s, 1H), 7.03 (dd, 1H), 4.56 (m, 1H), 3.87(s, 3H), 3.84 (m, 2H), 3.76 (s, 2H), 1.85 (m, 1H), 1.30 (d, 3H) 0.86 (d, 6H).

 $C_{18}H_{23}NO_4S$  (MW = 349, Mass Spectroscopy (MH<sup>+</sup> 350)).

#### Example A55

#### Synthesis of N-[(3-phenyl-1,2,4-thiadiazol-5-yl)acetyl]alanine iso-butyl ester

15 Following General Procedure I' above, and using (3-phenyl-1,2,4-thiadiazol-5-yl)acetic acid (CAS# 90771-06-5) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

20 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.47$  (m, 5H), 4.66 (m, 1H), 4.16 (s, 2H), 3.91 (m, 2H), 1.93 (m, 1H), 1.48 (d, 3H) 0.93 (d, 6H).

 $C_{17}H_{21}N_3O_3S$  (MW = 347, Mass Spectroscopy (MH<sup>+</sup> 348)).

#### Example A56

Synthesis of N-[2-phenyloxazol-4-yl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using (2-phenyloxazol-4yl)acetic acid (CAS# 22086-89-1) and alanine iso-butyl ester (prepared
following General Procedure J' above), the title compound was prepared. The

reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

#### Example A57

5 Synthesis of N-[(3-methylphenyl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using 3-methylphenylacetic acid (Aldrich) and alanine *iso*-butyl ester (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.21$  (m, 1H), 7.07 (m, 3H), 4.54 (m, 1H), 3.83 (m, 2H), 3.52 (s, 2H), 2.35 (s, 3H), 1.87 (m, 1H), 1.32 (d, 3H), 0.88 (d, 6H).  $C_{16}H_{23}NO_3$  (MW = 277, Mass Spectroscopy (MH<sup>+</sup> 278)).

15

20

10

#### Example A58

Synthesis of N-[(2,5-difluorophenyl)acetyl]alanine iso-butyl ester
Following General Procedure I' above, and using 2,5-difluorophenylacetic
acid (Aldrich) and alanine iso-butyl ester (prepared following General
Procedure J' above), the title compound was prepared. The reaction was
monitored by tlc on silica gel and purification was by filtration as described in
the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.08$ -6.94 (m, 3H), 4.57 (m, 1H), 3.91 (m, 2H), 3.56 (s, 2H), 1.92 (m, 1H), 1.41 (d, 3H) 0.91 (d, 6H).

25  $C_{15}H_{19}NO_3F_2$  (MW = 299, Mass Spectroscopy (MH<sup>+</sup> 300)).

30

#### Example A59

# Synthesis of N-[(3,5-diflurophenyl)acetyl]alanine iso-butyl ester

Following General Procedure I' above, and using 3,5-difluorophenylacetic acid (Aldrich) and alanine *iso*-butyl ester (prepared following General Procedure I' above), the title compound was prepared. The reaction was monitored by the on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.81$  (m, 2H), 6.74 (m, 1H), 6.06 (m, 1H), 4.57 (m, 1H), 3.92 (m, 2H), 3.51 (s, 2H), 1.94 (m, 1H), 1.36 (d, 3H) 0.87 (d, 6H).  $C_{15}H_{19}NO_3F_2$  (MW = 299, Mass Spectroscopy (MH<sup>+</sup> 300)).

#### Example A60

## Synthesis of N-[(3-thienyl)acetyl]alanine iso-butyl ester

15 Following General Procedure I' above, and using 3-thiopheneacetic acid (Aldrich) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.33$  (m, 1H), 7.14 (m, 1H), 7.01 (m, 1H), 6.09 (m, 1H), 4.58 (m, 1H), 3.88 (m, 2H), 3.60 (s, 2H), 1.91 (m, 1H), 1.37 (d, 3H) 0.92 (d, 6H).

Optical Rotation:  $[\alpha]_{23}$  -52 (c 1 MeOH) @ 589 nm.

25  $C_{13}H_{19}NO_3S$  (MW = 269, Mass Spectroscopy (MH<sup>+</sup> 269)).

#### Example A61

# Synthesis of N-[(4-methylphenyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure I' above, and using 4-methylphenylacetic acid (Aldrich) and L-alanine iso-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc

15

20

25

on silica gel and purification was by filtration as described in the general procedure.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.11$  (s, 4H), 5.93 (m, 1H), 4.58 (m, 1H), 3.88 (m, 2H), 3.54 (s, 2H), 2.33 (s, 3H), 1.89 (m, 1H), 1.32 (d, 3H), 0.89 (d, 6H).  $C_{16}H_{23}NO_3$  (MW = 277.35, Mass Spectroscopy (MH<sup>+</sup> 278)).

#### Example A62

# Synthesis of N-(phenylacetyl)-L-alanine S-1-(methoxycarbonyl) iso-butyl ester

Following General Procedure K' and using (S)-(+)-2-hydroxy-2-methylbutyric acid (Aldrich) in place of the amino acid, methyl (S)-(+)-2-hydroxy-2-methylbutyrate was prepared.

Methyl (S)-(+)-2-hydroxy-2-methylbutyrate was then coupled with carbobenzyloxy-L-alanine (Aldrich) using General Procedure E' to provide carbobenzyloxy-L-alanine S-1-(methoxycarbonyl) iso-butyl ester.

Carbobenzyloxy-L-alanine S-1-(methoxycarbonyl) iso-butyl ester (1.0 g) was then dissolved in 20 mL of methanol and 6N HCl (0.5 mL) and 10% palladium on carbon (0.1 g) were added. This reaction mixture was hydrogenated at 40 psi of hydrogen on a Parr apparatus for 5 hours at room temperature and then filtered through a pad of Celite. The filtrate was concentrated at reduced pressure to provide L-alanine S-1-(methoxycarbonyl) iso-butyl ester hydrochloride (98% yield).

L-Alanine S-1-(methoxycarbonyl) iso-butyl ester hydrochloride was then coupled to phenylacetic acid using General Procedure G' to provide the title compound.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.35 - 7.20$  (m, 5H), 6.22 (bd, 1H), 4.83 (d, 1H), 4.65 (p, 1H), 3.68 (s, 3H), 3.55 (s, 2H), 2.21 (m, 1H), 1.40 (d, 3H), 0.97 (d, 3H), 0.93 (d, 3H).

30  $^{13}$ C-nmr (CDCl<sub>3</sub>):  $\delta = 173.25$ , 171.18, 170.22, 135.11, 129.94, 129.50, 127.88, 52.67, 48.49, 43.98, 30.53, 19.21, 18.75, 17.58.

10

#### Example A63

## Synthesis of N-[(3-nitrophenyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure H' above and using 3-nitrophenylacetic acid (Aldrich) and L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared. The reaction was monitored by tlc on silica gel and purification was by recrystallization from butyl chloride.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.17$  (m, 2H), 7.68 (d, 1H), 7.52 (t, 1H), 6.18 (m, 1H), 4.48 (m, 1H), 3.94 (m, 2H), 3.67 (s, 2H), 1.93 (m, 1H), 1.42 (d, 3H), 0.91 (d, 3H).

Optical Rotation:  $[\alpha]_{23}$  -49 (c 5, MeOH).

#### Example A64

# Synthesis of N-[(3,5-difluorophenyl)acetyl]alanine ethyl ester

Following General Procedure G' and using 3,5-difluorophenylacetic acid (Aldrich) and alanine ethyl ester (Aldrich), the title compound was prepared as a solid with a melting point of 93°-95°C. The reaction was monitored by tlc on silica gel (Rf = 0.8 in EtOAC) and purification was by chromatography on silica gel using EtOAc as the eluant followed by recrystallization from 1-chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.30$  (d, 3H); 3.52 (s, 2H).  $C_{13}H_{15}NO_3F_2$  (MW = 271.26, Mass Spectroscopy (MH<sup>+</sup> 271)).

#### Example A65

Synthesis of N-[(3-nitrophenyl)acetyl] methionine ethyl ester

Following General Procedure G' above and using 3-nitrophenylacetic acid (Aldrich) and methionine ethyl ester hydrochloride (Aldrich), the title compound was prepared. The reaction was monitored by the on silica gel and purification was by recrystallization from butyl chloride.

30 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.18$  (s, 1H), 8.15 (d, 1H) 7.66 (d, 1H), 7.48 (t, 1H), 6.30 (m, 1H), 4.67 (m, 1H), 4.21 (t, 2H), 3.67 (s, 2H), 2.47 (t, 2H), 2.12 (m, 2 H), 2.08 (s, 3H), 1.27 (t, 3H).

Optical Rotation:  $[\alpha]_{23}$  -30 (c 5, MeOH).

5

10

20

#### Example A66

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine iso-butyl ester

Following General Procedure G' above and using 3-chlorophenylacetic acid (Aldrich) and alanine *iso*-butyl ester (prepared following General Procedure J' above), the title compound was prepared. The reaction was monitored by tlc on silica gel.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.29$  (m, 3H), 7.18 (m, 1H), 6.0 (m, 1H), 4.56 (m, 1H), 3.89 (m, 2H), 3.53 (s, 2H), 1.91 (m, 1H), 1.39 (d, 3 H), 0.91 (d, 3H). Optical Rotation:  $[\alpha]_{23}$  -45 (c 5, MeOH).

15  $C_{15}H_{20}NO_3Cl$  (MW = 297.78, Mass Spectroscopy (MH<sup>+</sup> 297)).

#### Example A67

# Synthesis of N-[(3-chlorophenyl)acetyl]alanine 2-(N,N-dimethylamino)ethyl ester

Following General Procedure C' above, and using N-(3-chlorophenyl-acetyl)alanine (from Example D' above) and 2-(N,N-dimethyl amino) ethanol (Aldrich), the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 0.1:2:0.79 NH<sub>4</sub>OH:EtOH:CHCl<sub>3</sub> as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): 7.37 (s, 1H), 7.33-7.2 (m, 3H), 4.675-4.6 (m, 1H), 4.5-4.37 (m, 1H), 4.25-4.13 (m, 1H), 3.6 (d, J = 7 Hz, 2H), 2.86 (bs, 2H), 2.3 (s, 6H), 1.23 (d, J = 9 Hz, 3H).

 $C_{15}H_{21}N_2O_3Cl$  (MW = 313.799, Mass Spectroscopy (M<sup>+</sup> 313)).

20

25

#### Example A68

# Synthesis of 2-[(3,5-dichlorophenyl)acetamido]hexanoic acid methyl ester

Following General Procedure F' above, an using 3,5-dichlorophenylacetic acid (from Example C' above) and L-norleucine methyl ester hydrochloride (Bachem), the title compound was prepared as a solid having a melting point of 77°-78°C. The reaction was monitored by tlc on silica gel (Rf = 0.70 in 40% EtOAC/hexanes) and purification was by flash chromatography on silica gel using 40% EtOAc/hexanes as the eluant.

NMR data was as follows:

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.54$ , 169.67, 138.43, 135.72, 128.33, 128.07, 78.04, 77.62, 77.19, 53.04, 52.90, 43.14, 32.57, 27.87, 22.81, 14.41.

#### Example A69

15 Synthesis of N-[(3,5-diclorophenyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure F' above, and using 3,5-dichlorophenylacetic acid (from Example C' above) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of  $115^{\circ}-116^{\circ}$ C. The reaction was monitored by tlc on silica gel (Rf = 0.40 in 3% methanol/dichloromethane) and purification was by flash chromatography on silica gel using 3% methanol/dichloromethane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.27$  (d, J = 2 Hz, 1H), 7.19 (s, 2H), 6.22 (d, J = 6 Hz, 1H), 4.59 (quint., J = 7 Hz, 1H), 3.9 (q, J = 4 Hz, 2H), 3.5 (s, 2H), 1.9 (m, 1H), 1.4 (d, J = 7 Hz, 3H), 0.91 (d, J = 7 Hz, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.45, 169.37, 138.31, 135.75, 128.39, 128.11, 78.04, 77.61, 77.19, 72.19, 54.03, 48.97, 43.12, 28.24, 19.52, 19.49, 19.09. C<sub>15</sub>H<sub>19</sub>NO<sub>3</sub>Cl<sub>2</sub> (MW = 331.9, Mass Spectroscopy (MH<sup>+</sup> 332)).

20

#### Example A70

## Synthesis of N-(cyclohexylacetyl)-L-alanine iso-butyl ester

Following General Procedure B' above, and using cyclohexylacetic acid (Aldrich) and L-alanine iso-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of 92°C-93°C. The reaction was monitored by tlc on silica gel (Rf = 0.39 in 1:3 EtOAc:hexane) and purification was by extraction with  $Et_2O$  followed by washes with aqueous  $K_2CO_3$  and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 0.93 (d, J = 6.7 Hz, 6H), 0.85-1.01 (m, 2H), 1.05-1.35 (m, 3H), 1.40 (d, J = 7.1 Hz, 3H), 1.60-1.85 (m, 6H), 1.95 (m, 1H), 2.06 (d, J = 7.0 Hz, 2H), 3.92 (m, 2H), 4.61 (m, 1H), 6.08 (bd, 1H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 18.7, 18.9, 26.0, 26.1, 27.6, 33.0, 35.3, 44.6, 47.9, 71.4, 171.8, 173.3.

15  $C_{15}H_{27}NO_3$  (MW = 269.39, Mass Spectroscopy (MH<sup>+</sup> 270)).

#### Example A71

## Synthesis of N-(cyclopentylacetyl)-L-alanine iso-butyl ester

Following General Procedure B' above, and using cyclopentylacetic acid (Aldrich) and L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of  $62^{\circ}$ C- $64^{\circ}$ C. The reaction was monitored by tlc on silica gel (Rf = 0.37 in 1:3 EtOAc:hexane) and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous  $K_2CO_3$  and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.87$  (d, J = 6.8 Hz, 6H), 1.01-1.17 (m, 2H), 1.34 (d, J = 7.2 Hz, 3H), 1.40-1.62 (m, 4H), 1.70-1.83 (m, 2H), 1.89 (m, 1H), 2.15 (m, 3H), 3.86 (m, 2H), 4.55 (m, 1H), 6.30 (d, J = 7.1 Hz, 1H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.4$ , 18.78, 18.80, 24.8 (very high), 27.5, 32.27, 32.32, 36.9, 42.5, 47.7, 71.2, 172.2, 173.2.

10

20

Elemental Analysis-Calc (%): C, 65.85; H, 9.87; N, 5.49; Found (%): C, 66.01; H, 10.08; N, 5.49.

 $C_{14}H_{25}NO_3$  (MW = 255.36, Mass Spectroscopy (MH<sup>+</sup> 256)).

#### Example A72

Synthesis of N-[(cyclohex-1-enyl)acetyl]-L-alanine iso-butyl ester

Following General Procedure B' above, and using cyclohex-1-enyl acetic acid (Alfa) and L-alanine *iso*-butyl ester hydrochloride (from Example B' above), the title compound was prepared as a solid having a melting point of  $49^{\circ}\text{C-}51^{\circ}\text{C}$ . The reaction was monitored by tlc on silica gel (Rf = 0.40 in 1:3 EtOAc:hexane) and purification was by extraction with Et<sub>2</sub>O followed by washes with aqueous  $K_2\text{CO}_3$  and aqueous HCl.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCI<sub>3</sub>):  $\delta = 0.91$  (d, J = 4.5 Hz, 3H), 0.93 (d, J = 6.7 Hz, 3H), 1.40 (d, J = 7.2 Hz, 3H), 1.52-1.70 (m, 4H), 1.97 (m, 3H), 2.06 (bs,

2H), 2.89 (s, 2H), 3.92 (m, 2H), 4.59 (m, 1H), 5.65 (s, 1H), 6.33 (d, J = 6.6 Hz, 1H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.7$ , 18.91, 18.93, 21.9, 22.7, 25.3, 27.6, 28.3, 46.1, 47.9, 71.4, 127.1, 132.5, 170.6, 173.1.

Elemental Analysis-Calc (%): C, 67.38; H, 9.42; N, 5.24; Found (%): C, 67.34; H, 9.54; N, 5.16.

 $C_{15}H_{25}NO_3$  (MW = 267.37, Mass Spectroscopy (MH<sup>+</sup> 268)).

#### Example A73

Synthesis of N-[(3-chlorophenyl)acetyl]alanine 3-methylbut-2-enyl thioester

Following General Procedure C' above, and using N-f(3-

chlorophenyl)acetyl] alanine and 3-methyl-2-butene thioester (TCI), the title compound can be prepared. The reaction was monitored by tlc on silica gel and purification was by liquid chromatography using 3:7 EtOAc:Hexane as the eluant.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 5.2-5.075$  (m, 1H), 4.37 (dq, J = 9 Hz, 1H), 3.56 (s), 3.43 (d, J = 12 Hz, 2H), 1.266 (d, J = 12 Hz, 6H) 1.3 (d, J = 9 Hz, 3H).

 $C_{16}H_{20}NO_2CIS$  (MW = 325.86, Mass Spectroscopy (M<sup>+</sup> 325)).

5

#### Example A74

### Synthesis of N-[(2-phenyl)-2-fluoroacetyl]alanine ethyl ester

Following General Procedure F' above, and using  $\alpha$ -fluorophenyl acetic acid (Aldrich) and alanine ethyl ester (Aldrich), the title compound was prepared. The reaction was monitored by tlc on silica gel (Rf = 0.75 in 1:1 EtOAc:hexane) and purification was by chromatography on silica gel using 1:2 ethyl acetate/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.14$  (q, 3H), 1.34 (d, 3H), 4.07 (m, 2H), 4.33 (m, 1H), 5.84 (d, 1H), 6.01 (d, 1H), 7.40-7.55 (m, 5H), 8.87 (m, 1H). C<sub>13</sub>H<sub>16</sub>NO<sub>3</sub>F (MW = 253.27, Mass Spectroscopy (MH<sup>+</sup> 253)).

15

20

10

# Example A75

## Synthesis of N-(3,5-difluorophenylacetyl)-L-phenylglycine methyl ester

Following General Procedure F above, and using 3,5-difluorophenylacetic acid (Aldrich) and L-phenylglycine methyl ester hydrochloride (Bachem), the title compound was prepared.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  =7.4-7.3 (m, 5H), 6.9-6.7 (m, 3H), 6.55 (d 1H, 7.1 Hz), 5.56 (d 1H 7 Hz), 3.72 (s 3H), 3.57 (s 2H)

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 197.6$ , 177.6, 171.8, 169.3, 136.7, 129.6, 129.3, 127.8, 113.0, 112.9, 112.7, 111.4, 103.8, 103.5, 65.1, 57.2, 53.5, 45.1, 43.3, 43.3

 $C_{17}H_{15}NO_3F_2$  (MW = 319.31, Mass Spectroscopy (MH +320)).

#### Example 76

10

15

Synthesis of N-(3,5-difluorophenylacetyl)-L-phenylglycine iso-butyl ester
The 3,5-difluorophenylacetic acid (Aldrich) was EDC coupled to Lphenylglycine methyl ester hydrochloride (Bachem) via General Procedure F
above.

The resulting compound was placed in a large excess of the desired alcohol. A catalytic amount of dry NaH was added, and the reaction was followed by tlc until the presence of starting material was no longer detected. The reaction was quenched with a few milliliters of 1N HCl, and after a few minutes of stirring saturated aqueous NaHCO, was added. The volume of the reaction mixture was reduced on a rotary evaporator until the excess alcohol was removed and then the remaining residue was taken up in ethyl acetate and additional water was added. The organic phase was washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator, and the crude product residue was then further purified by chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.35-7.3$  (m 5H), 6.8-6.7 (m 3H) 6.60 (d 1H, 7 Hz), 5.55 (d 1H 7.1 Hz), 3.9 (m 2H), 3.60 (s 2H), 1.85 (m 1H 7 Hz), 0.8 (q 6H 7 Hz)

20  $^{13}$ C-nmr (CDCl<sub>3</sub>):  $\delta = 171.3$ , 169.3, 165.4, 138.5, 137.0, 129.5, 129.2, 127.6, 113.1, 113.0, 112.8, 112.7, 103.8, 103.5, 103.2, 75.5, 57.2, 43.4, 43.3, 28.2, 19.3

 $C_{20}H_{21}NO_3F_2$  (MW = 361.39, Mass Spectroscopy (MH +362)).

#### Example A77

Synthesis of N-(cyclopentylacetyl)-L-phenylglycine methyl ester

Following General Procedure D' above, and using cyclopentylacetic acid (Aldrich) with L-phenylglycine methyl ester hydrochloride (Bachem) the title compound was prepared.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.35$  (s, 5H), 6.44 (bd, 1H), 5.6 (d, 1H), 3.72 (s, 3H), 2.24 (bs, 3H), 1.9-1.4 (m, 6H), 1.2-1.05 (m, 2H) <sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.3$ , 171.7, 136.7, 129.0, 128.6, 127.3, 56.2,

52.7, 42.5, 36.9, 32.40, 32.38, 24.8

5

10

25

#### Example A78

# Synthesis of N-(cyclopentylacetyl)-L-alanine methyl ester

Following General Procedure D' above, and using cyclopentylacetic acid (Aldrich) with L-alanine methyl ester hydrochloride (Sigma) the title compound was prepared.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.38$  (d, 1H), 4.50 (m, 1H), 3.65 (s, 3H), 2.13 (bs, 3H), 1.80-1.00 (m (includes d at 1.30, 3H), 11H)

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.7$ , 172.5, 52.1, 47.6, 42.3, 36.8, 32.15, 32.14,

15 18.0

 $C_{11}H_{19}NO_3$  (MW = 213.28, Mass Spectroscopy (MH<sup>+</sup> 214)).

#### Example A79

# Synthesis of N-(cyclopropylacetyl)-L-phenylglycine methyl ester

Following General Procedure D' above, and using cyclopropylacetic acid

(Aldrich) with L-phenylglycine methyl ester hydrochloride (Bachem), the title compound was prepared.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.35$  (m, 5H) 6.97 (bd, J = 7.2 Hz, 1H) 5.59 (d, J = 7.8 Hz, 1H), 3.71 (s, 3H), 2.17 (m, 2H), 1.05-0.95 (m, 1H), 0.62 (m, 2H), 0.02 (m, 2H)

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 171.9, 174.6, 136.6, 129.0, 128.5, 127.2, 56.1, 52.7, 41.0, 6.9, 4.37, 4.33

10

15

25

#### Example A80

## Synthesis of N-(cyclopropylacetyl)-L-alanine methyl ester

Following General Procedure D' above, and using cyclopropylacetic acid (Aldrich) with L-alanine methyl ester hydrochloride (Sigma), the title compound was prepared.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.60$  (d, 1H), 4.55 (m, 1H), 3.69 (s, 3H), 2.10 (m, 2H), 1.34 (d, 3H), 0.95 (m, 1H), 0.58 (m, 2H) 0.15 (m, 2H)

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.7$ , 172.3, 52.3, 47.7, 41.0, 18.2, 6.7, 4.27, 4.22

#### Example A81

#### Synthesis of N-[(3-nitrophenyl)acetyl]-L-methionine iso-butyl ester

Following General Procedure H' above, and using nitrophenylacetic acid (Aldrich) and L-methionine (Aldrich), the title compound was prepared as a tan oil. The reaction was monitored by tlc on silica gel.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.16$  (m,2H) 7.67 (d,1H) 7.32 (t, 1H), 6.31 (bd, 1H), 4.69 (m, 1H), 3.90 (d, 2H), 3.68 (s, 2H), 2.47 (t, 2H), 2.15 (m, 1H), 2.02 (s, 3H), 1.90 (m, 2H), 0.91 (d, 6H).

20  $C_{17}H_{24}N_2O_5S$  (MW = 368.4, Mass Spectroscopy (MH<sup>+</sup> 368)).

The following General Procedures A"-B" and Examples B1-B2 illustrate the synthesis of N-(aryl/heteroarylacetyl)amino acid starting materials useful in this invention. Other N-(aryl/heteroarylacetyl)amino acids can be prepared using these procedures from commercially available or known starting materials.

## GENERAL PROCEDURE A"

## Acid Chloride Preparation

3,5-Difluorophenylacetic acid (30 g, 0.174 mol) (Aldrich) was dissolved in dichloromethane and this solution was cooled to 0°C. DMF (0.5 mL, catalytic) was added followed by the dropwise addition of oxalyl chloride (18 mL, 0.20 mol) over a 5 minute period. The reaction was stirred for 3 h and then rotoevaporated at reduced pressure to a residue which was placed on a high vacuum pump for 1 h to afford 3,5-difluorophenylacetyl chloride as a thin yellow oil. Other acid chlorides can be prepared in a similar manner.

10

15

20

5

## GENERAL PROCEDURE B"

## Schotten-Bauman Procedure

3,5-Difluorophenylacetyl chloride (from General Procedure A") was added dropwise to a 0°C solution of L-alanine (Aldrich) (16.7 g, 0.187 mol) in 2 N sodium hydroxide (215 mL, 0.43 mol). The reaction was stirred for 1 h at 0°C and then overnight at room temperature. The reaction was diluted with water (100 mL), then extracted with ethyl acetate (3 x 150 mL). The organic layer was then washed with brine (200 mL), dried over MgSO<sub>4</sub>, and rotoevaporated at reduced pressure to a residue. Recrystallization of the residue from ethyl acetate/hexanes afforded the desired product (34.5 g, 82% yield). Other acid chlorides may be used in this procedure to provide for intermediates useful in this invention.

## Example B1

## Synthesis of N-(Phenylacetyl)-L-alanine

Following General Procedure B" above, title compound was prepared from phenylacetyl chloride (Aldrich) and L-alanine (Aldrich) as a solid having a melting point of 102-104°C.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 9.14 (br s, 1H), 7.21-7.40 (m, 5H), 6.20 (d, J = 7.0 Hz, 1H), 4.55 (m, 1H), 3.61 (s, 2H), 1.37 (d, J = 7.1 Hz, 3H). <sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 176.0, 171.8, 134.0, 129.4, 127.5, 48.3, 43.2, 17.9.

5

## Example B2

## Synthesis of N-(3,5-Difluorophenylacetyl)-L-alanine

Following General Procedure B" above, the title compound was prepared from 3,5-difluorophenylacetyl chloride (from General Procedure A" above) and L-alanine (Aldrich).

10

15

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta$  = 8.32 (br s, 0.3H), 6.71 (m, 2H), 6.60 (m, 1H), 4.74 (br s, 1.7H), 4.16 (m, 1H), 3.36 (s, 2H), 1.19 (d, J = 7.3 Hz, 3H). <sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta$  = 175.9, 172.4, 164.4 (dd, J = 13.0, 245.3 Hz), 141.1, 113.1 (dd, J = 7.8, 17.1 Hz), 102.9 (t, J = 25.7 Hz), 49.5, 42.7, 17.5.

The following General Procedures A"'-C" and Examples C1-C8 illustrate the synthesis of dipeptide ester starting materials useful in this invention. Other dipeptide esters can be prepared using these procedures from commercially available or known starting materials.

20

25

## GENERAL PROCEDURE A"

## **EDC** Coupling Procedure

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at 0°C or room temperature was charged with THF, carboxylic acid (1.0 eq), an amine or amine hydrochloride (1.1 eq.), 1-hydroxybenzotriazole hydrate (1.15-1.2 eq.), N,N-diisopropylethylamine (2.2-2.9 eq.), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.15-1.2 eq.). The cooling bath was removed and the mixture allowed to warm to room temperature with stirring for 10-20 hours. The mixture was diluted with EtOAc

10

15

20

25

and washed with 0.5 N aqueous HCl (2x), dilute aqueous NaHCO<sub>3</sub> (1x), brine (1x) and dried over either Na<sub>2</sub>SO<sub>4</sub> or MgSO<sub>4</sub>. The drying agent was removed by filtration and the filtrate concentrated *in vacuo*. The residue was either used without further purification or purified using standard procedures, such as flash chromatography on silica gel and/or recrystallization.

# GENERAL PROCEDURE B" Removal of the N-tert-Boc Protecting Group

The *N-tert*-Boc-amine was dissolved in a suitable dry solvent (such as 1,4-dioxane or ethyl acetate) and the solution was cooled in an ice bath. Gaseous HCl was introduced into the solution until the mixture was saturated with HCl. The mixture was then stirred until the reaction was complete. The resulting mixture was concentrated under reduced pressure to yield the amine hydrochloride. The amine hydrochloride was used without purification or was triturated using, for example, diethyl ether and the resulting solid was collected by filtration.

## GENERAL PROCEDURE C'' EEDO Coupling Procedure

A round bottom flask containing a magnetic stir bar under as atmosphere of nitrogen at room temperature was charged with THF, a carboxylic acid (1 eq.), an amine hydrochloride (1.1 eq.), and 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ) (1.1 eq). The reaction mixture was allowed to stir for 15 minutes and then 4-methylmorpholine (1.1 eq) was added and stirring was continued at room temperature for 15-20 hours. The reaction mixture was concentrated in vacuo and the resulting residue was partitioned between ethyl acetate and water. The organic phase was separated and washed with saturated aqueous NH<sub>4</sub>Cl (2x), saturated aqueous NaHCO<sub>3</sub> (2x), followed by brine (1x). The organic phase was then dried over Na<sub>2</sub>SO<sub>4</sub> and the drying agent was removed by filtration and the filtrate concentrated in vacuo. The residue was

10

15

either used without further purification or purified using standard procedures, such as flash chromatography on silica gel and/or recrystallization.

## Example C1

## Synthesis of N-(L-Methionine)-L-phenylglycine Methl Ester Hydrochloride

Following General Procedure A" and using N-(tert-butoxycarbonyl)-L-methionine (Sigma)-and L-phenylglycine methyl ester hydrochloride (Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B" to afford the title compound as a crude solid or foam.

## Example C2

## $Synthesis \ of \\ N-(2-Aminobutanoyl)-L-phenylglycine \ Methl \ Ester \ Hydrochloride$

Following General Procedure A" and using N-(tert-butoxycarbonyl)-2-aminobutyric acid (Sigma) and L-phenylglycine methyl ester hydrochloride (Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B" to afford the title compound as a crude solid or foam.

## Example C3

20

25

## Synthesis of N-(L-Leucine)-L-phenylglycine Methl Ester Hydrochloride

Following General Procedure A" and using N-(tert-butoxycarbonyl)-L-leucine (Sigma) and L-phenylglycine methyl ester hydrochloride (Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B" to afford the title compound as a crude solid or foam.

## Example C4

## Synthesis of N-(L-Phenylalanine)-L-phenylglycine Methl Ester Hydrochloride

Following General Procedure A'' and using N-(tert-butoxycarbonyl)-L-phenylalanine (Sigma) and L-phenylglycine methyl ester hydrochloride (Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B''' to afford the title compound as a crude solid or foam.

## Example C5

10

15

5

## Synthesis of N-(Glycine)-L-phenylglycine Methl Ester Hydrochloride

Following General Procedure A" and using N-(tert-butoxycarbonyl)glycine (Sigma) and L-phenylglycine methyl ester hydrochloride (Bachem), the Bocprotected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B" to afford the title compound as a crude solid or foam.

## Example C6

## Synthesis of N-(L-Phenylglycine)-L-phenylglycine Methl Ester Hydrochloride

Following General Procedure C" and using N-(tert-butoxycarbonyl)-Lphenylalanine (Sigma) and L-phenylglycine methyl ester hydrochloride
(Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam.
The resulting crude dipeptide was deprotected using General Procedure B" to
afford the title compound as a crude solid or foam.

25

## Example C7

Synthesis of N-(L-Valine)-L-phenylglycine Methl Ester Hydrochloride

10

Following General Procedure A" and using N-(tert-butoxycarbonyl)-L-valine (Sigma) and L-phenylglycine methyl ester hydrochloride (Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B" to afford the title compound as a crude solid or foam.

## Example C8

## Synthesis of

## N-[(S)-2-Aminocyclohexylacetyl)-L-phenylglycine Methl Ester Hydrochloride

Following General Procedure A" and using N-(tert-butoxycarbonyl)-(S)-aminocyclohexylacetic acid (e.g., Boc-L-cyclohexylglycine) and L-phenylglycine methyl ester hydrochloride (Bachem), the Boc-protected dipeptide was prepared as a crude solid or foam. The resulting crude dipeptide was deprotected using General Procedure B" to afford the title compound as a crude solid or foam.

The following Examples D1-D4 illustrate the synthesis of various intermediates useful as starting materials for this invention. Similar intermediates can be prepared using these procedures and commercially available or known starting materials.

## Example D1

20

25

## Synthesis of 3,5-Difluorophenyl- $\alpha$ -fluoroacetic Acid

Methyl 3,5-difluoromandelate was prepared following General Procedure G below and using commmerically available 3,5-difluoromandelic acid. The resultant α-hydroxy methyl ester was fluorinated according to the general procedure described in W. J. Middleton, et al., *Org. Synth. Col. Vol. VI*, 835. Specifically, a solution of diethylaminosulfur trifluoride (1.1 eq) in CH<sub>2</sub>Cl<sub>2</sub> was cooled to 0°C and treated with methyl 3,5-difluoromandelate (1.0 eq) as a solution in CH<sub>2</sub>Cl<sub>2</sub>. After 10 min. the cooling bath was removed and the

10

15

reaction was stirred at ambient temperature for 30 min. The reaction was monitored by tlc (Rf = 0.65, 1:1 ethyl acetate/hexanes). The mixture was then poured onto ice and the layers separated. The organic phase was washed with saturated aqueous NaHCO3 and brine. The organic layer was dried over Na2SO4, filtered, and concentrated in vacuo. The product was purified by LC2000 chromatograpy (180 mL/min) using 10% EtoAc/hexanes as the eluent. The resulting methyl 3,5-difluorophenyl- $\alpha$ -fluoroacetate was hydrolyzed by dissolving the ester in 70% aqueous dioxane and treating with lithium hydroxide (2.0 eq.). No starting material remained by tlc after 2 h. The dioxane was removed via rotary evaporation. The aqueous mixture was first washed with ethyl acetate and then acidified with 0.01 N HCl. The aqueous layer was extracted with ethyl acetate. The organic phase was washed with brine, dried over Na2SO4, filtered, and concentrated. The crude solid was recrystallized from ethyl acetate/hexanes affording 3,5-difluorophenyl- $\alpha$ -fluoroacetic acid as a white solid having a melting point of 90-110°C.

 $C_8H_5F_3O_2$  (MW = 190.1); mass spectroscopy: 190.1.

## Example D2

## Synthesis of (S)-2-Hydroxy-2-methyl-1-phenylprop-1-ylamine

(S)-2-Hydroxy-2-methyl-1-phenylprop-1-ylamine was prepared by adding
 5.0 equivalents of methyl magnesium bromide to a solution of L-phenylglycine methyl ester hydrochloride in THF at 0 C. The reaction mixture was stirred for 1 hour and then quenched with sodium bicarbonate. After standard work-up conditions, the residue was purified by silica gel chromatography using 10 %
 MeOH/CHCl<sub>3</sub> as the eluent.

### Example D3

Synthesis of Methyl (S)-2-Amino-2-(6-methoxy-2-naphthyl)acetate

(S)-2-(tert-Butoxycarbonylamino)-2-(6-methoxy-2-naphthyl)acetic acid was prepared from 2-(6-methoxy-2-naphthyl)acetic acid according to the general method described by D.A. Evans, et al., J. Amer. Chem. Soc., (1990), 112, 4011-4030. Briefly, (S)-3-(6-methoxy-2-naphthylacetyl)-4-benzyl-2ozazolidinone was converted to (S)-3-[(S)-6-methoxy-2-naphthyl- $\alpha$ -azidoacetyl)-5 4-benzyl-2-ozazolidinone via standard enolate azidation procedures using potassium 1,1,1,3,3,3-hexamethyldisilazane and trimethylsilyl azide at -78°C. Treatment of the azide derivative with 3 equivalents of lithium hydroxide in THF then provided (S)-2-azido-2-(6-methoxy-2-naphthyl)acetic acid. Reduction of this intermediate, as its sodium salt, in 1:1 1,4-dioxane/water (0.05 M) with 10 1 atm of hydrogen, 10% Pd/C at 25°C afforded (S)-2-azido-2-(6-methoxy-2naphthyl)acetic acid, which was then converted, without isolation, to its N-Boc derivative on treatment with 1.4 equivalents of di-tert-butyl dicarbonate and 0.47 equivalents of sodium carbonate. The product was isolated by the acidification to pH 2 with 1 N NaHSO<sub>4</sub> and extraction with three portions of 15 ethyl acetate. The product was recrystallized from ethyl acetate/hexanes to afford a white solid, m.p. = 176°C (shrink); 197-199°C (dec).

NMR data was as follows:

25

30

<sup>1</sup>HMR (DMSO- $d_6$ ):  $\delta = 12.78$  (s, 1H), 7.84-7.77 (m, 3H), 7.62 (d, J=8 Hz, 1H), 7.49 (d, J=8 Hz, 1H), 7.31 (d, J=2 Hz, 1H), 7.17 (dd, J=9, 2 Hz, 1H), 5.22 (d, J=8 Hz, 1H), 3.87 (s, 3H), 1.39 (s, 9H).

(S)-2-(tert-Butoxycarbonylamino)-2-(6-methoxy-2-naphthyl)acetic acid was then converted into the methyl ester using General Procedure G below. The methyl ester was then dissolved in CH<sub>2</sub>Cl<sub>2</sub> and this solution cooled to 0°C. Trifluoroacetic acid (50 molar eq.) was added and the reaction was allowed to warm to room temperature and stirring was continued for 2 hrs. The reaction mixture was then concentrated and the residue extracted into CH<sub>2</sub>Cl<sub>2</sub> and washed with sodium bicarbonate solution. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to yield methyl (S)-2-amino-2-(6-methoxy-2-naphthyl)acetate.

10

15

20

## Example D4

## Synthesis of Methyl 2-Amino-2-(thieno[2,3-b]thiophen-2-yl)acetate

To a 3.75 mole equivalents of sodium hydride (oil free) was added DMF and the resulting mixture was cooled to 0°C. A solution of methyl thieno[2,3-b]thiophen-2-carboxylate (1 mole eq.) and methyl methylsulfinyl methyl sulfide (1.1 mole eq.) in DMF was then added dropwise and the reaction mixture was stirred at 0°C for 30 min and then allowed to warm to room temperature and stirring was continued for 3 h. The reaction was then quenched with methanol and the product extracted into EtOAc. The organic extracts were washed with water followed by brine, and then dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give a gummy brown oil. The residue was slurried in diethyl ether and the resulting solid collected. The solid was then dissolved in hot ethyl acetate and decolorizing carbon was added. The mixture was then filtered and solvent removed to give a solid, which was used without further purification.

Acetic anhydride (10 mole eq.) and acetic acid (1.8 mole eq.) were mixed together and heated to 70°C for 15 min. and then cooled to 65°C. The solid sulfone from above was added in portions and the reaction was allowed to stir at 70°C for 30 min. and then cooled and concentrated. The resulting solid was taken up in ethyl acetate and washed with sodium bicarbonate solution, followed by 1 N Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution. The solution was then dried over MgSO<sub>4</sub>, filtered and concentrated to give methyl 2-keto-2-(thieno[2,3-b]thiophen-2-yl)thioacetate as a solid, which was used without further purification.

To the 2-keto compound (0.0165 moles) (4.0g) was added 270 mL of methanol and 16.5 mL of 1 N NaOH. The reaction was allowed to stir for 6 h at room temperature and then methoxyamine (1.38 g, 0.0165 moles) was added and stirring was continued for 18 h. The reaction mixture was then concentrated and the residue dissolved in ethyl acetate and washed with water. The aqueous layer was then acidified with in HCl and the oily product was

extracted into ethyl acetate and washed with brine. The organic layer was dried over MgSO<sub>4</sub>, filtered and concentrated to give 4.0 g of 2-(hydroxyimino)-2-(thieno[2,3-b]thiophen-2-yl)acetic acid as a yellow solid.

The methyl ester was then prepared using General Procedure G below and the oxime was reduced to an amino group using General Procedure R below to afford methyl 2-amino-2-(thieno[2,3-b]thiophen-2-yl)acetate.

## Example D5

## Synthesis of N-Methyl-N'-BOC-Leucinamide

A solution of 0.9968 g (4 mmol) of N-BOC-leucine (Bachem) and 1.2323 g (7.6 mmol) of CDI in 40 mL of THF was stirred for 1 hour, and then 0.5402 g (8 mmol) of methylamine hydrochloride (Aldrich) and 0.8092 g (8 mmol) of N-methylmorpholine were added. The mixture was stirred for 16 hours, evaporated at reduced pressure to dryness, and the residue was washed thproughly with water, 1N NaOH, water, followed by diethyl ether to yield 0.886 g (3.09 mmol, 70%) of the title compound.

## Example D6

## Synthesis of N-BOC-Norleucine amide

To a stirred mixture of 3.47 g (15 mmol) of BOC-norleucine (Bachem),

3.44 g (22.5 mmol) of 1-hydroxybenzotriazole monohydrate and 50 mL of dichloromethane at 0°C was added 3.45 g (1.2 mmol) of EDC. The resulting mixture was stirred at 0°C for 1 hour and then ammonia gas was bubbled through the mixture for 10 min. The cooling bath was allowed to warm to room temperature and the mixture stirred for 18 hours. The mixture was evaporated at reduced pressure to dryness, triturated with 20% Na<sub>2</sub>CO<sub>3</sub>. The resulting solid was collected by filtration and washed with water to yield 2.69 g (11.7 mmol, 78%) of the title compound.

## Example D7

## $Synthesis \ of \\ N-[N-(3,5-Difluor ophenylacetyl)-L-alaninyl]-L-alanine$

The title compound was prepared by dissolving 1.98 g (0.006 mols) of N
[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester (from Example 85 below) in 60 mL dioxane and 15 mL of H<sub>2</sub>O and adding LiOH (0.25 g, 0.006 mol) that has been dissolved in 15 mL of H<sub>2</sub>O. After stirring for 3 hours, the dioxane was removed under reduced pressure and the residue diluted with EtOAc, the layers were separated and the aqueous layer acidified to pH 2. The aqueous layer was back extracted with EtOAc (4 X 100 ml), and the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed under reduced pressure after filtration. The residue was recrystallized from EtOAc/isooctane giving 1.7 g (90 %). C<sub>14</sub>H<sub>16</sub>F<sub>2</sub>N<sub>2</sub>O<sub>4</sub> requires C, 53.50 H, 5.13 N, 8.91. Anal found C, 53.30 H, 5.26 N, 8.98.

15

## Example D8

## Synthesis of m-Nitrophenylacetyl-L-alanine 2,4,5-Trichlorophenyl Ester

m-Nitrophenylacetyl-L-alanine (1 eq.) and 2,4,5-tricholophenol (1.3 eq.) were stirred in dicholomethane. A 1.0 M solution of 1,3-

- dicyclohexylcarbodiimide in dichloromethane (1.2 eq.) was added and the mixture was stirred at ambient temperature for 16 hours. The resulting mixture was filtered and the filtrate was concentrated under reduced pressure. The resulting oil was purified by silica gel chromatography using 1:2 ethyl acetate/hexanes as the eluant to provide the title compound as a pink solid. For
- 25 C<sub>17</sub>H<sub>13</sub>Cl<sub>3</sub>N<sub>2</sub>O<sub>5</sub>: Calc. 47.30% C, 3.04% H, 6.49% N. Found 47.57% C, 3.18% H, 6.47% N.

### Example D9

Synthesis of D,L-α-Methylphenylglycine Ethyl Ester

10

15

20

The title was prepared using the procedures described in J.J. Fitt and H. W. Gschwend, J. Org. Chem., 42, No. 15, 2639 (1977). More specifically, D,L-phenylglycine (Aldrich) was stirred in dimethylformamide dimethylacetal and the mixture was heated at reflux under an atmosphere of dry nitrogen for 4 hours. After cooling, the mixture was concentrated under reduced pressure to provide a yellow oily solid. The mixture was slurried in diethyl ether and filtered through Celite. The filtrate was concentrated to an orange oil which was purified by vacuum distillation to provide a yellow oil which solidified. The yellow solid was stirred in dry THF at -20 C under dry nitrogen. Lithium bis(trimethylsilyl)amide (1.05 eq, 1.0M solution in THF) was added dropwise. The resulting mixture was allowed to warm to -10 C and stirring was continued for 1 hour at that temperature. Methyl iodide (1.05 eq) was added and the mixture was allowed to warm ambient temperature with stirring. After 14 hours, the mixture was concentrated. The residue was partitioned between aqueous potassium carbonate and chloroform. The organic portion was dried (sodium sulfate) and concentrated under reduced pressure. The product was purified by silica gel chromatography to yield a yellow oil. The yellow oil was stirred in absolute ethanol. Dry zinc chloride (4 eq.) was added and the mixture was heated at reflux. After 14 hours, the mixture was concentrated under reduced pressure to provide a yellow oil. The oil was partitioned between aqueous potassium carbonate and chloroform. The organic portion was dried (sodium sulfate) and concentrated under reduced pressure. The title compound was purified by silica gel chromatography.

## Example D10

25

30

## Synthesis of D,L-Phthalimidoalanine Ethyl Ester Hydrochloride

N-(Diphenylmethylene)glycine ethyl ester (1 eq.) (Aldrich) was stirred in dry THF at -78°C under an atmosphere of dry nitrogen. Lithium bis(trimethylsiyl)amide (1.02 eq, 1.0 M solution in THF) was added dropwise. The resulting mixture was stirred 1 hour at -78°C. A THF solution of N-(bromomethyl)phthalimide (1.1 eq) (Aldrich) was added and the mixture was

10

15

20

allowed to warm to ambient temperature and then stirring was continued for 1 hour. Hydrochloric acid (600mL, 2N) was added and the mixture was stirred for 20 minutes. The THF was removed on a rotoevaporator. The resulting aqueous mixture was washed with diethyl ether, and then concentrated (to 100 mL) to yield a thick slurry. A white solid was collected, washed with cold water and dried in a vacuum oven to yield the title compound which was used without further purification.

## Example D11

## Synthesis of N-(3-Nitrophenylacetyl)-L-alanine

The title compound was prepared by dissolving 9.27 g (0.0348 mols) of the N-(3-nitrophenylacetyl)-L-alanine methyl ester in 60 mL of dioxane and 15 mL of H<sub>2</sub>O and adding LiOH (3.06 g, 0.0731 mol) that has been dissolved in 15 mL of H<sub>2</sub>O. After stirring for 4 hours, the dioxane was removed under reduced pressure and the residue diluted with EtOAc, the layers were separated and the aqueous layer acidified to pH 2. The aqueous layer was back extracted with EtOAc (4 X 100 ml), the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed under reduced pressure after filtration. The residue was recrystallized from EtOAc/isooctane giving 7.5 g (85 %).  $C_{11}H_{12}N_2O_5$  requires C, 52.38 H, 4.80 N, 11.11. Anal found C, 52.54 H, 4.85 N, 11.08.  $[\alpha]_{23} = -29.9$  @ 589 nm.

### Example D12

## Synthesis of Methyl 2-Amino-2-(3-fluorophenyl)acetate Hydrochloride

Potassium cyanide (6.3, 0.1 mol) and ammonium carbonate (15.7 g, 0.2 mol) were dissolved in 50 mL of water (in a well ventilated fume hood). 3-Fluorobenzaldehyde (5.0 g, 0.04mol) was dissolved in 50 mL of EtOH and added to the reaction. After stirring at reflux under nitrogen atmosphere for 17 hours, the reaction was cooled to 23°C, the pH adjusted to 2.0 by the addition of 5 N HCl and cooled to 5°C. The resulting hydantoin was collected, rinsed

WO 98/22494 PCT/US97/20804

-- 156 --

with cold water and vacuum dried giving 3.59 of an off-white solid. The hydantoin was hydrolyzed at reflux using 1 N NaOH giving 2-amino-2-(3-fluorophenyl)acetic acid which was esterified via Procedure H in methanol to give the title compound.

5

10

15

## Example D13

## Synthesis of

## N-[N-(S)-2-Aminobutanoyl]-L-phenylglycine tert-Butyl ester

A mixture of N-[N-(benzyloxycarbonyl)-(S)-2-aminobutanoyl]-L-phenylglycine tert-butyl ester (4.13 g) (prepared from N-(benzyloxycarbonyl)-(S)-2-aminobutanoic acid (Novabiochem) and L-phenylglycine tert-butyl ester hydrochloride (Novabiochem) using General Procedure D) and 20% Pd(OH)<sub>2</sub>/C (0.360 g) in EtOH (200 mL) was shaken in a Parr Apparatus under a hydrogen atmosphere (40 psi) for 4 hours. The solids were removed by filtration through a plug of Celite, while rinsing with EtOH. The filtrate was concentrated to an off-white oil, which was used without further purification. <sup>1</sup>H-NMR in CDCl<sub>3</sub> revealed that ~10% trans-esterification to the ethyl occurred during this reaction. The ethyl ester was removed by flash chromatography after subsequent reaction of this compound.

## Example D14

20

25

30

## Synthesis of N-[N-L-Valinyl]-L-phenylglycine tert-Butyl ester

A mixture of N-[N-(benzyloxycarbonyl)-L-valinyl]-L-phenylglycine tert-butyl ester (4.63 g) (prepared from N-(benzyloxycarbonyl)-L-valine (Aldrich) and L-phenylglycine tert-butyl ester hydrochloride (Novabiochem) using General Procedure D) and 20% Pd(OH)<sub>2</sub>/C (0.360 g) in EtOH (200 mL) was shaken in a Parr Apparatus under a hydrogen atmosphere (40 psi) for 4 hours. The solids were removed by filtration through a plug of Celite, while rinsing with EtOH. The filtrate was concentrated to an off-white solid, which was used without further purification.  $^1$ H-NMR in CDCl<sub>3</sub> revealed that  $\sim 1$  % trans-esterification to the ethyl occurred during this reaction. The ethyl ester

was removed by flash chromatography after subsequent reaction of this compound.

## Example D15

## Synthesis of (S)-Phenylglycinol Methyl Ether

,

10

5

(S)-(+)-2-phenylglycinol (1 eq.) (Aldrich) was stirred in dry THF under an atmosphere of dry nitrogen. Sodium hydride (1 eq.) was added and the resulting mixture was stirred for 1 hour at ambient temperature. A THF solution of iodomethane (1 eq.) was added and the mixture was stirred for 1 hour. The mixture was concentrated to provide a residue which was taken up in water and extracted with chloroform. The organic extracts were concentrated under reduced pressure to yield the title compound as an oil which was purified by silica gel chromatography to yield a crude product which was used without further purification.

15

20

25

## Example D16

## Synthesis of (S)-2-Hydroxy-2-methyl-1-phenylprop-1-ylamine

To a stirred, cooled (0°C) suspension of 5.6 g (27.8 mmol) of L-phenylglycine methyl ester hydrochloride (Aldrich) in 200 mL of dry THF was added methylmagnesium bromide (46.3 mL, 138.9 mmol, 3.0 M in diethyl ether). During the addition, the internal temperature increased to 24°C. Stirring was continued for 1 hour, after which the reaction was carefully quenched by addition of saturated sodium bicarbonate solution. The reaction mixture was partitioned between ethyl acetate and aqueous sodium bicarbonate solution, back extracting the aqueous layer with 3 volumes of ethyl acetate. The combined organics were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The product was purified by flash chromatography on silica gel, eluting with 10% methanol in chloroform (neutralized with ammonium hydroxide) to afford 1.96 g to the title compound.

10

15

20

25

-- 158 ---

## Example D17

## Synthesis of 5-Chloro-2-thiophenecarboxaldehyde

A solution of 2-chlorothiophene (Aldrich; 1 molar eq.) in THF was cooled to -78°C and treated with *n*-butyllithium (1.6M in hexanes; 1.1 molar eq.) in a dropwise manner. The resultant yellow solution was stirred at -78°C for 40 minutes. Dimethylformamide (1.1 molar eq.) was added dropwise and the reaction stirred and additional 30 minutes. The mixture was diluted with methylene chloride and washed with 10% acetic acid, 1 M potassium carbonate, and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified by HPLC eluting with 15% ethyl acetate/hexanes to afford the title compound.

## Example D18

## Synthesis of (S)-(-)- $\alpha$ -Methylbenzylisocyanide

Prepared according to the general procedure of Wolber, E. K. A.; Ruchardt, C. Chem. Ber. 1991, 124, 1667. To a suspension of 1,1'-carbonyldiimidazole (1.6 molar eq.; Aldrich) in acetonitrile at 0°C was treated with methanesulfonic acid (3.2 molar eq.; Aldrich) in a dropwise fashion. A very thick suspension results. S-(-)-α-Methylbenzyl formamide (1 molar eq.; from Example D19 below) was added as a solution in acetonitrile via cannulation. The mixture was stirred overnight at ambient temperature. The suspension was filtered, washing with acetonitrile. The filtrate was concentrated and purified via flash chromatography eluting with 30% ethyl acetate/hexanes. The oil was further purified via bulb-to-bulb distillation (80°C, 0.04 mm Hg) giving a pale yellow oil in 51% yield. Calcd for C<sub>9</sub>H<sub>9</sub>N: C, 82.41; H, 6.92; N, 10.68. Found: C, 82.56; H, 6.82; N, 10.71.

Example D19

Synthesis of (S)-(-)- $\alpha$ -Methylbenzyl formamide

10

15

(S)-(-)- $\alpha$ -Methylbenzylamine (1 molar eq.) was treated with ethyl formate (80 molar eq.; Aldrich). A precipitate formed immediately. The suspension was heated to reflux (55°C) for 3 hours. The precipitate went into solution upon heating. The solution was cooled to ambient temperature and concentrated via rotary evaporation. The resultant solid was used without purification.

## Example D20

## Synthesis of 3-(Phenyl)benzaldehyde

A solution of 3-bromobiphenyl (Aldrich; 1 molar eq.) in dry THF was cooled to -78°C and treated with *tert*-butyllithium (Aldrich; 1.7 M in hexanes, 2 molar eq.) in a dropwise manner. The reaction was allowed to stir at -78°C for 40 minutes. Dimethylformamide (Aldrich; 2.5 molar eq.) was added and stirring continued an additional 20 minutes. The mixture was partitioned in a separatory funnel between methylene chloride and water. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via HPLC eluting with 5% ethyl acetate/hexanes. The desired aldehyde was obtained in 71% yield.

## Example D21

20

25

## Synthesis of 4-(Cyclohexyl)benzaldehyde

18-Crown-6 (Aldrich; 4 molar eq.) and pyridinium chlorochromate (Aldrich; 4 molar eq.) were added together in chloroform and stirred for 20 minutes. 4-Cyclohexylbenzylalcohol (from Example D22 below; 1 molar eq.) was added and stirring continued for 3 hours. Ether was added and the mixture filtered through a plug of silica eluting with ether. The solvent was removed via rotary evaporation. The residue was dissolved in ether and washed with water. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated.

10

15

20

25

30

## Example D22

## Synthesis of 4-(Cyclohexyl)benzyl Alcohol

To a solution of 4-cyclohexylbenzoic acid (Aldrich; 1 molar eq.) in toluene was added diiso-butylaluminum hydride (Aldrich; 1 M in toluene; 4 molar eq.) over a 2 hour period. After addition was complete, the reaction was heated to 60°C for 1 hour. The reaction was cooled to 5°C and quenched with saturated aqueous ammonium chloride. The layers were separated and the aqueous layer extracted with ethyl acetate. The combined organics were filtered to remove salts and concentrated.

## Example D23

## Synthesis of 3,5-Difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetic Acid

A solution of ethyl 3,5-difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetate (from Example D24 below; 1 molar eq.) in 50% aqueous ethanol was treated with lithium hydroxide (1.5 molar eq.). The solution was stirred for 3 hours at ambient temperature then concentrated via rotary evaporation. The residue was taken up in water; a small amount of 1 N NaOH was added to make basic. The aqueous mixture was extracted with ether. The aqueous layer was acidified to pH 3 with 1 N HCl. The acid was extracted thrice with methylene chloride. The combined methylene chloride extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated.

## Example D24

## Synthesis of Ethyl 3,5-Difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetate

Ethyl 3,5-difluorophenyl- $\alpha$ -ketoacetate (Rieke Metals, Inc. #14014; 1 molar eq.) was treated with (diethylamino)sulfur trifluoride (DAST) (2.5 molar eq.). The reaction was stirred at ambient for 72 hours then heated to 50°C for 6 hours. The mixture was poured over ice and extracted with methylene chloride. The organic layer was washed with saturated sodium bicarbonate,

10

dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via HPLC eluting with 2% ethyl acetate/hexanes.

## Example D25

## Synthesis of

N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-D,L-phenylglycine

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine methyl ester (from Example 111 below) was hydrolyzed according to Procedure AF. The acid was recrystallized from isooctane/EtOAc providing a mixture of diastereomers at the phenylglycine center. Elemental analysis; C<sub>19</sub>H<sub>18</sub>F<sub>2</sub>N<sub>4</sub>O<sub>4</sub> requires C, 60.63 H, 4.82 N, 7.44. Found; C, 60.65 H, 5.02 N, 7.37. Mass spectroscopy (MH<sup>+</sup> 377).

## Example D26

## Synthesis of 3-(4-Iodophenyl)propylamine

- N-(3-bromopropyl)phthalimide (1 eq., Aldrich) and 4-iodophenol (1 eq., Aldrich) and potassium carbonate (2 eq.) was stirred in acetonitrile. The mixture was heated at reflux. After 64 hour, the reaction mixture was concentrated to a thick mixture which was slurried in water. A white solid was collected, washed with water and vacuum dried.
- The white solid was stirred in ethanol. Anhydrous hydrazine (2 eq.) was added and mixture was heated at reflux for 18 hours. The reaction mixture was concentrated to yield a solid which was treated with 1N NaOH and extracted with CHCl<sub>3</sub>. The organic portion was dried, concentrated then diluted with ether. The mixture was treated with dry HCl. The title compound was collected as a white solid and vacuum dried.

## Example D27

Synthesis of 2-Amino-1-phthalimidopentane Hydrochloride

10

15

20

25

2-Amino-1-pentanol was stirred in a mixture of chloroform and saturated aqueous sodium bicarbonate. Di-*tert*-butyl dicarbonate (1.05 eq.) was added in one portion and the mixture was stirred until starting material was consumed. The organic portion was separated, dried (sodium sulfate) and concentrated. The crude material was purified by silica gel chromatrography using 1:1 ethyl acetate/hexanes.

The product was dissolved in THF. Triethylamine (1.1 eq.) was added and the mixture was cooled in an ice bath. Methanesulfonyl chloride (1.1 eq.) was added dropwise and the mixture was stirred until starting material was consumed. The mixture was concentrated under reduced pressure then was partitioned between ethyl acetate and water. The organic portion was separated, dried (sodium sulfate) and concentrated to yield a white solid which was chromatographed on silica gel using 30% ethyl acetate in hexanes and finally crystallized from 1-chlorobutane/hexanes.

The crystalline product was stirred in dry DMF and potassium phtalimide (1.1 eq.) was added. The mixture was stirred for 18 hours then was concentrated under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic portion was dried and concentrated to yield a white solid. The solid was taken up in chloroform and filtered through a plug of silica. Eluent containing product was concentrated to yield the crude product as a white solid.

The white solid was taken up in dry dioxane and resulting solution was saturated with gaseous HCl. After stirring for 30 minutes, the mixture was concentrated to yield a white solid which was triturated in ether. The title compound was collected, washed with ether and dried in a vacuum oven.

Example D28

Synthesis of D,L-3,5-Difluorophenylglycine

KOH (11.76 grams), LiCl (2.95 grams), saturated aqueous ammonia (20 mL), and benzyltriethylammonium chloride (0.805 grams) were stirred and chilled in CH<sub>2</sub>Cl<sub>2</sub> (17 mL). Gaseous ammonia was bubbled into this mixture with chilling (0°C) to saturation. To the resulting mixture was added 3,5-difluorobenzaldehyde (5.0 grams) (Lancaster) and chloroform (4.46 mL), dissolved in CH<sub>2</sub>Cl<sub>2</sub> (17.5 mL) with concurrent saturation with ammonia gas. The resulting mixture was stirred cold for 4 hours and at 22.5°C for 96 hours. Water (60 mL) and CH<sub>2</sub>Cl<sub>2</sub> (20 mL) were added; the layers separated, and the aqueous layer was extracted 3 times more with CH<sub>2</sub>Cl<sub>2</sub>. The aqueous layer was reduced *in vacuo* by 50%. The pH was adjusted to 6.5 with cold conc. HCl whereupon white crystals of D,L-3,5 difluorophenylglycine formed (3.4343 grams).

## Example D29

15

20

25

5

10

## Synthesis of L-3,5-Difluorophenylglycine Methyl Ester Tartate Salt

3.43 Grams of D,L-3,5 difluorophenylglycine (from Example D28 above) was slurried in 50 mL methanol and 2.5 mL conc. H<sub>2</sub>SO<sub>4</sub>. The reaction mixture was heated under gentle reflux for 18 hours. The mixture was chilled in and ice bath and the pH of the solution was adjusted to 7.0 with saturated aqueous ammonia. The volatile organic solvents were removed *in vacuo* and the aqueous portion was extracted three times with CH<sub>2</sub>Cl<sub>2</sub>; the combined organic layers dried, filtered, and reduced *in vacuo* to provide 2.680 grams of crude ester. This ester, benzaldehyde (1.4085 grams), and (-) tartaric acid (1.9921 grams), were dissolved in 20.5 mL of hot ethanol and stirred slowly for 72 hours as the title compound crystallized. The product was filtered and dried to provide 3.4805 grams of the (-) tartarate salt.

Example D30

Synthesis of N-(3,5-Difluorophenylacetyl)-L-3,5-difluorophenylglycine

10

15

20

25

L-3,5-Difluorophenylglycine (0.4291 g) (prepared from L-3,5-difluorophenylglycine (-)-tartarate salt (from Example D29 above) by neutralization) and 3,5-difluoroacetic acid 0.367 gram were dissolved in THF. EEDQ coupling using General Procedure AN afforded 0.7441 grams of the title compound as the methyl ester. The ester was dissolved in 1,4-dioxane (10 mL), chilled and LiOH.H<sub>2</sub>0 (89.0 mg) in water (10 mL) was added slowly and the mixture was stirred for 2 hours at 22.5°C. EtOAc (30 mL) and 1N HCl were added and the aqueous layer extracted two times. The combined organic layers were dried (MgSO<sub>4</sub>) and reduced *in vacuo* to provide the title compound (700.8 mg).

Each of the compounds set forth in the following examples was prepared by one of the following general procedures, unless otherwise indicated.

## GENERAL PROCEDURE A

## EDC Coupling Procedure I

To a 1:1 mixture of the corresponding carboxylic acid and amino ester/amide in CH<sub>2</sub>Cl<sub>2</sub> or DMF at 0°C was added 1.5 equivalents triethylamine, followed by 2.0 equivalents hydroxybenzotriazole monohydrate, then 1.25 equivalents of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC). The reaction mixture was stirred overnight at room temperature and then transferred to a separatory funnel. The mixture was washed with water, saturated aqueous NaHCO<sub>3</sub>, 1 N aqueous hydrochloric acid, and saturated aqueous sodium chloride, and then dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to yield the crude product.

## GENERAL PROCEDURE B

## **EDC Coupling Procedure II**

The carboxylic acid was dissolved in methylene chloride in a round-bottomed flask. The amino acid (1 eq.), N-methylmorpholine (5 eq.) and hydroxybenzotriazole monohydrate (1.2 eq.) were added in sequence. A

cooling bath was applied to the round-bottomed flask until the solution reached 0°. At that time, 1.2 eq. of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride was added. The solution was then allowed to stir overnight and come to room temperature under N<sub>2</sub> pressure. The reaction mixture was then washed with saturated aqueous Na<sub>2</sub>CO<sub>3</sub>, 0.1 M citric acid, and brine before drying with Na<sub>2</sub>SO<sub>4</sub> and removing the solvents to yield the crude product. Pure products were typically obtained by flash chromatography in an appropriate solvent.

## GENERAL PROCEDURE C

10

15

20

5

## EDC Coupling Procedure III

A round-bottomed flask was charged with the appropriate carboxylic acid (1.0 eq), hydroxybenzotriazole hydrate (1.1 eq) and the appropriate amine (1.0 eq) in THF under a nitrogen atmosphere. An appropriate amount (1.1 eq. for the free amine and 2.2 eq. for amine hydrochloride salt) of a suitable base, such as Hunig's base was added to the stirred mixture, followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.1 eq). After stirring for about 4 h to 17 h at room temperature, the solvent was removed at reduced pressure and the residue taken up in EtOAc (or a similar solvent)/H<sub>2</sub>O. The extracts were washed with saturated NaHCO<sub>3</sub>, 1 N aqueous hydrochloric acid, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. In some cases, the isolated product required further purification using standard procedures, such as chromatography and/or recrystallisation.

# GENERAL PROCEDURE D EDC Coupling Procedure IV

25

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at 0°C was charged with THF, an amine or amine hydrochloride (1.0 eq.), carboxylic acid (1.1 eq.), 1-hydroxybenzotriazole hydrate (1.15-1.2 eq.), N,N-diisopropylethylamine (2.3 eq.), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.15-1.2 eq.).

10

15

20

The cooling bath was removed and the mixture allowed to warm to room temperature with stirring for 10-20 hours. The mixture was diluted with EtOAc and washed with 0.5 N aqueous. HCl (2x), dilute aqueous NaHCO<sub>3</sub> (1x), brine (1x) and dried over either Na<sub>2</sub>SO<sub>4</sub> or MgSO<sub>4</sub>. The drying agent was removed by filtration and the filtrate concentrated *in vacuo*. The residue was either used without further purification or purified by standard procedures, such as flash chromatography on silica gel and/or recrystallization.

# GENERAL PROCEDURE E EDC Coupling Procedure V

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at 0°C or room temperature was charged with THF, carboxylic acid (1.0 eq), an amine or amine hydrochloride (1.0-1.1 eq.), 1-hydroxybenzotriazole hydrate (1.1-1.2 eq.), N,N-diisopropylethylamine (2.2-2.9 eq.), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.1-1.2 eq.). The cooling bath was removed and the mixture allowed to warm to room temperature with stirring for 10-20 hours. The mixture was diluted with EtOAc and washed with 0.5 N aqueous HCl (2x), dilute aqueous NaHCO<sub>3</sub> (1x), brine (1x) and dried over either Na<sub>2</sub>SO<sub>4</sub> or MgSO<sub>4</sub>. The drying agent was removed by filtration and the filtrate concentrated *in vacuo*. The residue was either used without further purification or purified using standard procedures, such as flash chromatography on silica gel and/or recrystallization.

# GENERAL PROCEDURE F EDC Coupling Procedure VI

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at 0°C was charged with THF, carboxylic acid (1.0 eq.), an amine or amine hydrochloride (1.1 eq.), N,N-diisopropylethylamine (2.2-2.3 eq.), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.1-1.2 eq.). The cooling bath was removed and the mixture allowed to warm to room temperature with stirring for 10-20 hours. The mixture was

diluted with EtOAc and washed with 0.2 N aqueous HCl (2x), dilute aqueous NaHCO<sub>3</sub> (1x), brine (1x) and dried over either Na<sub>2</sub>SO<sub>4</sub> or MgSO<sub>4</sub>. The drying agent was removed by filtration and the filtrate concentrated *in vacuo*. The residue was either used without further purification or purified using standard procedures, such as flash chromatography on silica gel and/or recrystallization.

## GENERAL PROCEDURE G

## Methyl Ester Preparation

To 1-methyl-3-nitro-1-nitrosoguanidine (1.2 eq.) in diethyl ether cooled to 0°C was added 40% KOH until bubbling ceased. This mixture was then decanted into a plastic tube containing KOH pellets as a drying agent. The solution was then added to the appropriate carboxylic acid and the mixture was stirred until the reaction was complete (as determined, for example, by tlc). The reaction was then quenched with acetic acid and extracted into EtOAc. Removal of the solvent afforded the desired methyl ester.

15

20

10

5

## GENERAL PROCEDURE H

## Carboxylic Acid Ester Preparation

To the appropriate amino acid or carboxylic acid in the appropriate alcohol was bubbled anhydrous HCL gas until the solution was saturated. The reaction was stirred overnight at 25°C and the solvent was then removed under reduced pressure. The residue was then dissolved in EtOAc and this solution was washed with sodium bicarbonate solution. The organic layer were then dried over sodium sulfate, filtered and solvent removed to afford the desired ester.

## GENERAL PROCEDURE I

## tert-Butyl Ester Preparation I

25

To a solution of an N-CBZ-protected amino acid in CH<sub>2</sub>Cl<sub>2</sub> was added 1.5 equivalents of N,N'-diisopropyl-O-t-butylisourea (prepared by standard literature methods such as those found in *Synthesis* (1979), p. 561), and the reaction was heated to reflux for 17 h. An additional 1.5 equivalents of isourea

were then added, and reflux was continued for another 7 h. The reaction was then cooled to room temperature and filtered through a bed of Celite 545, then stripped to dryness to leave a clear oil. The residue was dissolved in hexanes and filtered to remove solids, and the filtrate was washed with saturated aqueous NaHCO<sub>3</sub>, water, saturated aqueous NaCl, and dried over MgSO<sub>4</sub>. The solution was concentrated under reduced pressure to leave the product.

## GENERAL PROCEDURE J

### tert-Butyl Ester Preparation II

The reaction was conducted in a sealed tube using the appropriate carboxylic acid, a catalytic amount of H<sub>2</sub>SO<sub>4</sub> (0.03 eq.) and an excess of condensed *iso*-butylene in dioxane or CH<sub>2</sub>Cl<sub>2</sub> at -20°C. The reaction times varied from about 48 hours to about 120 hours. When the reaction was complete, the solvent was removed under reduced pressure and the residue dissolved in diethyl ether. This solution was washed with sodium bicarbonate solution and the organic layer dried over sodium sulfate, filtered and solvent removed. The resulting product was purified using standard procedures, such as HPLC or titration using, for example, diethyl ether/hexanes.

### GENERAL PROCEDURE K

### Amide Preparation I

20

25

5

10

15

To a solution of 3 equivalents of the desired amine in 1,2-dichloroethane was added 5.2 equivalents triethylaluminum subsurface. After stirring for 30 minutes at room temperature, a solution of the desired ester dissolved in 1,2-dichloroethane was added. The reaction was refluxed until tlc showed complete conversion, typically 3h. The reaction was then cooled to 0°C and quenched with 10% aqueous hydrochloric acid (Note: the acid should be added slowly as some foaming occurs during its addition). The mixture was transferred to a separatory funnel and the layers were separated. The aqueous phase was washed with ethyl acetate, and the organic phases were washed with saturated

10

15

20

25

aqueous NaCl, dried over MgSO<sub>4</sub>, and concentrated under reduced pressure to leave the crude product.

Alternatively, if the product is acid soluble, after the reaction is quenched, the reaction volume was reduced to about one-third of its initial volume under reduced pressure. To the resulting solution was added 20% aqueous potassium sodium tartrate (Rochelle's salt) and ethyl acetate. The pH of the solution was then adjusted to -13, and the aluminum salts dissolved in the aqueous solution. The organic phase was separated, and the aqueous phase was extracted with ethyl acetate. The combined organic solution was washed with saturated aqueous NaCl, dried over MgSO<sub>4</sub>, and concentrated under reduced pressure to leave the crude product.

## GENERAL PROCEDURE L

## Amide Preparation II

The carboxamide was prepared from its corresponding ester using the procedure described in Hogberg, T., et.al., J. Organic Chem., 1987, 52, 2033-2036.

## GENERAL PROCEDURE M

### Amide Preparation III

To the appropriate carboxylic acid (1.0 eq.) in THF was added N-methylmorpholine (1.1 eq.) and the solution was cooled to -20°C to 0°C. isobutyl chloroformate (1.1 to 2.1 eq.) was then added and the reaction mixture was stirred at -20°C to 0°C for 30 min. A mixture of the appropriate amino acid, water and 1.5 eq. of potassium carbonate was then added, and the resulting mixture was allowed to warm to room temperature and stir for 2 hrs. The reaction mixture was then poured into water and washed with EtOAc. The pH of the water layer was then adjusted to 2.0 with 5 N HCl and the water layer was extracted with EtOAc. The combined organic layers were dried over sodium sulfate, filtered and the solvent removed under reduced pressure. The

resulting crude amide was used without further purification or purified using standard procedures such as chromatography or titration using, for example, diethyl ether/hexanes or EtOAc/hexanes.

### GENERAL PROCEDURE N

5

10

### Hydrolysis of Carboxylic Acid Esters

To the ester in a 1:1 mixture of CH<sub>3</sub>OH/H<sub>2</sub>O was added 2-5 equivalents of K<sub>2</sub>CO<sub>3</sub>. The mixture was heated to 50°C for 0.5-1.5 h until tlc showed complete reaction. The reaction was cooled to room temperature and the methanol was removed on a rotary evaporator. The pH of the remaining aqueous solution was adjusted to about 2, and ethyl acetate was added to extract the product. The organic phase was then washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to yield the product.

## GENERAL PROCEDURE O

15

## Removal of N-Carbobenzyloxy (CBZ) Protecting Groups

The N-CBZ-protected compound was dissolved in ethanol in a hydrogenation flask and a catalytic amount of 10% Pd/C was added. The mixture was hydrogenated at 20 psi H<sub>2</sub> on a Parr shaker for 30 min. The reaction was then filtered through a pad of Celite 545 and stripped free of solvent on a rotary evaporator to yield the product.

20

25

## GENERAL PROCEDURE P

## Removal of the N-tert-Boc Protecting Group

The *N-tert*-Boc-amine was dissolved in a suitable dry solvent (such as 1,4-dioxane or ethyl acetate) and the solution was cooled in an ice bath. Gaseous HCl was introduced into the solution until the mixture was saturated with HCl. The mixture was then stirred until the reaction was complete. The resulting mixture was concentrated under reduced pressure to yield the amine hydrochloride. The amine hydrochloride was used without purification or was

WO 98/22494 PCT/US97/20804

-- 171 --

triturated using, for example, diethyl ether and the resulting solid was collected by filtration.

## GENERAL PROCEDURE Q

## Halide Exchange (Finkelstein) Reaction

5

10

15

20

25

The corresponding alkyl bromide or alkyl chloride was dissolved in 20 mL of methyl ethyl ketone and 1 eq. of NaI was added. The reaction was heated to 60°C and stirred overnight. The cooled reaction mixture was extracted with dichloromethane (2 x 30 mL) and the combined extracts were roto-evaporated at reduced pressure to give the crude product. Pure products were typically obtained by flash chromatography in an appropriate solvent.

### GENERAL PROCEDURE R

### Oxime Reduction I

To the oxime ester in the alcohol corresponding to the ester was added formic acid (500 eq.) and water (500 eq.). The reaction mixture cooled to 5°C and zinc dust (3.8 eq.) was added in portions over 20 min. The reaction was then allowed to warm to room temperature and stirring was continued for 3 hours. The reaction was then filtered over HYFLO and the solvent removed under reduced pressure. The residue was dissolved in EtOAc and this solution washed with saturated sodium bicarbonate solution. The organic layer was then dried over sodium sulfate, filtered and solvent removed to afford the product.

### GENERAL PROCEDURE S

### Reduction of Esters to Alcohols

To a 0°C solution of the starting ester in anhydrous THF was added 1.0 equivalents of LiBH<sub>4</sub> in THF. The reaction was stirred at room temperature overnight and then quenched with water. The THF was removed on a rotary evaporator and ethyl acetate was added. The phases were separated and the organic phase was washed with saturated aqueous sodium chloride solution,

10

15

20

25

dried over magnesium sulfate, and concentrated under reduced pressure to afford the alcohol product.

## GENERAL PROCEDURE T

## CDI Coupling Procedure

A solution of the appropriate acid (3.3 mmol) and 1,1'-carbodiimidazole (CDI) in 20 mL THF was stirred for 2 h. The amino acid ester hydrochloride (3.6 mmol) was added, followed by 1.5 mL (10.8 mmol) of triethylamine. The reaction mixture was stirred overnight and then dissolved in 100 mL of diethyl ether, washed with 10% HCl three times, brine once, 20% potassium carbonate once and brine once. The solution was dried over magnesium sulfate, filtered, and evaporated at reduced pressure to yield the product.

# GENERAL PROCEDURE U EDC Coupling Procedure VII

A mixture of the appropriate carboxylic acid (1 eq.), 1-hydroxybenzotriazole (1.6 eq.), the appropriate amine (1 eq.), N-methylmorpholine (3 eq.) and dichloromethane (or DMF for insoluble substrates), cooled in an ice-water bath, was stirred until a clear solution was obtained. EDC (1.3 eq.) was added to the reaction mixture and the cooling bath was allowed to warm to ambient temperature over 1-2 h. The reaction was then stirred overnight. The reaction mixture was then evaporated at reduced pressure to dryness under vacuum and 20% aqueous potassium carbonate was added to the residue. The mixture was shaken vigorously and allowed to stand for hours or overnight, if necessary, until the oily product to solidify. The solidified product was then filtered off, washed thoroughly with 20% potassium carbonate, water, 10% HCl, and water to give the product. No racemization was observed using this procedure.

GENERAL PROCEDURE V

O-Acylation of Alcohols

To a solution of the alcohol (e.g., N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide from Example 228 below) in pyridine was added 4 equivalents of acetic anhydride and the reaction was stirred at room temperature for 2.5 h. The reaction was quenched onto ice and then ethyl acetate was added and the phases were separated. The organic phase was washed with 10% HCl, water, saturated aqueous NaCl, and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to yield the product.

## GENERAL PROCEDURE W

10

15

5

## O-Esterification of Alcohols

To a suspension of 0.95 equivalents of NaH in THF was added an alcohol (e.g., N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide from Example 228 below) dissolved in THF. This solution was cooled to 0°C, then 1.1 equivalents of an acyl chloride (e.g. trimethylacetyl chloride) was added. The reaction was stirred at room temperature overnight, then was quenched with water and ethyl acetate. The organic phase was washed with water, saturated aqueous NaCl, and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to yield the crude product.

## GENERAL PROCEDURE X

20

25

## **BOP Coupling Procedure**

A solution of the carboxylic acid (1.0 eq.) and N-methyl morpholine (1.5 eq.) in dichloromethane was cooled to -20°C under nitrogen. BOP (1.05 eq.) was added in one portion and the reaction mixture was maintained at -20°C for 15 minutes. The appropriate alcohol (1.2 eq.) was added and the reaction mixture was allowed to warm to room temperature and stirring was continued for 12 hours. The reaction mixture was then poured into water and extracted with ethyl acetate (3x) and the combined organic layers were washed with saturated aqueous citric acid (2x), saturated aqueous sodium bicarbonate (2x),

10

15

20

25

brine (1x), and then rotoevaporated at reduced pressure to provide the crude product.

### GENERAL PROCEDURE Y

## **BOC** Removal Using TFA

The Boc-protected compound was added to a 1:1 mixture of dichloromethane and trifluoroacetic acid (TFA) and the reaction mixture was stirred until tlc indicated complete conversion, typically 2 hours. The solution was then stripped to dryness. The residue was suspended in dichloromethane and again stripped to dryness to remove excess TFA. The residue was placed under high vacuum for several hours to afford the desired TFA salt.

## GENERAL PROCEDURE Z

### Amide Preparation IV

The trichlorophenyl ester (1 eq.) was stirred in DMF or THF and the oxime or amine (1.2 eq.) was added. The mixture was stirred at ambient temperature for 1-4 hours. In cases where the hydrochloride salt form of an amine was used, a suitable base such as diisopropylethylamine (1.2 eq.) was also added. The resulting mixture was concentrated under reduced pressure to yield an oil or residue which was used without further purification or was purified by standard procedures, such as silica gel chromatography and/or recrystallization.

## GENERAL PROCEDURE AA

## Sodium Borohydride Reduction

The ketone was dissolved in MeOH and treated with 1.0 equivalent of sodium borohydride. The reaction was stirred until tlc showed the starting material was consumed, typically 1 hour. The reaction mixture was then evaporated at reduced pressure and chromatographed to afford the alcohol product.

10

15

20

## GENERAL PROCEDURE AB

## Preparation Amino Acid Derivatives Using Chiral Amines

(S)-(+)- $\alpha$ -Methylbenzyl amine was added dropwise to a solution of 4-(phenyl)benzaldehyde (1 molar eq.) in THF followed by the addition of 1.0 molar eqivalent of zinc chloride. The reaction mixture was allowed to stir at room temperature for 5 h. The cloudy mixture was then cooled to -30°C and treated with tert-butylisocyanide (1.05 molar eq.). After 20 minutes, N-(3,5difluorophenylacetyl)-L-alanine was added and stirring was continued at -30°C for 120 h. The reaction mixture was then poured into a seperatory funnel and diluted with CH2Cl2, washed with sodium bicarbonate. The organic layer was then washed with 0.5 N HCl, followed by brine. The organic layer was then dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give N-tert-butyl-N'-[N-(3,5difluorophenylacetyl)-L-alaninyl]-N'-(S)- $\alpha$ -methylbenzyl-2-amino-2-(4phenylphenyl)acetamide, as a mixture of isomers. At this stage, the isomers were typically separated by HPLC chromatography using, for example, a gradient of 30 to 35% EtOAc/hexanes. The  $\alpha$ -methylbenzyl protecting group was then removed from the S,S isomer by added 10 molar equivalents of triethylsilane and 20 molar equivalents of trifluoroacetic acid to the S,S isomer. The reaction was then heated to 37°C for 3 h and then poured into ethyl acetate and washed with sodium bicarbonate. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was purified by recrystallisation from ethyl acetate or ethyl acetate/hexanes. Various other aldehydes and carboxylic acids can be used in this procedure to provide for a variety of compounds useful in this invention.

25

30

## GENERAL PROCEDURE AC

## Oxime Reduction II

To a solution of the oxime ester in the alcohol corresponding to the ester was added a catalytic amount of acetic acid and 0.1 mole equivalent of 10% Pd/C. The reaction vessel (Parr shaker) was charged with hydrogen to 40 PSI and this mixture was shaken for 3 h. The reaction mixture was then filtered

over HyFlo and concentrated. The residue was dissolved in ethyl acetate and washed with a saturated solution of sodium bicarbonate. The organic layer was then dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give the desired amine.

## GENERAL PROCEDURE AD

5

10

15

20

25

## Mitsunobu Reaction

To a solution of N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester in 20 mL of THF was added 1.3 equivalents each of triphenylphosphine and diethyl azodicarboxylate (DEAD), and 1.0 equivalents of an alcohol. The mixture was stirred a room temperature overnight and the solvent was then removed. The residue was purified by standard procedures, such as chromatography and/or recrystallization.

# GENERAL PROCEDURE AE O-Alkylation of Tyrosine Derivatives

To a solution of N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester in 20 mL of acetone was added 1.3 equivalents of an alkyl bromide and 3.0 equivalents of potassium carbonate as a fine powder and a catalytic amount of sodium iodide. The reaction mixture was stirred at room temperature overnight and then partitioned between DCM and water. The organic layer was dried over anhydrous sodium sulfate, stripped of solvents and purified using standard procedures, such as chromatography and/or recrystallization.

# GENERAL PROCEDURE AF Hydrolysis of Carboxylic Acid Esters

A solution of the carboxylic acid ester (1.0 eq.) and lithium hydroxide (1.1 eq.) in 1:2 water/dioxane was stirred at 23°C for 1 hour. The reaction mixture was then acidified to pH 3 with 1 N HCl and extracted with ethyl acetate.

Concentration of the ethyl acetate extracts provided the product. In some cases,

20

25

the product was further purified using standard procedures, such as chromatography and/or recrystallization.

## GENERAL PROCEDURE AG

## Methyl Ester Formation from Amino Acids

The amino acid (amino acid or amino acid hydrochloride) is suspended in methanol and chilled to 0°C. HCl gas is bubbled through this solution for 5 minutes. The reaction is allowed to warm to room temperature then stirred for 4 hours. The solvents are then removed to afford the desired amino acid methyl ester hydrochloride. This product is usually used without further purification.

## GENERAL PROCEDURE AH <u>EEDO Coupling Procedure</u>

A round bottom flask containing a magnetic stir bar under as atmosphere of nitrogen at room temperature was charged with THF, the carboxylic acid (1 eq.), the amine hydrochloride (1.1 eq.) and EEDQ (1.1 eq.) and the reaction mixture was allowed to stir for 15 minutes. 4-Methylmorpholine (1.1 eq.) was added to the reaction and stirring was continued at room temperature for 15-20 hours. The reaction mixture was then concentrated *in vacuo* and the resulting residue was partitioned between ethyl acetate and water. The organic phase was separated and washed with saturated aq. NH<sub>4</sub>Cl (2x), saturated aq. NaHCO<sub>3</sub> (2x), followed by brine (1x). Organic phase dried over Na<sub>2</sub>SO<sub>4</sub>. The drying agent was removed by filtration and the filtrate concentrated *in vacuo*. The residue was either used without further purification or purified using standard procedures, such as flash chromatography on silica gel and/or recrystallization.

GENERAL PROCEDURE AI

N-tert-BOC Protection of Amino Acids

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at room temperature was charged with dioxane, water, 1.0 N aq. sodium hydroxide, and the amino acid (1 eq). Stirring was initiated and the flask was cooled in an ice bath. Di-t-butyldicarbonate (1.1 eq) was added to the reaction mixture, followed by removal of the ice bath and slow warming to room temperature over 1 hour. The reaction was partially concentrated on the rotary evaporator followed by the addition of ethyl acetate. The flask was recooled in an ice bath and the mixture was acidified to a pH of 2-3 through the addition of potassium bisulfate. The reaction was transferred into a seperatory funnel and the organic layer was separated. The aqueous layer was extracted with ethyl acetate and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>. The drying agent was removed by filtration and the filtrate concentrated in vacuo. The solid was used without further purification.

### GENERAL PROCEDURE AJ

15

20

5

10

## Removal of N-Carbobenzyloxy (CBZ) Protecting Groups

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at room temperature was charged with methanol, tetrahydrofuran,  $20\% \text{ Pd}(OH)_2/C$  (1 mass eq.), and the CBZ-protected dipeptide. Stirring was initiated and the flask was purged (3x) with hydrogen. The reaction mixture was allowed to stir at room temperature overnight under an atmosphere of hydrogen. The reaction was filtered and the filtrate was concentrated *in vacuo*. The resulting solid was used as is or purified via silica gel chromotography.

### GENERAL PROCEDURE AK

## Addition of N-Carbobenzyloxy (CBZ) Protecting Groups

25

A round bottom flask containing a magnetic stir bar under an atmosphere of nitrogen at room temperature was charged with water, sodium carbonate (2.2 eq.), and amino acid (1.0 eq.). The slurry was stirred at room temperature for 1 hour. Benzylchloroformate was added to the reaction and stirring was continued overnight. The reaction mixture was extracted with  $CH_2Cl_2$  (3x) and

10

15

20

the combined organic extracts were acidified to a pH of 2-3. The resulting solid was isolated via vacuum filtration.

#### GENERAL PROCEDURE AL

#### Preparation of Amino Acid Derivatives Using Chiral Amines II

A solution of aryl aldehyde (1 molar eq.) in THF was treated with S-(-)- $\alpha$ methylbenzylamine (1 molar eq.), followed by MgSO<sub>4</sub>. The reaction mixture was stirred for 1 hour then treated with tert-butylisocyanide (1.5-2.0 molar eq.) and N-(3,5-difluorophenylacetyl)-L-alanine (1.5-2.0 molar eq.). The reaction was allowed to stir for 60 hours. The reaction was diluted with methylene chloride and washed with 0.01 N HCl and saturated aqueous NaHCO<sub>3</sub>. Each aqueous wash was back-extracted with methylene chloride. The combined organics were washed with brine, dried over Na2SO4, filtered and concentrated to give N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-N'-R- $\alpha$ methylbenzyl-2-amino-2-DL-(aryl)acetamide. At this stage, the isomers were separated if possible by HPLC chromatography using, for example, a gradient of 20 to 25% ethyl acetate/hexanes. The  $\alpha$ -methylbenzyl protecting group was then removed from the peptide by adding 10 molar equivalents of triethylsilane and 20 molar equivalents of trifluoroacetic acid to the compound. The reaction was heated to 37°C for 3 hours and then poured into ethyl acetate and washed with sodium bicarbonate. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified by trituration with ether or ether/hexanes. Various other aldehydes, isocyanides, and carboxylic acid can be used in this procedure to provide for a variety of compounds useful in this invention.

25

#### GENERAL PROCEDURE AM

Preparation of Amino Acid Derivatives Using Chiral Amines III

A solution of an aromatic aldehyde (3 molar eq.) and S-(-)- $\alpha$ -methylbenzylamine (1 molar eq.) in methanol was treated with titanium(IV) isopropoxide (1.5 molar eq.). After stirring the mixture at ambient temperature

10

15

for 6 hours, tert-butylisocyanide (1.1 molar eq.) was added followed by N-(3,5difluorophenylacetyl)-L-alanine (1.2 molar eq.) 40 minutes later. The reaction mixture was stirred for 72 hours. The methanol was removed via rotary evaporation. The residue was dissolved in methylene chloride and washed with 0.01 N HCl. The emulsion was filtered through celite washing with methylene chloride. The layers were separated; the organic phase was washed with saturated NaHCO<sub>3</sub> and brine. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to give N-tert butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-N'-R- $\alpha$ -methylbenzyl-2-amino-2-DL-(aryl)acetamide. At this stage the isomers were separated if possible by HPLC chromatography using, for example, a gradient of 20 to 25% ethyl acetate/hexanes. The  $\alpha$ -methylbenzyl protecting group was then removed from the peptide by adding 10 molar equivalents of triethylsilane and 20 molar equivalents of trifluoroacetic acid to the compound. The reaction was heated to 37°C for 3 hours and then poured into ethyl acetate and washed with sodium bicarbonate. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified by trituration with ether or ether/hexanes. Various other aldehydes, isocyanides, and carboxylic acid can be used in this procedure to provide for a variety of compounds useful in this invention.

20

25

### GENERAL PROCEDURE AN EEDO Coupling Procedure II

To a 1:1 mixture of the corresponding carboxylic acid and amino ester/amide in THF at 0°C was added 1.1 equivalents of EEDQ. The reaction mixture was stirred for 18 hours at 22.5°C. The solvent was removed under reduced pressure or under a stream of nitrogen and the residue dissolved in EtOAc. The organic solution was washed 1 time with saturated NaHCO<sub>3</sub> solution, 1 time with N HCl, and dried over MgSO<sub>4</sub>. The organic solution was reduced *in vacuo* to yield the product.

#### GENERAL PROCEDURE AO

#### Preparation of Primary Amides

A sealable pressure tube containing a magnetic stir bar under an atmosphere of nitrogen at room temperature was charged with a methyl ester (1 eq.), sodium cyanide (0.1 eq.) and a 7M solution of ammonia in methanol. The tube was sealed and heated to 45°C with stirring for 18 hours. The reaction was allowed to cool to room temperature and the resulting precipitate was isolated by vacuum filtration. The solid was either washed with methanol or recrystallyzed from ethyl acetate/methanol.

10

15

5

#### Example 1

## Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate

Following General Procedure A (without the 1N HCl wash) and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2) and norleucine methyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 142-143°C). The reaction was monitored by tlc (Rf = 0.71 in 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>, 0.22 in 50% EtOAc/hexanes) and the product was purified by silica plug chromatography using CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.90$  (d, J=7.69 Hz, 1H), 6.80 (m, 3H), 6.70 (m, 1H), 4.62 (quint, J=7.2 Hz, 1H), 4.48 (m, 1H), 3.72 (s, 3H), 3.51 (s, 2H), 1.78 (m, 1H), 1.60 (m, 1H), 1.36 (d, J=7.02 Hz, 3H), 1.25 (m, 4H), 0.85 (m, 3H).

25  $^{13}$ C-nmr (CDCl<sub>3</sub>):  $\delta = 173.23$ , 172.69, 169.97, 165.30, 165.12, 162.00, 139.01, 138.88, 138.76, 112.93, 112.83, 112.70, 112.60, 103.63, 103.30, 102.97, 52.94, 49.38, 43.28, 32.32, 27.95, 22.75, 19.23, 14.35.  $C_{18}H_{24}F_{2}N_{2}O_{4}$  (MW = 370.40); mass spectroscopy (MH<sup>+</sup>) 371.

#### Example 2

#### Synthesis of

#### N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-histidine Methyl Ester

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2) and L-histidine methyl ester dihydrochloride (Sigma), the title compound was prepared as a solid (mp =  $195-197^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.29 in 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>). NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 7.60$  (s, 1H), 7.00-6.81 (m, 4H), 4.70 (t, 1H), 4.39 (q, 1H), 3.72 (s, 3H), 3.60 (s, 2H), 3.22-3.00 (m, 2H), 1.38 (d, 3H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta$  = 175.46, 172.56, 172.94, 166.64, 166.47, 163.38, 163.20, 141.73, 141.60, 141.47, 136.85, 113.92, 113.82, 113.70, 113.59, 103.89, 103.55, 103.21, 54.55, 53.31, 51.00, 43.21, 43.19, 30.36, 18.44. C<sub>18</sub>H<sub>20</sub>F<sub>2</sub>N<sub>4</sub>O<sub>4</sub> (MW = 394.38); mass spectroscopy (MH<sup>+</sup>) 395.

15

10

5

#### Example 3

# Synthesis of N-Benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

Following General Procedure K and using methyl N-[N-(3,5-20 difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate (from Example 1 above) and benzylamine (Aldrich), the title compound was prepared as a solid (mp = >200°C). The reaction was monitored by tlc (Rf = 0.29 in 5% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by preparative plate chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.05$  (m,5H), 6.65 (m, 3H), 4.10 (m, 4H), 3.35 (d, 2H), 1.35 (m, 9H), 0.65 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 175.48, 174.75, 173.16, 166.64, 166.46, 163.37, 141.55, 141.42, 140.38, 130.04, 129.95, 129.05, 128.95, 128.73, 113.94,$ 

113.83, 113.71, 113.60, 103.90, 103.88, 103.56, 103.22, 55.43, 51.26, 44.53, 43.21, 33.38, 29.56, 23.91, 18.28, 14.78.

 $C_{24}H_{29}F_2N_3O_3$  (MW = 445.51); mass spectroscopy (MH<sup>+</sup>) 446.

#### Example 4

5

10

# Synthesis of N-2-(N,N-Dimethylamino) ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

Following General Procedure K and using methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate (from Example 1 above) and <math>N,N-dimethylethylenediamine (Aldrich), the title compound was prepared as a solid (mp = 182-187°C). The reaction was monitored by tlc (Rf = 0.51 in 15% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by preparative plate chromatography using 15% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.2$ , 172.0, 170.0, 165.4, 165.3, 163.9, 162.1, 162.0, 139.1, 138.8, 113.1, 112.8, 103.6, 103.3, 103.0, 58.1, 54.0, 49.7, 45.7, 43.3, 38.1, 33.2, 28.2, 23.0, 19.2, 14.4.

 $C_{21}H_{32}F_2N_4O_3$  (MW = 426.51); mass spectroscopy (MH<sup>+</sup>) 427.

#### Example 5

# Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

25

20

Following General Procedure K and using methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate (from Example 1 above) and 2-methoxyethylamine (Aldrich), the title compound was prepared as a solid (mp = >200°C). The reaction was monitored by tlc (Rf = 0.42 in 10%

CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by flash chromatography using 12% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.85$  (bd, J=8.79 Hz, 0.5H), 7.64 (bd, J=7.81 Hz, 0.5H), 7.35 (m, 1H), 7.16 (bd, J=7.27 Hz, 0.5H), 7.06 (bs, 0.5H), 6.83 (m, 2H), 6.68 (m, 1H), 4.70 (m, 2H), 3.56 (d, J=9.89 Hz, 2H), 3.40 (m, 7H), 1.57 (m, 10H), 0.84 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.62, 172.58, 172.14, 172.04, 170.02, 169.91, 165.33, 165.15, 162.08, 112.99, 112.92, 112.82, 112.77, 112.66, 112.59, 103.54, 103.34, 103.31, 103.29, 71.46, 71.44, 59.27, 59.24, 53.76, 49.64, 49.43, 43.29, 39.79, 33.26, 33.22, 28.10, 28.03, 22.97, 22.91, 19.71, 19.61, 19.56, 19.51, 14.46, 14.43.

 $C_{20}H_{29}F_2N_3O_4$  (MW = 413.47); mass spectroscopy (MH<sup>+</sup>) 414.

#### Example 6

15

20

30

5

10

# Synthesis of N-2-(N,N-Dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure K and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94 below) and N,N-dimethylethylenediamine (Aldrich), the title compound was prepared as a solid (mp = 174-182°C). The reaction was monitored by tlc (Rf = 0.31 in 10%  $CH_3OH/CH_2Cl_2$ ) and the product was purified by preparative plate chromatography using 10%  $CH_3OH/CH_2Cl_2$  as the eluent.

NMR data was as follows:

25  $^{1}$ H-nmr (CD<sub>3</sub>OD):  $\delta = 7.22$  (m, 5h), 6.85 (m, 3H), 4.51 (m, 1H), 4.18 (m, 1H), 3.57 (m, 2H), 3.50-2.45 (m, 6H), 2.39 (s, 6H), 1.26 (d, 2.4H), 1.10 (d, 0.6H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta$  = 176.03, 175.50, 174.20, 173.99, 173.50, 173.22, 166.63, 166.46, 163.36, 163.19, 141.65, 141.52, 141.39, 139.38, 139.00, 130.90, 130.74, 130.05, 130.01, 128.37, 128.30, 114.03, 113.93, 113.80, 113.70, 103.96, 103.62, 103.57, 103.28, 59.18, 59.14, 56.78, 56.51, 51.93,

51.74, 45.53, 45.47, 43.21, 43.18, 42.92, 38.84, 38.65, 37.94, 37.85, 18.09, 17.73.

 $C_{24}H_{30}F_2N_4O_3$  (MW = 460.53); mass spectroscopy (MH<sup>+</sup>) 461.

#### Example 7

5

10

# Synthesis of N-(4-Pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure K and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94 below) and 4-(aminomethyl)pyridine (Aldrich), the title compound was prepared as a solid (mp = >200°C). The reaction was monitored by tlc (Rf = 0.46 in 10%  $CH_3OH/CH_2Cl_2$ ) and the product was purified by recrystallization from ethyl acetate.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta$  = 8.37 (d, 2H), 7.25 (m, 5H), 7.11 (d, 2H), 6.85 (m, 3H), 4.56 (t, 1H), 4.29 (m, 3H), 3.64 (s, 2H), 3.08 (m, 2H), 1.30 (d, 3H). <sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta$  = 175.46, 174.04, 173.26, 166.60, 166.43, 163.34, 163.16, 150.97, 150.44, 141.59, 141.45, 138.84, 130.95, 130.13, 128.44, 124.28, 113.98, 113.87, 113.75, 113.64, 103.91, 103.57, 103.23, 62.08, 57.01, 43.33, 43.12, 38.93, 21.41, 18.16, 15.02.

 $C_{26}H_{26}F_2N_4O_3$  (MW = 480.52); mass spectroscopy (MH<sup>+</sup>) 481.

#### Example 8

# Synthesis of N-(3-Pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

25

Following General Procedure K and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94 below) and 4-(aminomethyl)pyridine (Aldrich), the title compound was prepared as a solid (mp = 199-210°C). The reaction was monitored by tlc (Rf = 0.46 in 10%)

 $CH_3OH/CH_2Cl_2$ , minor isomer Rf = 0.50) and the product was purified by preparative plate chromatography using 10%  $CH_3OH/CH_2Cl_2$  as the eluent.

NMR data was as follows:

38.77, 18.18, 17.79.

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 8.42$  (m, 2H), 7.61 (m, 1H), 7.29 (m, 6H), 6.90 (m, 3H), 4.61 (m, 1H), 4.33 (m, 3H), 3.58 (s, 1.5H), 3.54 (s, 0.5H), 3.10 (m, 2H), 1.33 (d, 2.25H), 1.15 (d, 0.75H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta = 176.00$ , 175.34, 174.03, 173.81, 173.23, 166.61, 166.44, 163.35, 163.17, 149.93, 149.20, 141.48, 139.20, 138.72, 138.10, 138.03, 136.88, 136.79, 130.89, 130.70, 130.06, 130.02, 128.40, 128.33, 125.71, 113.97, 113.87, 113.74, 113.64, 103.92, 103.58, 103.53, 103.23, 56.88, 56.66, 55.74, 53.21, 43.22, 43.15, 42.89, 42.06, 41.98, 39.04, 38.88,

 $C_{26}H_{26}F_2N_4O_3$  (MW = 480.52); mass spectroscopy (MH<sup>+</sup>) 481.

#### Example 9

15

20

30

5

10

# Synthesis of N-(4-Pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

Following General Procedure K and using methyl  $N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate (from Example 1 above) and 4-(aminomethyl)pyridine (Aldrich), the title compound was prepared as a solid (mp = 181-205°C). The reaction was monitored by tlc (Rf = 0.51 in 10% <math>CH_3OH/CH_2Cl_2$ ) and the product was purified by preparative plate chromatography using 10%  $CH_3OH/CH_2Cl_2$  as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 8.48$  (m, 0.8H), 8.42 (m, 1.2H), 7.37 (d, J=6.10, 0.8H), 7.28 (d, J=6.11, 1.2H), 6.85 (m, 3H), 4.39 (m, 4H), 3.61 (s, 0.8H), 3.53 (d, J=2.99, 1.2H), 2.05-1.25 (m, 9H), 0.90 (m, 3H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta = 176.61$ , 175.71, 175.33, 175.29, 173.32, 173.24, 166.49, 166.32, 163.22, 163.05, 151.30, 151.24, 150.55, 150.41, 141.54, 141.41, 124.35, 124.20, 113.95, 113.85, 113.72, 113.62, 103.86, 103.57,

WO 98/22494 PCT/US97/20804

-- 187 --

103.52, 103.18, 55.72, 55.64, 51.98, 43.38, 43.19, 42.82, 33.07, 32.57, 29.87, 29.67, 23.90, 23.82, 18.24, 17.86, 14.80.

 $C_{23}H_{28}F_2N_4O_3$  (MW = 446.50); mass spectroscopy (MH<sup>+</sup>) 447.

#### Example 10

5

10

15

# Synthesis of tert-butyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate

#### Step A - t-Butyl Ester Formation

To a solution of Z-norleucine-OH in CH<sub>2</sub>Cl<sub>2</sub> was added 1.5 equivalents of N,N'-diisopropyl-O-t-butylisourea (prepared by the method of Synthesis (1979) p.561 for review) and the reaction was heated to reflux for 17 hours. An additional 1.5 equivalents of isourea was then added, and reflux was continued for another 7 hours. The reaction was then cooled to room temperature and filtered through a bed of Celite 545, then stripped to dryness to leave a clear oil. The residue was dissolved in hexanes and filtered to remove solids, and the filtrate was washed with saturated aqueous NaHCO<sub>3</sub>, water, saturated aqueous NaCl, and dried over MgSO<sub>4</sub>. The solution was concentrated under reduced pressure to leave the product.

#### Step B - CBZ Removal

20

The CBZ-protected amino ester was dissolved in ethanol in a hydrogenation flask and a catalytic amount of 10% Pd/C was added. The mixture was hydrogenated at 20 psi H<sub>2</sub> on a Parr shaker for 30 min. The reaction was then filtered through a pad of Celite 545 and stripped free of solvent on a rotary evaporator to yield the product, norleucine *tert*-butyl ester hydrochloride.

#### 25 Step C

Following General Procedure D and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and the norleucine tert-butyl ester hydrochloride, the title compound was prepared as a semi-solid. The reaction was monitored by tlc (Rf = 0.41 in 50% EtOAc/hexanes) and the product was

purified by flash chromatography using 50% EtOAc/hexanes as the eluent, followed by preparative plate chromatography using 50% EtOAc/hexanes as the eluent.

NMR data was as follows:

5  $^{1}$ H-nmr (CDCl<sub>3</sub>):  $\delta = 7.63$  (d, J=7.7 Hz, 1H), 7.34 (d, J=7.7 Hz, 1H), 6.8 (m, 2H), 6.7 (m, 1H), 4.8 (m, 1H), 4.36 (q, J=5.6 Hz, 1H), 3.52 (s, 2H), 1.8-1.1 (m, 15H), 0.8 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.0$ , 171.8, 170.2, 165.1, 165.0, 161.9, 161.7, 139.6, 139.4, 139.3, 112.8, 112.7, 112.6, 112.5, 103.2, 102.9, 102.6, 82.3, 53.6, 49.3, 43.0, 32.2, 28.4, 27.8, 22.7, 19.4, 14.7, 14.2.

 $C_{21}H_{30}F_2N_2O_4$  (MW = 412.48); mass spectroscopy (MH<sup>+</sup>) 413.

#### Example 11

### Synthesis of N-[N-(Pent-4-enoyl)-L-alaninyl]-L-phenylalanine Methyl Ester

- Following General Procedure A and using N-(L-alaninyl)-L-phenylalanine methyl ester (prepared by coupling N-BOC-L-alanine (Sigma) and L-phenylalanine methyl ester (Sigma) using General Procedure A, followed by removal of the BOC-group using General Procedure Y) and pent-4-enoic acid (Aldrich), the title compound was prepared as a solid (mp = 125.5-126.5°C).
- The reaction was monitored by tlc (Rf = 0.32 in 50% EtOAc/hexanes; 0.51 in 10%  $CH_3OH/CH_2Cl_2$ ) and the product was purified by flash chromatography using 10%  $CH_3OH/CH_2Cl_2$  as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.27$  (bd, J=7.82 Hz, 1H), 7.25-7.05 (m, 5H), 6.72 (bd, J=7.57 Hz, 1H), 5.75 (m, 1H), 4.96 (m, 2H), 4.59 (quint, J=7.2 Hz, 1H), 3.65 (s, 3H), 3.05 (m, 4H), 2.40-2.18 (m, 4H), 1.28 (d, J=7.02 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.06$ , 172.77, 172.36, 137.47, 136.53, 129.76, 129.07, 116.09, 54.10, 52.87, 49.06, 38.31, 25.93, 30.03, 19.17.

30  $C_{18}H_{24}N_2O_4$  (MW = 332.40); mass spectroscopy (MNa<sup>+</sup>) 355.0.

10

20

25

30

#### Example 12

### Synthesis of N-[N-(Dec-4-enoyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure A and using N-(L-alaninyl)-L-phenylalanine methyl ester (prepared by coupling N-BOC-L-alanine (Sigma) and L-phenylalanine methyl ester (Sigma) using General Procedure A, followed by removal of the BOC-group using General Procedure Y) and dec-4-enoic acid (prepare from ethyl dec-4-enoate (ICM) using General Procedure N), the title compound was prepared as a solid (mp = 115.5-117.5°C). The reaction was monitored by tlc (Rf = 0.52 in 50% EtOAc/hexanes; 0.60 in 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by flash chromatography using 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.54$  (bd, J=7.69 Hz, 1H), 7.22-7.04 (m, 5H), 6.91 (bd, J=7.69 Hz, 1H), 5.37 (m, 2H), 4.73 (q, J=6.9 Hz, 1H), 4.63 (quint, J=7.2 Hz, 1H), 3.61 (s, 3H), 3.02 (m, 2H), 2.40-2.10 (m, 4H), 1.89 (m, 2H), 1.35-1.13 (m, 9H), 0.82 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.26$ , 173.05, 172.38, 136.65, 132.30, 129.74, 128.99, 128.68, 127.48, 54.19, 52.74, 48.97, 38.28, 36.70, 33.04, 31.93, 29.68, 29.09, 23.06, 19.23, 14.61.

 $C_{23}H_{34}N_2O_4$  (MW = 402.54); mass spectroscopy (MNa<sup>+</sup>) 425.0.

#### Example 13

#### Synthesis of

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4[3-(N,N-dimethylamino)propoxy]phenylalanine Methyl Ester

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2) and L-4-[3-(N,N-dimethylamino)propoxy]-phenylalanine methyl ester (prepared from N-BOC-L-tyrosine methyl ester (Bachem) and 3-dimethylamino-1-propanol (Aldrich) using a Mitsunobu procedure essentially as described in General Procedure AD, followed by removal of the BOC-group using General Procedure Y), the title compound was

15

20

30

prepared as a solid (mp = 153-155°C). The reaction was monitored by tlc (Rf = 0.36 in 10% MeOH/DCM/1%TEA) and the product was purified by acid/base washes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.973$ -6.947 (d, 2H); 6.794-6.766 (d, 2H); 6.743-6.714 (d, 2H); 6.735-6.676 (t, 1H); 4.761-4.735 (q, 1H); 4.511-4.463 (q, 1H); 3.967-3.924 (t, 2H); 3.703 (s, 3H); 3.473 (s, 2H); 3.019-2.977 (t, 2H); 2.443-2.394 (t, 2H); 2.233 (s, 6H); 1.944-1.897 (t, 2H); 1.319-1.296 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.292$ ; 172.256; 169.808; 158.747; 130.731; 127.887; 115.149; 112.900; 112.672; 66.690; 56.945; 54.039; 52.971; 49.400; 46.105; 43.302; 37.421; 28.129; 19.029.

 $C_{26}H_{33}F_2N_3O_5$  (MW = 505); mass spectroscopy (MH+) 506.

#### Example 14

#### Synthesis of

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4-[(tent-butyloxycarbonyl)methoxy]phenylalanine Methyl Ester

Following General Procedure AE and using *tert*-butyl bromoacetate (Aldrich) and N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester (from Example 15 below), the title compound was prepared as a solid (mp = 116-119°C). The reaction was monitored by tlc (Rf = 0.54 in 50% EtOAc/hexanes) and the product was purified by silica gel column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.648-7.615$  (d, 1H); 7.513-7.407 (d, 1H); 6.943-25 6.914 (d, 2H); 6.756-6.669 (d+t, 4H); 6.621-6.562 (t, 1H); 4.662-4.590 (q+quintex, 2H); 4.382 (s, 2H); 3.571 (s, 3H); 3.406 (s, 2H); 3.006-2.648 (m, 2H); 1.417 (s, 9H); 1.243-1.221 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.14; 173.001; 172.294; 170.273; 168.614; 168.546; 165.107; 161.816; 157.428; 139.493; 130.749; 129.385; 115.077; 112.803; 103.250; 828.270; 66.039; 54.361; 52.730; 49.172; 42.832; 37.288; 28.509; 19.018.

 $C_{27}H_{32}F_2N_2O_7$  (MW = 534); mass spectroscopy (MH<sup>+</sup>) 535.

#### Example 15

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-tyrosine Methyl Ester

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-tyrosine methyl ester (Bachem), the title compound was prepared as a solid (mp = 85-88°C). The reaction was monitored by tlc (Rf = 0.27 in 50% EtOAc/hexanes) and the product was purified by silica gel column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.036$  (b, 1H); 7.369-7.344 (d, 1H); 7.205-7.151 (d, 1H); 6.869-6.841 (d, 2H); 6.763-6.738 (d, 2H); 6.657-6.615 (m, 3H); 4.741-4.697 (q, 1H); 4.566-4.49l (q, 1H); 3.671 (s, 3H); 3.415 (s, 2H); 3.061-2.771 (dm, 2H); 1.271-1.250 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.049; 172.666; 172.444; 170.768; 165.211; 161.917; 156.098, 130.862; 127.542; 116.093, 112.990; 112.659; 103.236; 61.112; 54.306; 49.441; 42.947; 18.923.

 $C_{21}H_{22}F_2N_2O_5$  (MW = 420); mass spectroscopy (MH<sup>+</sup>) 421.

20

25

5

10

15

#### Example 16

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4-(carboxymethoxy)phenylalanine Methyl Ester

Following General Procedure N and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-4-[(tert-butyloxycarbonyl)methoxy]phenylalanine methyl ester (from Example 14 above), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.49 in 10% MeOH/DCM + 1% AcOH) and the product was purified by silica gel column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.817$  (s, 1H); 7.648-7622 (d, 1H); 7.544-7.520 (d, 1H); 6.956-6.914 (d, 2H); 6.762-6.703 (d+d, 4H); 6.650-6.590 (t, 1H); 4.678-4.636 (q, 1H); 4.567-4.503 (quinex + s, 3H); 3.622 (s, 3H); 3.431 (s, 2H); 2.987-2.811 (m, 2H); 1.241-1.219 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.618; 173.534; 172.215; 171.209; 171.108; 165.148; 164.973; 161.855; 161.683; 157.309; 139.052; 130.887; 129.376; 115.104; 112.895; 112.667; 103.083; 65.324; 54.155; 52.933; 50.538; 49.384; 42.683; 37.168; 18.678.

 $C_{23}H_{24}F_2N_2O_7$  (MW = 478); mass spectroscopy (MH<sup>+</sup>) 479.

10

20

25

5

#### Example 17

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)phenylalanine Methyl Ester

Following General Procedure AD and using N-[N-(3,5-

difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester (from Example 15 above) and 4-(2-hydroxyethyl)morpholine (Aldrich), the title compound was prepared as a solid (mp = 138-141°C). The reaction was monitored by tlc (Rf = 0.56 in 10%MeOH/DCM + 1%TEA) and the product was purified by silica gel column chromatography, followed by trituation using diethyl ether.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.974-6.945$  (d, 2H); 6.795-6.726 (d+t, 2H); 6.697-6.682 (t, 1H); 4.755-4.689 (q, 1H); 4.535-4.468 (quintex, 1H); 4.050-4.012 (t, 2H); 3.723-3.606 (t+s, 7H); 3.463 (s, 2H); 3.039-2.892 (m, 2H); 2.779-2.741 (t, 2H); 2.562-2.531 (t, 4H); 1.297-1.274 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 1721.477; 172.428; 172.303; 169.925; 158.397; 130.778; 128.504; 115.179; 112.988; 112.769; 112.659; 67.457; 66.249; 58.187; 54.631; 54.119; 52.956; 49.358; 43.202; 37.496; 19.028.  $C_{27}H_{33}F_2N_3O_6$  (MW = 533); mass spectroscopy (MH<sup>+</sup>) 534.

#### Example 18

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-6-(N,N-dimethylamino)hexanoate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and Nε,Nε-dimethyl-L-lysine methyl ester hydrochloride (Bachem), the title compound was prepared as a solid (mp = 123-126°C). The reaction was monitored by tlc (Rf = 0.22 in 10% MeOH/DCM + 1% TEA) and the product was purified by silica gel column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.019$ -6.993 (d, 1H); 6.828-6.801 (dd, 2H); 6.753-6.723 (m, 1H); 6.617-6.592 (d, 1H); 4.557-4.447 (q+q, 2H); 3.730 (s, 3H); 3.522 (s, 2H); 2.593-2.572 (m, 2H); 2.196 (s, 6H); 1.837-1.642 (m, 2H); 1.486-1.344 (m+d, 7H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.070$ ; 172.544, 169.809; 112.986; 112.655; 103.384: 59.393; 52.991; 49.368; 45.947; 43.427; 43.403; 43.375; 31.870; 27.376; 23.378; 19.155.

 $C_{20}H_{29}F_2N_3O_4$  (MW = 413); mass spectroscopy (MH<sup>+</sup>) 414.

20

15

#### Example 19

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-pyridyl)propionate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L25 alanine (from Example B2) and methyl (S)-2-amino-3-(2-pyridyl)propionate
hydrochloride (Synthetech), the title compound was prepared as a solid (mp =
121-124°C). The reaction was monitored by tlc (Rf = 0.39 in
10%MeOH/DCM) and the product was purified by silica gel column
chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.474-8.458$  (d, 1H); 7.767-7.631 (m, 1H); 7.625-7.574 (t, 1H); 7.178-7.102 (t+d, 2H); 6.818-6.811 (d, 2H); 6.734-6.667 (t, 1H); 6.593-6.542 (m, 1H); 4.933-4.873 (m, 1H); 4.566-4.496 (m, 1H); 3.646 (s, 3H); 3.499 (s, 2H); 3.375-3.196 (m, 2H); 1.393-1.370 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.453; 172.020; 169.527; 157.454; 149.608; 137.449; 124.366; 124.328; 122.694; 113.032; 112.992; 112.661; 103.333; 53.032; 52.997; 52.349; 52.252; 49.427; 49.405; 43.464, 43.437, 38.486; 19.548; 19.232.

 $C_{20}H_{21}F_2N_3O_4$  (MW = 405); mass spectroscopy (MH<sup>+</sup>) 406.

10

15

20

25

5

#### Example 20

## Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl](S)-2-amino-3-(3-pyridyl)propionate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2) and methyl (S)-2-amino-3-(3-pyridyl)propionate hydrochloride (Synthetech), the title compound was prepared as a solid (mp = 101-103°C). The reaction was monitored by tlc (Rf = 0.48 in 10% MeOH/DCM) and the product was purified by silica gel column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.492-8.396$  (m, 1H); 8.359-8.322 (m, 1H); 7.505-7.452 (m, 1H); 7.248-7.170 (m, 1H); 6.976-6.908 (m, 1H); 6.855-6.668 (m, 3H); 6.352-6.288 (m, 1H); 4.866-4.798 (m, 1H); 4.784-4.429 (m, 1H); 3.750 (s, 3H); 3.513 (s, 2H); 3.220-2.964 (m, 2H); 1.310-1.287 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.867; 171.831; 170.307; 161.942; 150.892; 150.753; 148.907; 148.750; 137.523; 137.388; 132.460; 124.106; 124.034; 112.981; 112.754; 103.228; 53.623; 53.461; 53.146; 49.368; 49.259; 43.137; 43.115; 43.086; 35.485; 18.664.

 $C_{20}H_{21}F_2N_3O_4$  (MW = 405); mass spectroscopy (MH<sup>+</sup>) 406.

#### Example 21

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]L-proline Methyl Ester

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-proline methyl ester hydrochloride (Bachem), the title compound was prepared as a viscous solid. The reaction was monitored by tlc (Rf = 0.57 in 10% MeOH/DCM) and the product was purified by acid/base washes.

10 NMR data was as follows:

15

20

25

30

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.524-7.498$  (d, 1H); 6.813-6.793 (d, 2H); 6.681-6.613 (m, 1H); 4.788-4.717 (m, 1H); 4.484-4.442 (m, 1H); 3.705-3.590 (m+s, 4H); 3.465 (s, 2H); 2.217-1.902 (m, 5H); 1.332-1.309 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.753$ ; 172.152; 169.843; 165.185; 161.894; 112.953; 112.850; 112.727; 112.624; 103.331, 102.996, 102.662; 59.352; 52.735; 47.495; 47.267; 43.069; 29.472; 25.403; 18.243.

 $C_{17}H_{20}F_2N_2O_4$  (MW = 354); mass spectroscopy (MH<sup>+</sup>) 355.

#### Example 22

# Synthesis of Methyl 1-[N-(3,5-Difluorophenylacetyl)L-alaninyl]piperidine-2-carboxylate

Following General Procedure B and using N-(phenylacetyl)-L-alanine (from Example B1 above) and methyl pipecolinate hydrochloride (Aldrich), the title compound was prepared as an oil. The reaction was monitored by tlc (Rf = 0.30 in 50% EtOAc/hexanes) and the product was purified by silica gel chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.2$  (m, 5H), 6.95 (dd, J=7.2, 15.2, 7.2 Hz, 1H), 5.21 (dd, J=5.0, 11.0, 5.0 Hz, 1H), 4.89 (q, J=7.1, 7.1 Hz, 1H), 3.7 (m, 1H), 3.59 (s, 3H), 3.47 (s, 2H), 3.1 (m, 1H), 2.16 (d, J=11.5 Hz, 1H), 1.4

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.6, 171.8, 170.7, 135.5, 129.8, 129.3, 127.6, 52.9, 52.8, 46.0, 43.9, 27.1, 26.8, 25.6, 21.4, 19.9, 18.5. C<sub>18</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> (MW = 332); mass spectroscopy (MH<sup>+</sup>) 333.

#### Example 23

5

10

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-(S)-2-amino-3-(4-pyridyl)propionate

#### Step A - Preparation of 3-(3-pyridyl)alanine methyl ester dihydrochloride

Sodium metal (1.40 g, 61 mmol) was dissolved in EtOH (100mL) and diethyl acetamidomalonate (6.62 g, 30.5 mmol) and 3-picolylchloride hydrochloride (5.00g, 30.5 mmol) were added. The mixture was heated to reflux for 6 hours, and then cooled and filtered to remove NaCl (washed with EtOH). The solvent was removed *in vacuo* and the mixture was taken up into saturated aqueous NaHCO<sub>3</sub> (100 mL) and extracted with EtOAc (3 x 100 m

L).

The solvent was removed and the residue purified by silica gel flash chromatography (95:5 CH<sub>2</sub>Cl<sub>2</sub>/MeOH) to give diethyl 2-(3-pyridylmethyl)-2-acetamidomalonate (2.84 g, 30%).

Diethyl 2-(3-pyridylmethyl)-2-acetamidomalonate was dissolved in 6N HCl (30 mL) and heated to reflux for 19 hours whereupon it was cooled to room

10

temperature and the HCl solution was removed by evaporation *in vacuo*. The intermediate amino acid dihydrochloride salt was taken up into MeOH (30 mL) saturated with HCl gas and stirred for 3.5 hours. The MeOH/HCl was removed by evaporation *in vacuo* to give 3-(3-pyridyl)alanine methyl ester dihydrochloride (2.235 g, 100%).

# Step B - Preparation of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4-pyridyl)propionate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (S)-2-amino-3-(4pyridyl)propionate hydrochloride (prepared by the method set forth above using

4-picolylchloride hydrochloride (prepared by the method set forth above using 4-picolylchloride hydrochloride), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.49 in 10% MeOH/DCM) and the product was purified by silica gel column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 8.423-8.335 (dd, 2H); 7.832-7.754 (q, 1H); 7.342-7.246 (dd, 1H); 7.032-6.972 (dd, 2H); 6.764-6.667 (t, 2H); 6.659-6.599 (m, 1H); 4.837-4.768 (m, 1H); 4.590-4.515 (m, 1H); 3.675 (s, 3H); 3.426 (s, 2H); 3.112-2.804 (m, 2H); 1.256-1.106 (dd, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.037$ ; 171.739; 170.258; 170.225; 165.201; 165.012; 161.904; 161.721; 150.183; 150.063; 146.115; 146.012; 139.100; 125.180; 125.122; 112.951; 112.915; 112.846; 103.492; 103.153; 53.088, 49.318; 42.977, 37.593; 37.547; 19.297; 18.882.

 $C_{20}H_{21}F_2N_3O_4$  (MW = 405); mass spectroscopy (MH<sup>+</sup>) 406.

#### Example 24

25

30

20

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-3-methoxypropionate hydrochloride (Bachem), the title compound was prepared as a solid (mp =

10

15

20

25

165-168°C). The reaction was monitored by tlc (Rf = 0.48 in 10% MeOH/DCM) and the product was purified by acid/base washes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.971$ -6.944 (d, 1H); 6.813-6.801 (m, 2H); 6.741-6.678 (m, 1H); 6.585-6.526 (m, 1H); 4.692-4.561 (quintex + q, 2H); 3.836-3.802 (m, 1H); 3.738 (s, 3H); 3.592-3.516 (m+ds, 3H); 3.312 (s, 3H); 1.408-1.355 (dd, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.705$ ; 172.680; 170.908; 113.019; 112.978; 112.687; 112.646; 103.347; 72.434; 72.405; 59.885; 59.837; 53.263; 53.240; 49.413; 49.329; 19.389; 18.9196.

 $C_{16}H_{20}F_2N_2O_5$  (MW = 358); mass spectroscopy (MH<sup>+</sup>) 359.

#### Example 25

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-3-morpholinopropionate

#### Step A methyl (2-N-CBZ-amino)-3-morpholino-propionate

To a solution of N-CBZ-dehydro-alanine methyl ester (Sigma) in acetonitrile was added 2.0 equivalent of morpholine and 0.25 equivalents of anhydrous ferric chloride. The mixture was stirred for 16 hours and monitored by TLC. The solvent was stripped off and the residue extracted with ethyl acetate and washed with 1N HCl. The aqueous layer was basified with 1N potassium carbonate to pH = 9 and extracted with ethyl acetate again, dried over sodium sulfate and rotovapped to dryness to give methyl (2-N-CBZ-amino)-3-morpholino-propionate as a clear tan oil. See Perez et al., Tetrahedron 51(3) 8355-62 (1995)

Step B Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-3-morpholinopropionate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-

10

L-alanine (from Example B2 above) and methyl 2-amino-3-morpholinopropionate hydrochloride (prepared by General Procedure O above from methyl (2-N-CBZ-amino)-3-morpholino-propionate), the title compound was prepared as a viscous solid. The reaction was monitored by tlc (Rf = 0.44 in 10% MeOH/DCM) and the product was purified by acid/base washes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.408-7.384$  (d, 1H); 7.247-7.173 (m, 1H); 6.774-6.614 (m+t, 3H); 4.605-4.468 (m, 1H); 3.667 (s, 3H); 3.642 (s, 2H); 3.576-3.561 (t, 4H); 3.479-3.461 (s+s, 2H); 2.639-2.618 (d, 2H); 2.395-2.366 (m, 4H); 1.344-1.307 (t, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.120; 172.245; 172.192; 170.275; 170.159; 165.189; 165.020; 161.897; 161.727; 139.167; 112.937; 112.863; 112.759; 112.610; 112.533; 103.103; 102.774; 67.379; 67.301; 59.346; 59.110; 54.030; 52.936; 51.116; 49.283; 43.053; 18.980; 18.921.

15  $C_{19}H_{26}F_2N_3O_5$  (MW = 413); mass spectroscopy (MH<sup>+</sup>) 414.

#### Example 26

# Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)phenylalaninamide

Following General Procedure K and using 2-methoxyethylamine (Aldrich) and N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)-phenylalanine methyl ester (from Example 17 above), the title compound was prepared as a solid (mp = 165-168°C). The reaction was monitored by tlc (Rf = 0.67 in 10% MeOH/DCM+1% TEA) and the product was purified by acid/base washes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 8.258-8.232 (d, 1H); 8.014-7.989 (d, 1H); 7.532-7.370 (t, 1H); 7.035-7.008 (d, 2H); 6.842-6.630 (m, 5H); 4.980-4.905 (m, 1H); 4.794-4.772 (m, 1H); 4.026-3.992 (t, 2H); 3.713-3.642 (t, 4H); 3.594-3.453 (dd, 2H); 3.404-3.267 (t, 2H); 3.179 (s, 3H); 2.930-2.914 (t, 2H); 2.763-2.731 (t, 2H); 2.538-2.502 (m, 4H); 1.335-1.314 (d, 3H).

15

20

25

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.956; 172.918; 171.756; 170.142; 161.677; 158.131; 130.973; 129.270; 114.968; 114.875; 112.908; 112.696; 112.571; 71.423; 71.367; 67.440; 66.164; 59.072; 58.232; 58.188; 54.636; 42.827; 42.800; 39.757; 39.642; 20.449; 20.135.

 $C_{29}H_{38}F_2N_4O_6$  (MW = 576); mass spectroscopy (MH<sup>+</sup>) 577.

#### Example 27

## Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionamide

Following General Procedure K and using 2-methoxyethylamine (Aldrich) and methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionate (from Example 24 above), the title compound was prepared as a solid (mp = 181-184°C). The reaction was monitored by tlc (Rf = 0.43 in 10% MeOH/DCM) and the product was purified by acid/base washes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.728$ -6.706 (d, 2H); 6.648-6.586 (t, 1H); 4.244-4213 (m, 1H); 4.092-4.068 (m, 1H); 3.553-3.503 (m, 2H); 3.393-3.347 (m, 2H); 3.210-3.073 (m+s, 7H); 3.053 (s, 3H); 1.183-1.138 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 176.31; 173.28; 172.59; 141.65; 114.02; 113.79; 113.69; 109.467; 103.528; 80.369; 73.210; 72.265; 72.011; 59.839; 59.801; 59.374; 55.584; 51.773; 51.731; 51.445; 42.915; 40.846; 17.751.

 $C_{18}H_{25}F_2N_3O_5$  (MW = 401); mass spectroscopy (MH<sup>+</sup>) 402.

#### Example 28

#### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]glycine Methyl Ester

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and glycine methyl ester hydrochloride (Bachem), the title compound was prepared as a solid (mp = 158-160°C). The

WO 98/22494 PCT/US97/20804

-- 201 --

reaction was monitored by tlc (Rf = 0.61 in 10%MeOH/DCM) and the product was purified by silica gel chromatography.

NMR data was as follows:

5

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.882-6.866$  (m, 1H); 6.827-6.794 (m, 2H); 6.748-6.7486.689 (t, 1H); 6.520-6.494 (d, 1H); 4.611-4.563 (quintex, 1H); 4.00-3.99 (d, 2H); 3.746 (s, 3H); 3.528 (s, 2H); 1.389-1.366 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.926$ ; 172.524; 170.524; 113.056; 112.951; 112.723; 103.769; 103.437; 103.214; 103.105; 85.309; 53.009; 49.333; 43.292; 41.692; 18.810.

10  $C_{14}H_{16}F_2N_2O_4$  (MW = 314); mass spectroscopy (MH<sup>+</sup>) 315.

#### Example 29

#### Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(4-pyridyl)propionamide

15 Following General Procedure K and using 2-methoxyethylamine and methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4-pyridyl)propionate (from Example 23 above), the title compound was prepared as a solid (mp = 202-206°C). The reaction was monitored by tlc (Rf = 0.72 in 10%MeOH/DCM) and the product was purified by acid/base washes.

20 NMR data was as follows:

> <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.214-8.198$  (d, 2H); 7.117-7.100 (d, 2H); 6.707-6.687 (m, 2H); 6.638-6.576 (t, 1H); 4.498-4.448 (m, 1H); 3.985-3.939 (q, 1H); 3.386 (s, 2H); 3.190-3.084 (m, 4H); 3.060 (s, 3H); 2.918-2.629 (m, 2H); 1.077-0.905 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 175.831$ ; 173.229; 150.440; 150.249; 126.887; 25 113.995; 113.662; 103.662; 103.529; 72.081; 59.370; 55.201; 51.674; 42.949; 40.889; 38.350; 17.933.

 $C_{22}H_{26}F_2N_4O_4$  (MW = 448); mass spectroscopy (MH<sup>+</sup>) 449.

10

15

20

#### Example 30

# Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(2-pyridyl)propionamide

Following General Procedure K and using 2-methoxyethylamine and methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-pyridyl)propionate (from Example 19 above), the title compound was prepared as a solid (mp = 183-187°C). The reaction was monitored by tlc (Rf = 0.39 in 10% MeOH/DCM) and the product was purified by recrystallization from MeOH/DCM.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.457-8.442$  (d, 1H); 8.029-8.005 (d, 1H); 7.642-7.585 (t, 1H); 7.395-7.379 (m, 1H); 7.267-7.141 (d+t, 2H); 6.828-6.802 (m, 2H); 6.754-6.679 (t, 1H); 6.604-6.581 (m, 1H); 4.871-4.809 (q, 1H); 4.532-4.485 (quintex, 1H); 3.537 (s, 2H); 3.342-3.118 (m, 6H); 3.248 (s, 3H); 1.394-1.371 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.360; 171.140; 158.43; 149.113; 137.59; 124.98; 122.54; 113.02; 112.69; 103.40; 71.376; 59.203; 53.143; 49.984; 43.355; 43.328; 39.685; 39.626; 19.295.

 $C_{22}H_{26}F_2N_4O_4$  (MW = 448); mass spectroscopy (MH<sup>+</sup>) 449.

#### Example 31

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-yl)propionate

25 Following General Procedure A and using N-(3,5-difluorophenylacetyl)L-alanine (from Example B2 above) and methyl (S)-2-amino-3-(thiazol-4yl)propionate hydrochloride (General Procedure H with methanol and HCl on
methyl (S)-2-amino-3-(thiazol-4-yl)propyl acid (Synthetech)), the title compound
was prepared as a solid (mp = 136-139°C). The reaction was monitored by tlc

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.737-8.731$  (d, 1H); 7.410-7.385 (d, 1H); 7.065-7.059 (d, 1H); 6.828-6.802 (m, 2H); 6.747-6.687 (m, 1H); 6.542-6.518 (d, 1H); 4.904-4.844 (q, 1H); 4.553-4.505 (quintex, 1H); 3.678 (s, 3H); 3.515 (s, 2H); 3.402-3.232 (dq, 2H); 1.384-1.361 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.497$ ; 171.726; 169.619; 153.831; 152.613; 116.431; 113.019; 112.688; 112.014; 103.396; 53.113; 52.625; 49.476; 43.460; 43.435; 32.850; 19.422.

 $C_{18}H_{19}F_2N_3O_4S$  (MW = 411); mass spectroscopy (MH<sup>+</sup>) 412.

10

25

5

#### Example 32

## Synthesis of Methyl 2-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-1,2,3,4-tetrahydroisoguinoline-3-carboxylate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)
L-alanine (from Example B2 above) and methyl 1,2,3,4-tetrahydroisoquinoline3-carboxylate (Aldrich), the title compound was prepared as a solid (mp = 3740°C). The reaction was monitored by tlc (Rf = 0.64 in 10% MeOH/DCM).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.500$ -7.475 (d, 1H); 7.161-7.057 (m, 4H); 6.815-20 6.795 (dm, 2H); 6.656-6.596 (t, 1H); 5.336-5.088 (m, 2H); 4.924-4.841 (m, 1H); 4.718-4.453 (m, 1H); 3.530 (s, 3H); 3.500 (s, 2H); 3.329-3.058 (m, 2H); 1.423-1.400, 1.327-1.304 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.428; 173.329; 171.690; 171.559; 169.558; 165.020; 161.899; 161.728; 139.368; 132.549; 128.912; 127.723; 126.648; 112.929; 103.360; 60.915; 53.318; 53.001; 46.377; 43.121; 31.027; 21.537; 19.545; 18.771; 14.716.

 $C_{22}H_{22}F_2N_2O_4$  (MW = 416); mass spectroscopy (MH<sup>+</sup>) 417.

25

30

#### Example 33

## Synthesis of N-(3-Methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure B and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine (prepared by coupling N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) with L-phenylalanine methyl ester hydrochloride (Sigma) using General Procedure E, followed by hydrolysis using General Procedure C) and 3-methoxybenzylamine (TCI), the title compound was prepared as a solid (mp = 117-130°C). The reaction was monitored by the (Rf = 0.8 in 3% MeOH/methylene chloride) and the product was purified by recrystallization from MeOH.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.4$  (t, 1H), 8.32 (d, 1H), 8.1 (d, 1H), 6.95-7.2 (m, 9H), 6.7 (m, 3H), 4.5 (m, 1H), 4.2 (m, 3H), 3.7 (s, 3H), 3.5 (s, 2H), 3.3 (d, 2H), 3.0 (m, 2H), 2.5 (s, 3H), 1.2 (m, 4H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.40$ , 171.08, 169.28, 159.62, 141.09, 138.06, 129.62, 129.51, 128.41, 126.63, 119.56, 112.97, 112.79, 112.59, 112.46, 55.31, 48.77, 40.69, 40.42, 40.28, 40.14, 40.03, 39.86, 39.70, 39.58, 39.46, 39.44, 39.31, 39.20, 39.03, 18.45.

 $C_{28}H_{29}N_3O_4F_2$  (MW = 509); mass spectroscopy (MH<sup>+</sup>) 509.

#### Example 34

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-(S)-2-amino-3-(1-naphthyl)propionate

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (S)-2-amino-3-(1-naphthyl)propionate hydrochloride (Bachem), the title compound was prepared as a solid (mp = 103-130°C). The reaction was monitored by tlc (Rf = 0.8 in 5% MeOH/methylene chloride) and the product was purified by flash column chromatography using 6% MeOH/methylene chloride as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.10$  (d, 1H), 7.85 (d, 1H), 7.71 (d, 1H), 7.50 (m, 3H), 7.35 (t, 1H), 7.20 (d, 1H), 6.70 (m, 4H), 6.30 (d, 1H), 4.90 (m, 1H), 4.45 (m, 1H), 3.3-3.7 (m, 8H), 1.7 (bs, 1H), 1.3 (d 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.43, 172.29, 169.77, 134.41, 132.61, 132.58, 129.51, 128.63, 128.33, 128.28, 128.06, 126.97, 126.80, 126.42, 126.29, 125.94, 125.86, 124.06, 123.90, 112.96, 112.63, 103.44, 78.03, 77.61, 77.19, 61.01, 54.02, 53.83, 52.99, 51.40, 49.33, 43.29, 35.64, 18.82, 14.77.

 $C_{24}H_{24}N_2O_4F_2$  (MW = 442); mass spectroscopy (MH<sup>+</sup>) 442.

10

15

20

25

5

#### Example 35

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-(S)-2-amino-3-(2-naphthyl)propionate

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (S)-2-amino-3-(2-naphthyl)propionate hydrochloride (Bachem), the title compound was prepared as a solid (mp =  $166^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.55 in 5% MeOH/methylene chloride) and the product was purified by preparative tlc using 5% MeOH/methylene chloride as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 1.3$  (d, 3H), 3.2 (m, 2H), 3.3 (s, 2H), 3.7 (s, 3H), 4.55 (m, 1H), 4.9 (quart, 1H), 6.7 (m, 4H), 7.05 (d, 1H), 7.20 (d, 1H), 7.45 (m, 2H), 7.55 (s, 1H), 7.80 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.43$ , 172.26, 169.86, 133.93, 133.76, 133.02, 128.86, 128.64, 128.23, 128.20, 127.69, 126.85, 126.45, 112.95, 112.62, 103.37, 78.05, 77.62, 77.20, 53.93, 53.05, 49.37, 43.12, 38.46, 18.81.  $C_{24}H_{24}N_2O_4F_2$  (MW = 442); mass spectroscopy (MH<sup>+</sup>) 442.

#### Example 36

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-(S)-2-amino-3-(2-thienyl)propionate

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (S)-2-amino-3-(2-thienyl)-propionate (Bachem), the title compound was prepared as a solid (mp = 145-147°C). The reaction was monitored by tlc (Rf = 0.9 in 100% EtOAc) and the product was purified by preparative tlc using EtOAc as the eluent.

NMR data was as follows:

20

25

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.15$  (d, 1H), 6.9 (t, 1H), 6.7-6.8 (m, 5H), 6.3 (d, 1H), 4.8 (m, 1H), 4.5 (m, 1H), 3.8 (s, 3H), 3.5 (s, 2H), 3.35 (d, 2H), 1.35 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.22$ , 171.56, 169.79, 137.47, 127.71, 125.55, 113.04, 112.71, 103.48, 78.03, 77.60, 77.18, 53.78, 53.25, 49.51, 43.41, 32.37, 18.97.

 $C_{19}H_{20}N_2O_4F_2S$  (MW = 410); mass spectroscopy (MH<sup>+</sup>) 410.

#### Example 37

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalanine Benzyl Ester

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylalanine benzyl ester hydrochloride (Bachem), the title compound was prepared as a solid (mp = 170-171°C). The reaction was monitored by tlc (Rf = 0.7 in 5% MeOH/methylene chloride) and the product was purified by recrystallization from MeOH.

NMR data was as follows:

<sup>1</sup>H-nmr (MeOH):  $\delta = 7.3$  (m, 10H), 6.9 (m, 3H), 5.2 (s, 2H), 4.75 (t, 30 J=7 Hz, 1H), 4.4 (quart, J=6 Hz, 1H), 3.6 (s, 2H), 3.1 (m, J= 6 Hz, 2H), 1.35 (d, J=7 Hz, 3H).

<sup>13</sup>C-nmr (MeOH):  $\delta$  = 175.29, 173.09, 172.78, 141.54, 138.35, 137.53, 130.88, 130.08, 130.05, 129.92, 128.42, 113.93, 113.83, 113.60, 103.90, 103.55, 103.21, 68.59, 55.87.

 $C_{27}H_{26}N_2O_4F_2$  (MW = 480); mass spectroscopy (MH<sup>+</sup>) 480.

5

10

15

20

25

#### Example 38

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-Bromopropyl Ester

Following General Procedure B and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine (prepared by coupling N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylalanine (Aldrich) using General Procedure B) and 3-bromo-1-propanol (Aldrich), the title compound was prepared as a solid (mp = 138-142°C). The reaction was monitored by the (Rf = 0.75 in 60% EtOAc/hexanes) and the product was purified by flash column chromatography using 60% EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.3$ -6.6 (m, 10H), 4.8 (m, 1H), 4.55 (m, 1H), 4.2 (t, J=6 Hz, 2H), 3.51 (s, 2H), 3.3 (m, 2H), 3.05 (m, J=6 and 8 Hz, 2H), 2.1 (m, 2H), 1.3-1.2 (m, J=7 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.49, 171.78, 171.71, 170.01, 169.96, 165.31, 162.02, 161.84, 138.91, 138.78, 138.66, 136.26, 136.19, 129.76, 129.72, 129.22, 129.18, 127.80, 113.04, 113.02, 112.93, 112.91, 112.82, 112.79, 112.71, 112.69, 103.72, 103.69, 103.36, 103.05, 103.03, 63.75, 63.70, 54.11, 53.91, 49.38, 49.32, 43.26, 38.56, 38.51, 31.92, 29.76, 29.71, 19.14, 19.06.

 $C_{23}H_{25}N_2O_4F_2Br$  (MW = 511.1); mass spectroscopy (MH<sup>+</sup>) 512.

#### Example 39

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-Iodopropyl Ester

15

Following General Procedure B and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine (prepared by coupling N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylalanine (Aldrich) using General Procedure B) and 3-iodo-1-propanol (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.45 in 5% MeOH/methylene chloride) and the product was purified by preparative tlc using 5% MeOH/methylene chloride.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.4$ -7.0 (m, 5H), 6.9-6.6 (m, 4H), 6.3 (m, 1H), 4.8 (m, 1H), 4.5 (m, 1H), 4.2 (t, 2H), 3.5 (s, 2H), 3.1 (m, 4H), 2.1 (m, 2H), 1.7 (s, 1H), 1.35-1.25 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.24$ , 171.72, 169.95, 136.12, 136.09, 129.77, 129.75, 129.28, 129.24, 127.87, 113.06, 113.02, 112.73, 112.70, 103.80, 103.49, 103.47, 65.73, 65.70, 54.00, 53.84, 49.42, 49.33, 43.38, 38.54, 38.50, 32.57, 18.97, 18.91.

#### Example 40

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-leucine tert-Butyl Ester

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and L-leucine tert-butyl ester hydrochloride
(Bachem), the title compound was prepared as a solid (mp = 128°C). The
reaction was monitored by tlc (Rf = 0.85 in 5% MeOH/methylene chloride)
and the product was purified by flash column chromatography using 5%
MeOH/methylene chloride as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.9$ -6.5 (m, 5H), 4.6 (m, 1H), 4.4 (m, 1H), 3.5 (s, 2H), 1.7-1.4 (m, 15H), 0.9 (t, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.41$ , 172.20, 169.87, 165.30, 162.00, 161.83, 139.01, 138.89, 112.92, 112.82, 112.69, 112.59, 103.62, 103.29, 102.95,

82.50, 78.03, 77.61, 77.18, 52.12, 49.39, 43.34, 41.86, 28.52, 25.42, 23.26, 22.46, 19.18.

 $C_{27}H_{30}N_2O_4F_2$  (MW = 412.48); mass spectroscopy (MH<sup>+</sup>) 413.

#### Example 41

5

10

15

# Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetamide

Following General Procedure L and using ethyl N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetate (from Example 65 below), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.1 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by recrystallization from EtOH.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.54$  (m, 1H), 8.43 (d, 1H), 7.77 (m, 1H), 7.59 (bs, 1H), 7.46 (m, 1H), 7.33 (m, 1H), 7.22 (m, 1H), 7.09 (m, 1H), 6.98 (m, 2H), 5.41 (m, 1H), 4.46 (m, 1H), 4.46 (m, 1H), 3.52 (s, 2H), 1.26 (m, 3H).  $C_{18}H_{17}N_3O_3F_2$  (MW = 376.3); mass spectroscopy (MH<sup>+</sup>) 377.

#### Example 42

20

25

30

# Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetamide

Following General Procedure L and using ethyl N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetate (from Example 53 below), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.1 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by recrystallization from EtOH.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.64$  (m, 1H), 8.55 (d, 1H), 8.52 (d, 1H), 8.41 (d, 1H), 7.79 (m, 1H), 7.37 (m, 1H), 7.32 (m, 1H), 7.09 (m, 1H), 6.98 (m, 2H), 5.42 (m, 1H), 4.42 (m, 1H), 3.53 (s, 2H), 1.26 (m, 3H).

10

15

 $C_{18}H_{17}N_4O_3F_2$  (MW = 376.3); mass spectroscopy (MH<sup>+</sup>) 377.

#### Example 43

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]- $N_c$ -(tent-butoxycarbonyl)-L-lysine Methyl Ester

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and  $N\epsilon$ -(tert-butoxycarbonyl)-L-lysine methyl ester (Bachem), the title compound was prepared as an oil. The reaction was monitored by tlc (Rf = 0.40 in 50% EtOAc/hexanes) and the product was purified by flash chromotography using 50% EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.80$  (d, 2H), 6.66 (t, 1H), 4.82 (bs, 1H), 3.73 (s, 3H), 3.52 (s, 2H), 3.04 (bs, 2H), 1.60-1.15 (m, 2H), 1.38 (s, 9H), 1.32 (d, 2H), 1.20-1.30 (m, 2H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.00$ , 172.80, 165.28, 165.11, 161.98, 161.78, 156.79, 138.95, 129.06, 128.72, 103.59, 103.26, 102.92, 79.81, 52.99, 52.76, 49.44, 43.25, 31.92, 29.98, 28.99, 22.95, 18.94.

 $C_{23}H_{33}F_2N_3O_6$  (MW = 485.53); mass spectroscopy (MH<sup>+</sup>) N/A.

#### Example 44

20

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-(S)-2-amino-4-phenylbutanoate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and methyl (S)-2-amino-4-phenylbutanoate

(prepared from (+)-α-amino-4-phenylbutyric acid (Bachem) using General

Procedure AG), the title compound was prepared as a solid (mp = 147149.5°C). The reaction was monitored by tlc (Rf = 0.32 in 50%

EtOAc/hexanes) and the product was purified by flash chromotography using

EtOAc/hexanes as the eluent.

NMR data was as follows:

WO 98/22494 PCT/US97/20804

-- 211 --

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.63$  (bd, 2H), 7.04 (m, 5H), 6.56-6.82 (m, 3H), 4.80 (p, 1H), 4.48 (q, 1H), 3.65 (s, 3H), 3.49 (s, 2H), 2.50-2.65 (m, 2H), 1.80-2.16 (m, 2H), 1.29 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.48$ , 172.89, 170.43, 165.17, 161.71, 140.91, 139.34, 129.07, 129.01, 128.89, 126.81, 126.76, 112.90, 112.67, 103.37, 103.03, 102.69, 52.86, 52.71, 49.36, 42.99, 33.79, 32.21, 19.34.

 $C_{22}H_{24}F_2N_2O_4$  (MW = 418.44); mass spectroscopy (MH<sup>+</sup>) 419.

#### Example 45

10

15

25

5

#### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]glycine 2-Phenylethyl Ester

Following General Procedure X and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine benzyl ester (from Example 73 below) using General Procedure O) and 2-phenylethanol (Aldrich), the title compound was prepared as a solid (mp = 154.0-155.2°C). The reaction was monitored by the (Rf = 0.15 in 15% EtOAc/hexanes) and the product was purified by flash chromotography using 15% EtOAc/hexanes as the eluent.

NMR data was as follows:

20 <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 7.35-7.20 (m, 5H), 6.76 (bs, 1H), 6.72-6.67 (m, 3H), 6.54 (bd, 1H), 4.58 (p, 1H), 4.34 (t, 2H), 3.96 (d, 2H), 3.52 (s, 2H), 2.93 (t, 2H), 1.26 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.9, 170.1, 169.9, 137.8, 129.4, 129.1, 127.3, 112.94, 103.4, 103.0, 65.5, 49.3, 43.2, 41.8, 35.4, 18.8.

 $C_{21}H_{22}N_2O_4F_2$  (MW = 404.42); mass spectroscopy (MH<sup>+</sup>) 405.

#### Example 46

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]glycine 3-Phenylpropyl Ester

Following General Procedure X and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine benzyl ester (from Example 73 below) using General Procedure O) and and 3-phenyl-1-propanol (Aldrich), the title compound was prepared as a solid (mp = 137°C). The reaction was monitored by tlc (Rf = 0.15 in 50% EtOAc/hexanes) and the product was purified by flash chromotography using 50% EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.55-7.32$  (m, 5H), 6.73 (d, 2H), 6.65 (m, 1H), 4.74 (p, 1H), 4.14 (t, 2H), 3.93 (m, 2H), 3.49 (s, 2H), 2.66 (t, 2H), 1.94 (p, 2H), 1.41 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 173.8, 170.5, 170.1, 165.2, 165.0, 161.9, 161.7, 141.5, 139.2, 129.1, 128.9, 126.7, 112.9, 112.8, 103.4, 103.1, 102.8, 65.4, 49.3, 42.9, 41.8, 32.6, 30.6, 19.3.

15  $C_{22}H_{24}N_2O_4F_2$  (MW = 418.44); mass spectroscopy (MH<sup>+</sup>) 419.

#### Example 47

# Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetamide

- Following General Procedure L and using ethyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetate (from Example 66), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.1 in CHCl<sub>3</sub>/MeOH 9:1) and the product was purified by silica gel chromatography using 9:1 CHCl<sub>3</sub>/MeOH as the eluent.
- NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.53$  (m, 2H), 8.88 (bs, 1H), 7.41 (m, 2H), 7.12 (m, 1H), 7.02 (m, 2H), 5.46 (m, 1H), 4.46 (m, 1H), 3.55 and 3.52 (s, 2H), 1.21 (m, 3H).

 $C_{18}H_{18}N_4O_3F_2$  (MW = 376.3); mass spectroscopy (MH<sup>+</sup>) 377.

#### Example 48

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-threonine Methyl Ester

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-threonine methyl ester hydrochloride (Bachem), the title compound was prepared as a solid.

NMR data was as follows:

67.5, 57.7, 52.1, 48.7, 42.8, 19.6, 18.3.

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 7.45 (d, J=8.9 Hz, 1H), 7.11-7.27 (m, 6H), 4.55 (quintet, J=7.2 Hz, 1H), 4.43 (dd, J=2.6,8.8 Hz, 1H), 4.20 (m, 1H), 3.62 (s, 3H), 3.46 (s, 2H), 1.29 (d, J=7.0 Hz, 3H), 1.04 (d, J=6.4 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.8, 171.1, 170.9, 134.5, 128.9, 128.4, 126.8,

 $C_{16}H_{22}N_2O_5$  (MW = 322.36); mass spectroscopy (MH<sup>+</sup>) 323.

#### Example 49

15

20

10

### $Synthesis of \\ N'-[N-(Phenylacetyl)-L-alaninyl]-L-leucinamide$

Following General Procedure T and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-leucinamide hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 207-209°C). The product was purified by extraction with EtOAc and washing with aqueous potassium carbonate and aqueous hydrochloric acid.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 7.00-7.12$  (m, 5H), 4.10-4.20 (m, 2H), 3.34 (s, 2H), 1.30-1.50 (m, 2H), 1.12-1.23 (m, 4H), 0.65-0.76 (m, 6H).

25  $^{13}$ C-nmr (CD<sub>3</sub>OD):  $\delta = 177.5$ , 174.9, 174.1, 136.8, 130.1, 129.6, 127.9, 52.8, 50.7, 43.4, 41.9, 25.8, 23.5, 21.8, 17.7.

#### Example 50

### Synthesis of N'-[N-(Phenylacetyl)-L-alaninyl]-L-alaninamide

10

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-alaninamide hydrochloride (Bachem), the title compound was prepared as a solid (mp = >260°C). The product was purified by washing with aqueous sodium hydroxide and aqueous hydrochloric acid.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.27$  (d, J=7.1 Hz, 1H), 7.88 (d, J=7.6 Hz, 1H), 7.26 (m, 6H), 6.99 (s, 1H), 4.25 (quintet, J=7.1 Hz, 1H), 4.16 (quintet, J=7.1 Hz, 1H), 3.46 (s, 2H), 1.19 (t, J=6.3 Hz, 6H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 174.1$ , 171.8, 170.0, 136.3, 129.0, 128.1, 126.3, 48.3, 47.9, 42.0, 18.3, 18.1.

#### Example 51

### $Synthesis of \\ N'-[N-(Phenylacetyl)-L-alaninyl]-L-phenylalanina mide$

Following General Procedure T and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-phenylalaninamide (Bachem), the title compound was prepared as a solid (mp = 224-225°C).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.24$  (d, J=7.2 Hz, 1H), 7.89 (d, J=8.2 Hz, 1H), 7.36 (s, 1H), 7.13-7.34 (m, 10H), 7.11 (s, 1H), 4.40 (m, 1H), 4.21 (quintet, J=7.1 Hz, 1H), 3.44 (d, 2H), 3.01 (dd, J=4.9, 13.7 Hz, 1H), 2.82 (dd, J=9.0, 13.7 Hz, 1H), 1.13 (d, J=6.9 Hz, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.7$ , 172.0, 170.0, 137.8, 136.3, 129.2, 129.0, 128.2, 128.0, 126.3, 126.2, 53.6, 48.5, 41.9, 37.3, 18.0.

#### Example 52

25

20

### Synthesis of N'-[N-(Phenylacetyl)-L-alaninyl)-L-valinamide

Following General Procedure T and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-valinamide hydrochloride (Bachem), the title compound was prepared as a solid (mp = >261°C).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.31$  (d, J=7.5 Hz, 1H), 7.62 (d, J=9.0 Hz, 1H), 7.38 (s, 1H), 7.15-7.30 (m, 5H), 7.05 (s, 1H), 4.34 (quintet, J=7.2 Hz, 1H), 4.08 (dd, J=6.4, 15.3 Hz, 1H), 3.45 (s, 2H), 1.91 (m, 1H), 1.19 (d, J=7.0 Hz, 3H), 0.79 (d, J=6.7 Hz, 3H), 0.76 (d, J=6.8 Hz, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.8$ , 172.1, 170.0, 136.3, 129.0, 128.2, 126.3, 57.2, 48.2, 42.0, 30.5, 19.2, 17.9, 17.8.

### Example 53

10

15

5

# Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-2-amino-2-(3-pyridyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-(3-pyridyl)acetate (prepared as described in P. Kolar et al., J. Heterocyclic Chem., 28, 1715 (1991) and references cited therein), the title compound was prepared as a solid (mp = 146-157°C). The reaction was monitored by tlc (Rf = 0.1 in CHCl<sub>3</sub>/MeOH 98:2) and the product was purified by silica gel chromatography using 959:5 CHCl<sub>3</sub>/MeOH as the eluent, followed by recrystallization from chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.60$  (m, 1H), 8.56 and 8.52 (m, 1H), 7.91 (m, 1H), 7.63 (m, 1H), 7.22 (m, 1H), 6.90 (m, 1H), 6.74 (m, 2H), 5.55 (m, 1H), 4.69 (m, 1H), 4.17 (m, 2H), 3.50 and 3.41 (s, 2H), 1.33 and 1.29 (d, 3H), 1.21 (m, 3H), 1.18 (m, 3H).

25  $C_{20}H_{21}N_3O_4F_2$  (MW = 405.4); mass spectroscopy (MH<sup>+</sup>) 405.

### Example 54

## $Synthesis \ of \\ N-Methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide$

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from 30 Example B1 above) and N-methyl-L-leucinamide (prepared from N-methyl-N'-

BOC-L-leucinamide (from Example D5 above) using General Procedure Y), the title compound was prepared as a solid (mp = 233-235°C). The product was purified by recrystallization from MeOH.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>/CD<sub>3</sub>OD):  $\delta = 7.25$ -7.40 (m, 5H), 4.36 (quartet, J=7.2 Hz, 1H), 4.27 (dd, J=5.1, 14.6 Hz, 1H), 3.56 (s, 2H), 2.72 (s, 3H), 1.40-1.61 (m, 2H), 1.32 (d, J=7.1 Hz, 3H), 0.89 (d, J=6.2 Hz, 3H), 0.86 (d, J=6.2 Hz, 3H).

### Example 55

10

15

20

25

5

# Synthesis of N,N-Dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and N,N-dimethyl-L-phenylalaninamide (prepared by coupling N-BOC-L-phenylalanine (Bachem) with dimethylamine hydrochloride (Aldrich) using General Procedure B, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 152-155°C). The product was purified by extraction with EtOAc, washing with aqueous sodium carbonate and aqueous hydrochloric acid, and trituration with Et<sub>2</sub>O.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.49$  (d, J=8.2 Hz, 1H), 7.20-7.26 (m, 8H), 7.14 (m, 2H), 6.45 (d, J=7.5 Hz, 1H), 5.08 (quartet, J=8.0 Hz, 1H), 4.60 (quintet, J=7.3 Hz, 1H), 3.56 (s, 2H), 2.95 (m, 2H), 2.86 (s, 3H), 2.61 (s, 3H), 1.26 (d, J=6.9 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 171.6$ , 170.8, 170.4, 136.0, 134.7, 129.3, 129.2, 128.9, 128.8, 128.3, 127.1, 50.2, 48.7, 43.4, 39.5, 36.8, 35.6, 18.8.

### Example 56

## Synthesis of N,N-Dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and N,N-dimethyl-L-leucinamide (prepared by coupling N-BOC-L-leucine (Bachem) with dimethylamine hydrochloride (Aldrich) using General Procedure B, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 130-132°C).

The product was purified by extraction by EtOAc, washing with aqueous

sodium carbonate and aqueous hydrochloric acid, and trituration with Et2O.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 7.23-7.36 (m, 5H), 7.04 (d, J=8.7 Hz, 1H), 6.30 (d, J=7.6 Hz, 1H), 4.92 (m, 1H), 4.56 (quintet, J=7.2 Hz, 1H), 3.56 (s, 2H), 3.07 (s, 3H), 2.94 (s, 3H), 1.33-1.64 (m, 3H), 1.27 (d, J=6.9 Hz, 3H), 0.94 (d, J=6.4 Hz, 3H), 0.88 (d, J=6.5 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.0$ , 171.7, 170.4, 134.6, 129.2, 128.8, 127.2, 48.7, 47.3, 43.5, 42.1, 36.9, 35.8, 24.6, 23.3, 21.8, 18.6.

### Example 57

20

25

15

## Synthesis of N,N-Dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and N,N-dimethyl-L-valinamide (prepared by coupling N-BOC-L-valine (Bachem) with dimethylamine hydrochloride (Aldrich) using General Procedure B, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 147-149°C). The product was purified by extraction by EtOAc, washing with aqueous sodium carbonate and aqueous hydrochloric acid, and trituration with Et<sub>2</sub>O.

NMR data was as follows:

30 <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.24-7.38$  (m, 5H), 6.64 (d, 1H), 6.05 (d, 1H), 4.74 (dd, J=5.9, 8.9 Hz, 1H), 4.50 (quintet, J=7.1 Hz, 1H), 3.59 (s, 2H), 3.08 (s,

3H), 2.96 (s, 3H), 1.97 (m, 1H), 1.28 (d, J=7.0 Hz, 3H), 0.91 (d, J=6.8 Hz, 3H), 0.84 (d, J=6.8 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.3$ , 171.4, 170.4, 134.6, 129.0, 128.5, 126.8, 53.5, 48.5, 43.2, 37.3, 35.6, 31.2, 19.2, 18.6, 17.5.

 $C_{18}H_{27}N_3O_3$  (MW = 333.43); mass spectroscopy (MH<sup>+</sup>) 334.

### Example 58

## $Synthesis \ of \\ N-Methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide$

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and N-methyl-L-phenylalaninamide (prepared by coupling N-BOC-L-phenylalanine (Bachem) with methylamine hydrochloride (Aldrich) using General Procedure B, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid. The product was purified by washing with aqueous sodium carbonate and aqueous hydrochloric acid.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.23$  (d, J=7.0 Hz, 1H), 7.95 (d, J=8.2 Hz, 1H), 7.79 (d, J=4.4 Hz, 1H), 7.10-7.32 (m, 10H), 4.37 (quintet, J=5.4 Hz, 1H), 4.19 (quintet, J=7.1 Hz, 1H), 3.44 (s, 2H), 2.96 (dd, J=5.5, 13.7 Hz, 1H), 2.78 (dd, J=9.2, 13.7 Hz, 1H), 2.52 (d, J=4.4 Hz, 3H), 1.11 (d, J=7.0 Hz, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.0$ , 171.0, 170.1, 137.8, 136.3, 129.11, 129.07, 128.2, 128.1, 126.31, 126.26, 53.9, 48.5, 41.9, 37.5, 25.5, 18.0.

## Example 59

25

20

## Synthesis of N-Methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide

Following General Procedure U and using N-(phenylacetyl)-L-alanine (from Example B1 above) and N-methyl-L-valinamide (prepared by coupling N-BOC-L-valine (Bachem) with methylamine hydrochloride (Aldrich) using General

15

20

30

Procedure B, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid. The product was purified by washing with aqueous sodium carbonate and aqueous hydrochloric acid.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ): δ = 8.30 (d, J=7.6 Hz, 1H), 7.88 (d, J=4.7 Hz, 1H), 7.69 (d, J=9.1 Hz, 1H), 7.17-7.32 (m, 5H), 4.34 (quintet, J=7.2 Hz, 1H), 4.04 (dd, J=7.0, 8.9 Hz, 1H), 3.45 (s, 2H), 2.56 (d, J=4.6 Hz, 3H), 1.87 (m, 1H), 1.18 (d, J=7.0 Hz, 3H), 0.76 (d, J=6.6 Hz, 3H), 0.75 (d, J=6.7 Hz, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.0$ , 171.1, 170.0, 136.3, 129.0, 128.1, 126.3, 57.6, 48.2, 42.0, 30.6, 25.4, 19.2, 18.1, 17.9.

### Example 60

## Synthesis of N-Methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

Following General Procedure U and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-methyl-L-norleucinamide (prepared by coupling N-BOC-L-norleucine (Bachem) with methylamine hydrochloride (Aldrich) using General Procedure B, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid. The product was purified by washing with aqueous sodium carbonate and aqueous hydrochloric acid.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.37$  (d, 7.1, 1H), 7.88 (d, 8.1, 1H), 7.78 (d, 4.4, 1H), 7.08 (t, 9.5, 1H), 6.98 (d, 6.90, 2H), 4.27 (quintet, 7.0, 1H), 4.13 (quartet, 5.5, 1H), 3.51 (s, 2H), 2.54 (d, 4.4, 3H), 1.58 (m, 1H), 1.46 (m, 1H), 1.19 (m, 7H), 0.81 (t, 6.5, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.0$ , 171.9, 169.0, 162.2(dd, J=13.6, 244.0 Hz), 140.7, 112.2(dd, J=8.3, 17.0 Hz), 101.9(t, J=25.5 Hz), 52.4, 48.4, 41.3, 31.8, 27.4, 25.5, 21.8, 17.9, 13.8.

 $C_{18}H_{25}N_3O_3F_2$  (MW = 369.42); mass spectroscopy (MH<sup>+</sup>) 384.

### Example 61

## Synthesis of N,N-Dimethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

Following General Procedure U and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and N,N-dimethyl-L-norleucinamide
(prepared by coupling N-BOC-L-norleucine (Bachem) with dimethylamine
hydrochloride (Aldrich) using General Procedure B, followed by removal of the
BOC-group using General Procedure Y), the title compound was prepared as a
solid (mp = 138-140°C). The product was purified by extraction with EtOAc
and washing with aqueous sodium carbonate and aqueous hydrochloric acid.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.11$  (d, 8.1, 1H), 6.81 (m, 2H), 6.71 (m, 1H), 6.60 (d, 7.6, 1H), 4.89 (q, J=5.0, 1H), 4.57 (quint, J=7.1, 1H), 3.53 (s, 2H), 3.08 (s, 3H), 2.97 (s, 3H), 1.70 (m, 1H), 1.55 (m, 1H), 1.20-1.38 (m, 7H), 0.85 (t, 6.9, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 171.6$ , 171.5, 168.9, 163.0 (dd, J=12.9, 247.3 Hz), 138.4, 112.2 (dd, J=7.8, 17.0 Hz), 102.7 (t, J=25.0 Hz), 49.1, 48.9, 42.9, 37.1, 35.8, 32.6, 27.1, 22.4, 19.1, 13.8.

 $C_{19}H_{27}N_3O_3F_2$  (MW = 383.44); mass spectroscopy (MH<sup>+</sup>) 384.

### Example 62

# Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

- Following General Procedure U and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-norleucinamide (prepared from N-BOC-L-norleucinamide (from Example D6 above) using General Procedure Y), the title compound was prepared as a solid (mp = >215°C). The product was purified by precipitation from water.
- 30 NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.37$  (d, 7.4, 1H), 7.83 (d, 8.0, 1H), 7.29 (s, 1H), 6.95-7.14 (m, 4H), 4.29 (quintet, J=7.2, 1H), 4.14 (quartet, J=5.0, 1H), 3.52 (s, 2H), 1.61 (m, 1H), 1.46 (m, 1H), 1.21 (m, 7H), 0.82 (m, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 173.6$ , 171.9, 168.9, 162.0 (dd), 140.7, 112.2 (dd, J=7.5, 16.6 Hz), 101.9 (t), 52.2, 48.3, 41.3, 31.8, 27.4, 21.8, 18.0, 13.8.

 $C_{17}H_{23}N_3O_3F_2$  (MW = 355.39); mass spectroscopy (MH<sup>+</sup>) 356.

### Example 63

10

15

20

5

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-methoxyphenyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(3-methoxyphenyl)acetate hydrochloride (prepared by the Bucherer Modification of the Strecker procedure as described in J. P. Greenstein et al., "The Chemistry of Amino Acids", Vol. 1, p. 698, Wiley, New York (1961)), the title compound was prepared as a solid (mp =  $163-170^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.45 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 97:3 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.27$  (m, 1H), 7.18 and 7.06 (m, 1H), 6.87-6.67 (m, 6H), 6.25 (m, 1H), 5.46 (m, 1H), 4.58 (m, 1H), 3.82 (s, 3H), 3.71 and 3.69 (s, 3H), 3.53 and 3.48 (s, 3H), 1.39 and 1.30 (d, 3H).

 $C_{21}H_{22}N_2O_5F_2$  (MW = 420.42); mass spectroscopy (MH<sup>+</sup>) 421.

25

## Example 64

Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-methoxyphenyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(4-methoxyphenyl)acetate hydrochloride (prepared by the Bucherer Modification of the Strecker procedure as described in J. P. Greenstein et al., "The Chemistry of Amino Acids", Vol. 1, p. 698, Wiley, New York (1961)), the title compound was prepared as a solid (mp = 170-174°C). The reaction was monitored by tlc (Rf = 0.1 in 98:2 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 98:2 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

 $C_{21}H_{22}N_2O_5F_2$  (MW = 420.42); mass spectroscopy (MH<sup>+</sup>) 421.

### Example 65

15

20

25

30

5

## Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-(2-pyridyl)acetate hydrochloride (prepared as described in P. Kolar et al., J. Heterocyclic Chem., 28, 1715 (1991) and references cited therein), the title compound was prepared as a solid (mp = 123-125°C). The reaction was monitored by tlc (Rf = 0.1 in 98:2 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 95:5 CHCl<sub>3</sub>/MeOH as the eluent, followed by recrystallization from chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.53$  (m, 1H), 7.70 (m, 2H), 7.48 (m, 1H), 7.27 (m, 1H), 6.86 (m, 2H), 6.74 (m, 1H), 6.52 (m, 1H), 5.58 (m, 1H), 4.67 (m, 1H), 4.18 (m, 2H), 3.54 and 3.50 (s, 2H), 1.48 and 1.39 (d, 3H), 1.21 (m, 3H).

 $C_{20}H_{21}N_3O_4F_2$  (MW = 405.4); mass spectroscopy (MH<sup>+</sup>) 405.

10

20

25

### Example 66

# Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-(4-pyridyl)acetate hydrochloride (prepared as described in P. Kolar et al., J. Heterocyclic Chem., 28, 1715 (1991) and references cited therein), the title compound was prepared as a solid (mp = 175-181°C). The reaction was monitored by tlc (Rf = 0.1 in 98:2 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 95:5 CHCl<sub>3</sub>/MeOH as the eluent, followed by recrystallization from chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.59$  (m, 2H), 7.39 (m, 1H), 7.26 (m, 2H), 6.80 (m, 3H), 6.21 (m, 1H), 5.51 (m, 1H), 4.62 (m, 1H), 4.21 (m, 2H), 3.57 and 3.51 (s, 2H), 1.38 (m, 3H), 1.23 (m, 3H).

 $C_{20}H_{21}N_3O_4F_2$  (MW = 405.4); mass spectroscopy (MH<sup>+</sup>) 405.

### Example 67

## Synthesis of N-[N-(Cyclohexylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure U and using cyclohexylacetic acid (Aldrich) and N-(L-alaninyl)-L-phenylalanine methyl ester (prepared by coupling N-BOC-L-alanine (Bachem) with L-phenylalanine methyl ester hydrochloride (Bachem) using General Procedure U, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 156-158°C). The reaction was monitored by tlc (Rf = 0.25 in 1:1 EtOAc/hexanes).

NMR data was as follows:

30 <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 0.95 (m, 2H), 1.10-1.38 (m, 3H), 1.33 (d, J=7.0 Hz, 3H), 1.60-1.86 (m, 6H), 2.02 (d, J=7.5 Hz, 2H), 3.10 (m, 2H), 3.71 (s,

3H), 4.49 (m, 1H), 4.81 (m, 1H), 6.10 (d, J=7.3 Hz, 1H), 6.65 (d, J=7.7 Hz, 1H), 7.11 (m, 2H), 7.26 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.4$ , 26.0, 26.1, 33.0, 33.1, 35.3, 37.8, 44.5, 48.5, 52.4, 53.3, 127.1, 128.6, 129.2, 135.6, 171.6, 172.0, 172.2.

 $C_{21}H_{30}N_2O_4$  (MW = 374.48); mass spectroscopy (MH<sup>+</sup>) 375.

### Example 68

## Synthesis of N-[N-(Cyclopentylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure U and using cyclopentylacetic acid (Aldrich) and N-(L-alaninyl)-L-phenylalanine methyl ester (prepared by coupling N-BOC-L-alanine (Bachem) with L-phenylalanine methyl ester hydrochloride (Bachem) using General Procedure U, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 137-139°C). The reaction was monitored by tlc (Rf = 0.23 in 1:1 EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 1.13$  (m, 2H), 1.33 (d, J=7.0 Hz, 3H), 1.58 (m, 4H), 1.80 (m, 2H), 2.17 (m, 3H), 3.10 (m, 2H), 3.71 (s, 3H), 4.50 (m, 1H), 4.83 (m, 1H), 6.12 (d, J=7.4 Hz, 1H), 6.69 (d, J=7.7 Hz, 1H), 7.2 (m, 2H), 7.25 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 18.3, 24.9, 32.4, 32.5, 37.0, 37.7, 42.7, 48.4, 52.3, 53.3, 127.1, 128.5, 129.2, 135.7, 171.6, 172.0, 172.6. C<sub>20</sub>H<sub>28</sub>N<sub>2</sub>O<sub>4</sub> (MW = 360.46); mass spectroscopy (MH<sup>+</sup>) 361.

25

30

20

### Example 69

## Synthesis of N-[N-(Cyclohex-1-enylacetyl)-L-alaninyl]L-phenylalanine Methyl Ester

Following General Procedure U and using cyclohex-1-enylacetic acid (Alfa) and N-(L-alaninyl)-L-phenylalanine methyl ester (prepared by coupling N-BOC-

L-alanine (Bachem) with L-phenylalanine methyl ester hydrochloride (Bachem) using General Procedure U, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 139-142°C). The reaction was monitored by tlc (Rf = 0.27 in 1:1

5 EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 1.31$  (d, J=7.0 Hz, 3H), 1.58 (m, 4H), 1.89 (m, 2H), 2.04 (br s, 2H), 2.83 (s, 2H), 3.00-3.20 (m, 2H), 3.71 (s, 3H), 4.47 (m, 1H), 4.81 (m, 1H), 5.60 (s, 1H), 6.26 (d, J=7.3 Hz, 1H), 6.67 (d, J=7.7 Hz, 1H), 7.11 (m, 2H), 7.26 (m, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.1, 21.9, 22.7, 25.3, 28.3, 37.7, 46.0, 48.4,$ 52.3, 53.3, 127.1, 127.2, 128.5, 129.1, 132.2, 135.7, 171.0, 171.6, 171.8.  $C_{21}H_{28}N_2O_4$  (MW = 372.47); mass spectroscopy (MH<sup>+</sup>) 373.

### Example 70

15

20

10

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-1-aminocyclopropane-1-carboxylate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 1-aminocyclopropane-1-carboxylate hydrochloride (Sigma), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.3 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 97:3 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

25 'H-nmr (CDCl<sub>3</sub>):  $\delta = 6.96$  (bs, 1H), 6.82 (m, 2H), 6.69 (m, 1H), 6.48 (d, 1H), 4.50 (m, 1H), 3.67 (s, 3H), 3.54 (s, 2H), 1.58 (m, 2H), 1.40 (d, 2H), 1.12 (m, 2H).

Optical Rotation:  $[\alpha]_{23} = -18^{\circ}$  (c 1, MeOH).

WO 98/22494 PCT/US97/20804

-- 226 --

### Example 71

## Synthesis of N-2-(N, N-Dimethylamino) ethyl-N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and N,N,N'-trimethylethylenediamine (Aldrich), the title compound was prepared as a solid.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.37$  (m, 2H), 8.19 (d, 1H), 8.08 (d, 2H), 7.10 (m, 1H), 6.99 (m, 2H), 4.67 (m, 1H), 4.30 (m, 1H), 3.52 (s, 2H), 3.01 and 2.86 (s, 3H), 2.47 (t, 1H), 2.31 (t, 1H), 2.15 (s, 6H), 1.19 (m, 6H).

Optical Rotation:  $[\alpha]_{23} = -85^{\circ}$  (c 1, MeOH).

 $C_{19}H_{28}N_4O_3F_2$  (MW = 398.45); mass spectroscopy (MH<sup>+</sup>) 398.

### Example 72

15

20

25

5

10

## Synthesis of N-[N-(Cyclopropylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure U and using cyclopropylacetic acid (Lancaster) and N-(L-alaninyl)-L-phenylalanine methyl ester (prepared by coupling N-BOC-L-alanine (Bachem) with L-phenylalanine methyl ester hydrochloride (Bachem) using General Procedure U, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp =  $128-131^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.14in 1:1 EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.17$  (m, 2H), 0.59 (m, 2H), 0.92 (m, 1H), 1.35 (d, J=7.0 Hz, 3H), 2.11 (m, 2H), 3.05 (dd, J=6.7, 13.9 Hz, 1H), 3.16 (dd, J=5.5, 13.9 Hz, 1H), 3.73 (s, 3H), 4.52 (m, 1H), 4.82 (m, 1H), 6.47 (d, J=7.1 Hz, 1H), 6.70 (d, J=7.5 Hz, 1H), 7.12 (m, 2H), 7.28 (m, 3H).

30 <sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 4.6, 6.9, 18.2, 37.7, 41.2, 48.4, 52.4, 53.2, 127.1,$ 128.5, 129.2, 135.7, 171.7, 171.9, 172.3.

 $C_{18}H_{24}N_2O_4$  (MW = 332.40); mass spectroscopy (MH<sup>+</sup>) 333.

### Example 73

## Synthesis of N-[N-(3,5-Difluorophenylacety])-L-alaninyl]glycine Benzyl Ester

5

10

15

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and glycine benzyl ester (prepared from N-BOC-glycine (Bachem) and benzyl alcohol (Aldrich) using General Procedure X, followed by removal of the BOC-group using General Procedure Y), the title compound was prepared as a solid (mp = 167.5°C). The reaction was monitored by tlc (Rf = 0.35 in 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by flash chromotography using 2% MeOH/CH2Cl2 as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.12$  (m, 5H), 6.71 (m, 3H), 6.60 (m, 2H), 4.95 (s, 2H), 4.18 (q, 1H), 3.76 (dd, 2H), 3.35 (s, 2H), 1.13 (d, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 176.0, 172.9, 171.5, 166.46, 163.30, 141.54,$ 137.70, 130.11, 129.88, 113.98, 113.87, 113.75, 113.64, 103.89, 103.55, 103.21, 68.44, 50.93, 43.25, 42.61, 18.65.

 $C_{20}H_{20}N_2O_4F_2$  (MW = 390.39); mass spectroscopy (MH<sup>+</sup>) 391.

20

25

#### Example 74

## Synthesis of N-[N-(Isovaleryl)-L-phenylglycinyl]-L-alanine Ethyl Ester

Following General Procedure C and using N-(isovaleryl)-L-phenylglycine (prepared from isovaleric acid (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure C, followed by hydrolysis using General Procedure AF) and L-alanine ethyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 198-201°C). The reaction was monitored by tlc (Rf = 0.3 in 1:1 EtOAc/hexanes) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by rerystallization from EtOAc.

30

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:5 mixture of diastereomers):  $\delta = 1.25$  and 1.30 (two d, 3H), 5.57 (d, 1H), 5.60 (d, 1H).

 $C_{18}H_{26}N_2O_4$  (MW = 334.42); mass spectroscopy (MH<sup>+</sup>) 335.

5

#### Example 75

## Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure Z and using N-(3-nitrophenylacetyl)-L-alanine 2,4,5-trichlorophenyl ester (from Example D8 above) and L-phenylalanine methyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 154-158°C). The reaction was monitored by tlc (Rf = 0.3 in 1:1 EtOAc/hexanes) and the product was purified by silica gel chromatography using 50-100% EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:3 mixture of diastereomers):  $\delta = 1.00$  and 1.18 (two d, 3H), 2.96 (m, 2H).

 $C_{21}H_{23}N_3O_6$  (MW = 413.43); mass spectroscopy (MH<sup>+</sup>) 413.

Example 76

20

25

# Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]L-alanine Ethyl Ester

Following General Procedure Z and using N-(3-nitrophenylacetyl)-L-alanine 2,4,5-trichlorophenyl ester (from Example D8 above) and L-alanine ethyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 193-195°C). The reaction was monitored by tlc (Rf = 0.4 in EtOAc) and the product was purified by silica gel chromatography using EtOAc as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (m, 9H), 3.65 (s, 2H); 4.05 (m, 2H).

30 Optical Rotation:  $[\alpha]_{20} = -27.3^{\circ}$  @ 589nm, (c = 1.02, DMSO).

 $C_{16}H_{21}N_3O_6$  (MW = 351.36); mass spectroscopy (MH<sup>+</sup>) 352.

### Example 77

## Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]glycine Ethyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (prepared from 3-nitrophenylacetic acid (Aldrich) and L-alanine ethyl ester hydrochloride (Sigma) using General Procedure C, followed by hydrolysis using General Procedure AF) and glycine ethyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 164-165°C). The product was purified by silica gel chromatography using EtOAc as the eluent, followed by recrystallization from EtOAc.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (m, 6H), 4.08 (q, 2H); 4.32 (m, 1H).

Optical Rotation:  $[\alpha]_{20} = -25^{\circ} @ 589 \text{ nm}$ , (c = 1.00, DMSO).

 $C_{15}H_{19}N_3O_6$  (MW = 337.33); mass spectroscopy (MH<sup>+</sup>) 338.

### Example 78

# Synthesis of N-Hydroxy-N'-[N-(3-nitrophenylacetyl)-L-alaninyl]-D,L-threoninamide

- Following General Procedure Z and using N-(3-nitrophenylacetyl)-L-alanine 2,4,5-trichlorophenyl ester (from Example D8 above) and D,L-threonine hydroxamate (Sigma), the title compound was prepared as a solid (mp = 180-183°C). The reaction was monitored by tlc (Rf = 0.25 in 15% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 15%
- 25 MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from EtOAc.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 1.22$  (m, 3H); 0.98 (m, 3H).

 $C_{15}H_{20}N_4O_7$  (MW = 368.35); mass spectroscopy (MH<sup>+</sup>) 368.

10

20

25

30

-- 230 --

## Example 79

## Synthesis of N-[N-(Isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl Ester

Following General Procedure C and using N-(isovaleryl)-L-phenylglycine (prepared from isovaleric acid (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure C, followed by hydrolysis using General Procedure AF) and L-alanine iso-butyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 2-methyl-1-propanol (Aldrich) using General Procedure C (with catalystic DMAP), followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 181-186°C). The reaction was monitored by tlc (Rf = 0.4 in 1:1 EtOAc/hexanes) and the product was purified by silica gel chromatography using 1:1 EtOAc/hexanes as the eluent.

NMR data was as follows:

### Example 80

Synthesis of Methyl N-[N-(3-nitrophenylacetyl)-L-alaninyl]-2-amino-3-(3-hydroxyphenyl)propionate

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (prepared from 3-nitrophenylacetic acid (Aldrich) and L-alanine ethyl ester hydrochloride (Sigma) using General Procedure C, followed by hydrolysis using General Procedure AF) and methyl 2-amino-3-(3-hydroxyphenyl)propionate (prepared from 2-amino-3-(3-hydroxyphenyl)propionate (Biosynth AG, Switzerland) and methanol using General Procedure H), the title compound was prepared as a solid (mp = 155-159°C). The reaction was monitored by tlc (Rf = 0.4 in EtOAc) and the product was purified by silica gel chromatography using EtOAc as the eluent.

NMR data was as follows:

-- 231 --

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 1.02$  and 1.20 (two d, 3H); 3.62 (2 s, 3H).

 $C_{21}H_{23}N_2O_7$  (MW = 429.43); mass spectroscopy (MH<sup>+</sup>) 429.

### Example 81

5

10

## Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-tyrosine Ethyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (prepared from 3-nitrophenylacetic acid (Aldrich) and L-alanine ethyl ester hydrochloride (Sigma) using General Procedure C, followed by hydrolysis using General Procedure AF) and L-tyrosine ethyl ester (Sigma), the title compound was prepared as a solid (mp = 117-119°C). The reaction was monitored by tlc (Rf = 0.5 in EtOAc) and the product was purified by silica gel chromatography using EtOAc as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.07$  (t, 3H); 1.20 (d, 3H); 9.23 (s, 1H). 15 Optical Rotation:  $[\alpha]_{20} = -13.1^{\circ} @ 589 \text{ nm}, (c = 1.08, DMSO).$  $C_{22}H_{25}N_3O_7$  (MW = 443.46); mass spectroscopy (MH<sup>+</sup>) 443/444.

### Example 82

## Synthesis of N-[N-(Isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl Ester

20

25

Following General Procedure C and using N-(isovaleryl)-L-isoleucine (prepared from isovaleric acid (Aldrich) and L-isoleucine methyl ester hydrochloride (Aldrich) using General Procedure C, followed by hydrolysis using General Procedure AF) and L-alanine iso-butyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 2-methyl-1-propanol (Aldrich) using General Procedure C (with catalystic DMAP), followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 142-146°C). The reaction was monitored by tlc (Rf = 0.4 in 1:1

20

EtOAc/hexanes) and the product was purified by silica gel chromatography using 1:1 EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:4 mixture of diastereomers):  $\delta = 1.26$  (d, 3H), 7.70, 7.80 (doublets, 1H); 8.30, 8.40 (doublets, 1H).

 $C_{18}H_{34}N_2O_4$  (MW = 342.48); mass spectroscopy (MH<sup>+</sup>) 343.

## Example 83

Step A --Synthesis of N-[N-[N-(tert-Butoxycarbonyl)-L-valinyl]-D,L-phenylglycinyl]-L-alanine iso-butyl Ester

Following General Procedure A and using N-[N-BOC-L-valinyl]-D,L-phenylglycine (prepared by coupling N-BOC-L-valine (Bachem) and L-phenylglycine methyl ester hydrochloride (Sigma) using General Procedure C, followed by hydrolysis of the methyl ester using General Procedure AF) and L-alanine iso-butyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 2-methyl-1-propanol (Aldrich) using General Procedure C (with catalytic DMAP), followed by removal of the BOC-group using General Procedure P), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.3 in 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 1.25$  (d, 3H); 5.58 (d, 1H).

 $C_{25}H_{39}N_3O_6$  (MW = 477.61); mass spectroscopy (MH<sup>+</sup>) 478.

Step B --Synthesis of N-[N-(L-Valinyl)-L-phenylglycinyl]-L-alanine isobutyl Ester Hydrochloride

Following General Procedure P and using the product from Example 83 - Step A above, the title compound was prepared as a solid (mp = 225-232°C). The product was purified by trituration in Et<sub>2</sub>O.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:2 mixture of diastereomers):  $\delta = 1.26$ , 1.32 (doublets, 3H); 5.60, 5.65 (doublets, 1H).

 $C_{20}H_{32}N_3O_4Cl$  (MW = 413.94); mass spectroscopy (MH<sup>+</sup>) 378 (free base).

Step C --Synthesis of N-[N-[N-(Isovaleryl)-L-valinyl]-L-phenylglycinyl]-L-alanine iso-butyl Ester

Following General Procedure C and using isovaleric acid (Aldrich) and the product from Example 83 - Step B above, the title compound was prepared as a solid (mp = 217-221°C). The reaction was monitored by tlc (Rf = 0.25 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub>) as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:3 mixture of diastereomers):  $\delta = 5.52$ , 5.58 (doublets, 1H).

 $C_{25}H_{39}N_3O_5$  (MW = 461.60); mass spectroscopy (MH<sup>+</sup>) 462.

15

20

25

5

10

## Example 84

## Synthesis of N-[N-(Isovaleryl)-L-phenylalaninyl]-L-alanine iso-butyl Ester

Following General Procedure C and using isovaleric acid (Aldrich) and N-(L-phenylalaninyl)-L-alanine iso-butyl ester hydrochloride (prepared from N-BOC-L-phenylalanine (Sigma) and L-alanine iso-butyl ester hydrochloride (prepared as described in Example 83A above) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 135-138°C). The reaction was monitored by tlc (Rf = 0.3 in 3% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 0.75$  (d, 3H), 0.84 (d, 3H); 0.90 (d, 6H); 1.33 (d, 3H).

Ontical Potation: [at]  $= \pm 4.71^{\circ} \otimes 590 \text{ nm}$  (a = 1.02 DMSO)

 $C_{21}H_{32}N_2O_4$  (MW = 376.50); mass spectroscopy (MH<sup>+</sup>) 376.

## Example 85

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-alanine Ethyl Ester

Following General Procedure C and using 3,5-difluorophenylacetic acid (Oakwood) and L-alanine ethyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 197-199°C). The reaction was monitored by tlc (Rf = 0.6 in EtOAc) and the product was purified from bi-products by silica gel chromatography using EtOAc as the eluent, followed by recrystallization from EtOAc. 10

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.22$  (m, 9H); 3.52 (s, 2H).

Optical Rotation:  $[\alpha]_{20} = -76.1^{\circ}$  @ 589 nm, (c = 1.01, DMSO).

 $C_{16}H_{20}N_2O_4F_2$  (MW = 342.34); mass spectroscopy (MH<sup>+</sup>) 343.

15

20

5

## Example 86

## Synthesis of Ethyl 1-[N-(3-Nitrophenylacetyl)-L-alaninyllindoline-(S)-2-carboxylate

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (prepared from 3-nitrophenylacetic acid (Aldrich) and L-alanine ethyl ester hydrochloride (Sigma) using General Procedure C, followed by hydrolysis using General Procedure AF) and ethyl (S)-indoline-2-carboxylate (prepared from (S)indoline-2-carboxylic acid (Aldrich) and ethanol using General Procedure H), the title compound was prepared as a solid. The reaction was monitored by tlc

## Example 87

## Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-alaninamide hydrochloride (Sigma), the title compound was prepared as a solid (mp = 285-288°C). The reaction was monitored by tlc (Rf = 0.35 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from EtOH.

10 NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.21$  (m, 6H); 7.95 (d, 1H); 8.37 (d, 1H). Optical Rotation:  $[\alpha]_{20} = -26.84^{\circ}$  @ 589 nm, (c = 1.01, DMSO).  $C_{14}H_{17}N_3O_3F_2$  (MW = 313.31); mass spectroscopy (MH<sup>+</sup>) 314.

## Example 88

15

20

5

# Synthesis of N-Methoxy-N-methyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide

Following General Procedure C and using N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine (prepared from N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine ethyl ester (from Example 74 above) using General Procedure AF) and N, O-dimethylhydroxylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.6 in EtOAc) and the product was purified by silica gel chromatography using EtOAc as the eluent.

NMR data was as follows:

O TI NO (2011 040 40)

. ....

## Example 89

# Synthesis of N-iso-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester (from Example 85 above) using General Procedure AF) and iso-butylamine (Aldrich), the title compound was prepared as a solid (mp = 258-260°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 0.80$  (d, 6H); 1.20 (m, 6H).

Optical Rotation:  $[\alpha]_{20} = -30.4^{\circ} @ 589 \text{ nm}, (c = 1.01, DMSO).$ 

 $C_{18}H_{25}N_3O_3F_2$  (MW = 369.41); mass spectroscopy (MH<sup>+</sup>) 369.

### Example 90

## Synthesis of N,N-Di-n-propyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester (from Example 85 above) and di-n-propylamine (Aldrich), the title compound was prepared as a solid (mp = 137-146°C). The reaction was monitored by tlc (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:2 mixture of diastereomers):  $\delta = 3.50$  (s, 2H), 4.30 (m, 1H), 4.63 (m, 1H).

30  $C_{20}H_{29}N_3O_3F_2$  (MW = 397.46); mass spectroscopy (MH<sup>+</sup>) 397.

15

20

## Example 91

## Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-valinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-valinamide hydrochloride (Sigma), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

 $C_{16}H_{21}N_3O_3F_2$  (MW = 341.36) mass spectroscopy (MH<sup>+</sup>) 342.

## Example 92

# Synthesis of N-(4-Nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-diffuorophenylacetyl)-L-alanine (from Example B2 above) and N-(4-nitrophenyl)-L-alaninamide hydrochloride (Fluka), the title compound was prepared as a solid (mp = 242-244°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.24$  (d, 3H); 1.33 (d, 3H).

25 Optical Rotation:  $[\alpha]_{20} = -5.18^{\circ}$  @ 589 nm, (c = 1.00, DMSO).  $C_{20}H_{20}N_4O_5F_2$  (MW = 434.40); mass spectroscopy (MH<sup>+</sup>) 434.

## Example 93

Synthesis of N'-[N-[N-(Isovaleryl)-L-phenylglycinyl]-

Following General Procedure C and using N-(isovaleryl)-L-phenylglycine (prepared from isovaleric acid (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure C, followed by hydrolysis using General Procedure AF) and N'-(L-alaninyl)-L-phenylalaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and Lphenylalaninamide (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 272-276°C). The reaction was monitored by tlc (Rf = 0.25 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 1.07, 1.17$ (doublets, 3H); 5.40, 5.52 (doublets, 1H).

 $C_{25}H_{32}N_4O_4$  (MW = 452.55); mass spectroscopy (MH<sup>+</sup>) 453.

15

25

5

10

### Example 94

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure C and using 3,5-difluorophenylacetic acid 20 (Oakwood) and N-(L-alaninyl)-L-phenylalanine methyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylalanine methyl ester hydrochloride (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 173-175°C). The reaction was monitored by tlc (Rf = 0.6 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 4% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.17$  (d, 3H); 3.48 (s, 2H).

30 Optical Rotation:  $[\alpha]_{20} = -32.47^{\circ}$  @ 589 nm, (c = 1.01, MeOH).  $C_{21}H_{22}N_2O_4F_2$  (MW = 404.41); mass spectroscopy (MH<sup>+</sup>) 404.

## Example 95

## Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure C and using 3,5-difluorophenylacetic acid

(Oakwood) and N'-(L-alaninyl)-L-phenylalaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylalaninamide (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 252-253°C). The reaction was monitored by tlc (Rf = 0.5 in 15% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 15% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from EtOH.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.15$  (d, 3H); 3.51 (s, 2H).

Optical Rotation:  $[\alpha]_{20} = -24.4^{\circ}$  @ 589 nm, (c = 1.01, DMSO).

 $C_{20}H_{21}N_3O_3F_2$  (MW = 389.41); mass spectroscopy (MH<sup>+</sup>) 389.

### Example 96

## $Synthesis of \\ N-iso-butyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alanina mide$

Following General Procedure C and using N-[N-(isovaleryl)-Lphenylglycinyl]-L-alanine (prepared from N-[N-(isovaleryl)-L-phenylglycinyl]-Lalanine ethyl ester (from Example 74 above) using General Procedure AF) and
iso-butylamine (Aldrich), the title compound was prepared as a solid (mp =
227-232°C). The reaction was monitored by tlc (Rf = 0.3 in 5%
MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using
5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:4 mixture of diastereomers):  $\delta = 1.58$  (m, 1H); 1.95 (m, 1H); 5.55 (d, 1H).

 $C_{20}H_{31}N_3O_3$  (MW = 361.48); mass spectroscopy (MH<sup>+</sup>) 361.

25

### Example 97

# Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94) using General Procedure AF) and 2-methoxyethylamine (Aldrich), the title compound was prepared as a solid (mp = 206-208°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.14$  (d, 3H); 4.22 (m, 1H); 4.45 (m, 1H). Optical Rotation:  $[\alpha]_{20} = -25^{\circ}$  @ 589 nm, (c = 1.00, DMSO).  $C_{23}H_{27}N_3O_4F_2$  (MW = 447.49); mass spectroscopy (MH<sup>+</sup>) 447.

## Example 98

# Synthesis of N-(4-Nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester (from Example 85 above) using General Procedure AF) and 4-nitrobenzylamine (Aldrich), the title compound was prepared as a solid (mp = 257-259°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from EtOH/acetonitrile.

NMR data was as follows:

 . 5

10

15

20

25

 $C_{21}H_{22}N_4O_5F_2$  (MW = 448.43); mass spectroscopy (MH<sup>+</sup>) 448.

### Example 99

# Synthesis of N-(4-Nitrophenyl)-N'-[N-(N-(isovaleryl)-L-phenylglycinyl]-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(isovaleryl)-L-phenylglycinyl]-L-phenylglycinyl]-L-alanine (prepared from N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine ethyl ester (from Example 74 above) using General Procedure AF) and N-(4-nitrophenyl)-L-alaninamide hydrochloride (Fluka), the title compound was prepared as a solid (mp = 255-257°C). The reaction was monitored by tle (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:2 mixture of diastereomers):  $\delta = 5.45$ , 5.55 (doublets, 1H); 10.20, 10.54 (singlets, 1H).

 $C_{25}H_{31}N_4O_6$  (MW = 497.56); mass spectroscopy (MH<sup>+</sup>) 497.

## Example 100

## Synthesis of N-(4-Nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(4-nitrophenyl)-L-phenylalaninamide hydrochloride (Lancaster), the title compound was prepared as a solid (mp = 253-254°C). The reaction was monitored by tlc (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 8% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.17$  (d, 3H); 10.52 (s, 1H).

30 Optical Rotation:  $[\alpha]_{20} = +40.6^{\circ}$  @ 589 nm, (c = 1.00, DMSO).

10

15

 $C_{26}H_{24}N_4O_5F_2$  (MW = 510.50); mass spectroscopy (MH<sup>+</sup>) 510.

## Example 101

## Synthesis of N-Benzyl-N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester (from Example 85 above) using General Procedure AF) and N-benzyl-N-methylamine (Aldrich), the title compound was prepared as a solid (mp = 167-169°C). The reaction was monitored by tlc (Rf = 0.4 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:3 mixture of diastereomers):  $\delta = 3.52$  (singlets, 2H); 2.95 (s, 2H).

Optical Rotation:  $[\alpha]_{20} = -55.8^{\circ}$  @ 589 nm, (c = 1.01, DMSO).  $C_{22}H_{25}N_3O_3F_2$  (MW = 417.45); mass spectroscopy (MH<sup>+</sup>) 417.

## Example 102

20

## Synthesis of N-(3,5-Difluorobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(3,5-difluorobenzyl)-L-alaninamide

bydrochloride (prepared from N-BOC-L-alanine (Sigma) and 3,5-difluorobenzylamine (Lancaster) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 267-269°C). The reaction was monitored by tlc (Rf = 0.25 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel

WO 98/22494

-- 243 --

chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.21$  (d, 3H), 1.24 (d, 3H).

Optical Rotation:  $[\alpha]_{20} = +26.9^{\circ} @ 589 \text{ nm}, (c = 1.01, DMSO).$ 

 $C_{21}H_{21}N_3O_3F_4$  (MW = 439.41); mass spectroscopy (MH<sup>+</sup>) 439.

## Example 103

## Synthesis of N-(3-Nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

10

15

5

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and N-(3-nitrobenzyl)-L-alaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 3-nitrobenzylamine hydrochloride (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 245-247°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

20 NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.21$  (d, 3H); 1.25 (d, 3H).

Optical Rotation:  $[\alpha]_{20} = -32.8^{\circ}$  @ 589 nm, (c = 1.00, DMSO).

 $C_{21}H_{22}N_4O_5F_2$  (MW = 448.43); mass spectroscopy (MH<sup>+</sup>) 449.

## Example 104

25

30

## Synthesis of N-Benzyl-N'-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and N-benzyl-L-alaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and benzylamine (Aldrich) using

General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 260-262°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (d, 3H); 1.24 (d, 3H). Optical Rotation:  $[\alpha]_{20} = -29.3^{\circ}$  @ 589 nm, (c = 1.03, DMSO).  $C_{21}H_{23}N_3O_3F_2$  (MW = 403.43); mass spectroscopy (MH<sup>+</sup>) 403.

10

25

5

#### Example 105

## Synthesis of N-(4-Nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94) using General Procedure AF) and 4-nitrobenzylamine hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 248-250°C). The reaction was monitored by tlc (Rf = 0.4 in 12% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 12% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.15$  (d, 3H); 7.35 (d, 2H); 8.12 (d, 2H). Optical Rotation:  $[\alpha]_{20} = -27.6^{\circ}$  @ 589 nm (c = 1.01, DMSO).  $C_{27}H_{26}N_4O_5F_2$  (MW = 524.52); mass spectroscopy (MH<sup>+</sup>) 524.

Example 106

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-tryptophan Methyl Ester

10

30

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-tryptophan methyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 191-193°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (d, 3H); 3.55 (s, 3H). Optical Rotation:  $[\alpha]_{20} = -8.82^{\circ}$  @ 589 nm (c = 1.02, DMSO).  $C_{23}H_{23}N_3O_4F_2$  (MW = 443.45); mass spectroscopy (MH<sup>+</sup>) 443.

### Example 107

# Synthesis of N-(4-Methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(4-methoxybenzyl)-L-alaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 4-methoxybenzylamine hydrochloride (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 234-236°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from EtOH/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (d, 6H); 3.51 (s, 2H); 3.72 (s, 3H). Optical Rotation:  $[\alpha]_{20} = +27.9^{\circ}$  @ 589 nm (c = 1.00, DMSO).  $C_{22}H_{25}N_3O_4F_2$  (MW = 433.46); mass spectroscopy (MH<sup>+</sup>) 433.

### Example 108

Synthesis of N-[N-(Phenylacetyl)-L-phenylglycinyl]-L-alanine Ethyl Ester

Following General Procedure C and using phenylacetic acid (Aldrich) and N-(L-phenylglycinyl)-L-alanine ethyl ester hydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and L-alanine ethyl ester hydrochloride (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 208-210°C). The reaction was monitored by tlc (Rf = 0.4 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

10 NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 3.55$  (s, 2H); 5.55 (d, 1H). Optical Rotation:  $[\alpha]_{20} = +44.8^{\circ}$  @ 589 nm (c = 1.02, DMSO).

 $C_{21}H_{24}N_2O_4$  (MW = 368.43); mass spectroscopy (MH<sup>+</sup>) 369.

## Example 109

15

20

25

5

# Synthesis of N-[N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalaninyl]-L-phenylglycine Methyl Ester

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine (prepared from N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94) using General Procedure AF) and L-phenylglycine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 203-207°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by trituration using 1-chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.13$  (d, 3H); 3.62 (s, 3H). Optical Rotation:  $[\alpha]_{20} = +42.1^{\circ}$  @ 589 nm (c = 1.03, DMSO).  $C_{29}H_{29}N_3O_5F_2$  (MW = 537.56); mass spectroscopy (MH<sup>+</sup>) 537.

## Example 110

# Synthesis of N-[N-(Cyclohexylacetyl)-L-phenylglycinyl]L-alanine Ethyl Ester

Following General Procedure C and using cyclohexylacetic acid (Aldrich) and N-(L-phenylglycinyl)-L-alanine ethyl ester hydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and L-alanine ethyl ester hydrochloride (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 196-198°C). The reaction was monitored by tlc (Rf = 0.3 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by trituration using 1-chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 2.08$  (d, 2H); 5.56 (d, 1H).

Optical Rotation:  $[\alpha]_{20} = +26.3^{\circ}$  @ 589 nm (c = 1.01, DMSO).

 $C_{21}H_{30}N_2O_4$  (MW = 374.48); mass spectroscopy (MH<sup>+</sup>) 375.

### Example 111

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine Methyl Ester

20

25

15

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylglycine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 198-200°C). The reaction was monitored by tlc (Rf = 0.4 in 4% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using

MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 4% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.26$  (d, 3H); 3.64 (s, 3H).

30 Optical Rotation: (DMSO)  $[\alpha]_{20} = +69.9^{\circ}$  @ 589 nm (c = 1.01, DMSO).

10

15

 $C_{20}H_{20}N_2O_4F_2$  (MW = 390.39); mass spectroscopy (MH<sup>+</sup>) 391.

## Example 112

## Synthesis of N-[N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-alaninyl]-L-phenylglycine Methyl Ester

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(L-alaninyl)-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 243-245°C). The reaction was monitored by tlc (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.19$  (d, 3H); 1.24 (d, 3H). Optical Rotation:  $[\alpha]_{20} = +38.2^{\circ}$  @ 589 nm (c = 1.02, DMSO).  $C_{23}H_{25}N_3O_5F_2$  (MW = 461.46); mass spectroscopy (MH<sup>+</sup>) 461.

### Example 113

20

25

30

## Synthesis of N-(2-Phenylethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(2-phenylethyl)-L-alaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and phenethylamine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 241-243°C). The reaction was monitored by tlc (Rf = 0.3 in 8% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 8% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.14$  (d, 3H); 1.21 (d, 3H).

Optical Rotation:  $[\alpha]_{20} = -33.7^{\circ} @ 589 \text{ nm (c} = 1.00, DMSO).$ 

 $C_{22}H_{25}N_3O_3F_2$  (MW = 417.45); mass spectroscopy (MH<sup>+</sup>) 417.

5

10

15

### Example 114

## Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-tryptophanamide

Following General Procedure C and using 3,5-difluorophenylacetic acid (Oakwood) and N'-(L-alaninyl)-L-tryptophanamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-tryptophanamide hydrochloride (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 199-202°C). The reaction was monitored by tlc (Rf = 0.3 in 15% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 15% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.17$  (d, 3H); 4.26 (m, 1H); 4.44 (m, 1H). Optical Rotation:  $[\alpha]_{20} = -31.0^{\circ}$  @ 589 nm (c = 1.05, DMSO).  $C_{22}H_{22}N_4O_3F_2$  (MW = 428.44); mass spectroscopy (MH<sup>+</sup>) 428.

20

### Example 115

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-cyclohexylpropionate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and methyl (S)-2-amino-3cyclohexylpropionate (Novabiochem), the title compound was prepared as a
solid (mp = 116-119°C). The reaction was monitored by tlc (Rf = 0.4 in 4%
MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using
4% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1chlorobutane/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.22$  (d, 3H); 3.62 (s, 3H).

Optical Rotation:  $[\alpha]_{20} = -21.2^{\circ}$  @ 589 nm (c = 1.01, DMSO).

 $C_{21}H_{27}N_2O_4F_2$  (MW = 410.46); mass spectroscopy (MH<sup>+</sup>) 411.

5

25

30

## Example 116

# Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4-nitrophenyl)propionamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L
alanine (from Example B2 above) and N-(2-methoxyethyl)-(S)-2-amino-3-(4nitrophenyl)propionamide hydrochloride (prepared from N-BOC-L-4nitrophenylalanine (Advanced Chemtech) and 2-methoxyethylamine (Aldrich)
using General Procedure C, followed by removal of the BOC-group using
General Procedure P), the title compound was prepared as a solid (mp = 263265°C). The reaction was monitored by tlc (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>)
and the product was purified by silica gel chromatography using 10%
MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from
EtOH/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.15$  (d, 3H); 4.23 (m, 1H); 4.54 (m, 1H). Optical Rotation:  $[\alpha]_{20} = -19.9^{\circ}$  @ 589 nm (c = 1.00, DMSO).  $C_{23}H_{26}N_4O_6F_2$  (MW = 492.48); mass spectroscopy (MH<sup>+</sup>) 493.

#### Example 117

Synthesis of

N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-serine Ethyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (prepared from 3-nitrophenylacetic acid (Aldrich) and L-alanine ethyl ester hydrochloride (Sigma) using General Procedure C, followed by hydrolysis using General Procedure AF) and L-serine ethyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 179-181°C). The reaction was

10

15

20

monitored by tlc (Rf = 0.2 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (m, 6H); 4.30 (m, 1H); 4.41 (m, 1H); 5.04 (t, 1H).

Optical Rotation:  $[\alpha]_{20} = -19.7^{\circ}$  @ 589 nm (c = 1.01, DMSO).  $C_{16}H_{21}N_3O_7$  (MW = 367.36); mass spectroscopy (MH<sup>+</sup>) 368.

#### Example 118

### Synthesis of N-[(R)- $\alpha$ -Methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(R)- $\alpha$ -methylbenzyl-L-alaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and (R)- $\alpha$ -methylbenzylamine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 240-242°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 9% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.19$  (t, 6H); 1.31 (d, 3H). Optical Rotation:  $[\alpha]_{20} = +1.0^{\circ}$  @ 589 nm (c = 1.00, DMSO).  $C_{22}H_{25}N_3O_3F_2$  (MW = 417.45); mass spectroscopy (MH<sup>+</sup>) 417.

25

#### Example 119

## Synthesis of N-[(S)- $\alpha$ -Methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-30 alanine (from Example B2 above) and N-(S)- $\alpha$ -methylbenzyl-L-alaninamide

15

20

hydrochloride (prepared from *N*-BOC-L-alanine (Sigma) and (R)- $\alpha$ -methylbenzylamine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 293-295°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (m, 6H); 1.30 (d, 3H).

10 Optical Rotation:  $[\alpha]_{20} = -65.9^{\circ}$  @ 589 nm (c = 1.05, DMSO).

 $C_{22}H_{25}N_3O_3F_2$  (MW = 417.45); mass spectroscopy (MH<sup>+</sup>) 417.

#### Example 120

### Synthesis of N-(4-Fluorobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(4-fluorobenzyl)-L-alaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 4-fluorobenzylamine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 257-259°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 9% MeOH/CHCl<sub>3</sub> as the eluent, followed by trituration using 1-chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (m, 6H); 3.52 (s, 2H).

Optical Rotation:  $[\alpha]_{20} = -28.7^{\circ}$  @ 589 nm (c = 1.00, DMSO).

 $C_{21}H_{22}N_3O_3F_3$  (MW = 421.42); mass spectroscopy (MH<sup>+</sup>) 421.

30

#### Example 121

#### Synthesis of N-(4-Pyridylmethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-5 alanine (from Example B2 above) and N-(4-pyridylmethyl)-L-alaninamide dihydrochloride (prepared from N-BOC-L-alanine (Sigma) and 4-(aminomethyl)pyridine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was 10 prepared as a solid (mp = 244-247°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

15 <sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.21$  (d, 3H); 1.26 (d, 3H). Optical Rotation:  $[\alpha]_{20} = -30.3^{\circ} @ 589 \text{ nm (c} = 1.00, DMSO).$  $C_{20}H_{22}N_4O_3F_2$  (MW = 404.42); mass spectroscopy (MH<sup>+</sup>) 405.

#### Example 122

#### Synthesis of N-(4-Trifluoromethylbenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and N-(4-trifluoromethylbenzyl)-Lalaninamide hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 4-(trifluoromethyl)benzylamine (Aldrich) using General Procedure C, followed by 25 removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 244-247°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 8% MeOH/CHCl<sub>3</sub> as the eluent, followed by triturated using 1-chlorobutane.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 3.52$  (s, 2H); 4.35 (d, 2H). Optical Rotation:  $[\alpha]_{20} = -27.4^{\circ}$  @ 589 nm (c = 1.05, DMSO).  $C_{22}H_{22}N_3O_3F_5$  (MW = 471.43); mass spectroscopy (MH<sup>+</sup>) 471.

#### Example 123

5

10

15

### Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-phenylpropionate

Following General Procedure C and using 3,5-difluorophenylacetic acid (Oakwood) and ethyl N-(L-alaninyl)-2-amino-2-phenylpropionate hydrochloride (prepared from N-BOC-L-alanine (Sigma) and D,L- $\alpha$ -methylphenylglycine ethyl ester (from Example D9 above) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 128-130°C). The reaction was monitored by tlc (Rf = 0.2 in 3% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 1.72$ , 1.77 (singlets, 3H); 3.52 (s, 2H).

 $C_{22}H_{24}N_2O_4F_2$  (MW = 418.44); mass spectroscopy (MH<sup>+</sup>) 418.

20

25

#### Example 124

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylalanine tert-Butyl Ester

Following General Procedure C and using 3,5-difluorophenylacetic acid (Oakwood) and N-(L-alaninyl)-L-phenylalanine tert-butyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylalanine tert-butyl ester hydrochloride (Advanced Chemtech) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a gel. The reaction was monitored by tlc (Rf = 0.5 in 4%

MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 4% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.19$  (d, 3H); 1.30 (s, 9H).

 $C_{24}H_{28}N_2O_4F_2$  (MW = 446.50); mass spectroscopy (MH<sup>+</sup>) 446.

#### Example 125

# Synthesis of Methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-methylpropionate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-aminoisobutyrate (prepared from 2-aminoisobutyric acid (Aldrich) using General Procedure H), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.25 in CHCl<sub>3</sub>/MeOH 95:5).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.32$  (m, 3H), 7.13 (m, 1H), 7.00 (m, 2H), 4.31 (m, 1H), 3.53 (m, 5H), 7.08 (m, 1H), 1.36 (s, 3H), 1.34 (s, 3H), 1.19 (d, 3H).

Optical Rotation:  $[\alpha]_{23} = -25^{\circ}$  (c 1, MeOH).

20  $C_{16}H_{20}N_2O_4F_2$  (MW = 342.34); mass spectroscopy (MH<sup>+</sup>) 343.

#### Example 126

# Synthesis of Ethyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]2-amino-2-cyclohexylacetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-cyclohexylacetate hydrochloride (prepared from cyclohexylglycine (Advanced Chemtech) using General Procedure H), the title compound was prepared as a solid (mp = 146-150°C). The reaction was monitored by tlc (Rf = 0.3 in 3% MeOH/CHCl<sub>3</sub>)

10

15

and the product was purified by silica gel chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 1.60$  (m, 6H); 3.50 (s, 2H).

 $C_{21}H_{28}N_2O_4F_2$  (MW = 410.46); mass spectroscopy (MH<sup>+</sup>) 410.

#### Example 127

Synthesis of N-(2-Methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-(2-methoxyethyl)-L-phenylglycinamide hydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and 2-methoxyethylamine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 252-254°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.22$  (d, 3H); 5.43 (d, 1H). Optical Rotation:  $[\alpha]_{20} = +6.17^{\circ}$  @ 589 nm (c = 1.04, DMSO).  $C_{22}H_{25}N_3O_4F_2$  (MW = 433.46); mass spectroscopy (MH<sup>+</sup>) 434.

#### Example 128

25

Synthesis of N-[N-(Isovaleryl)-2-amino-2-cyclohexylacetyl]-L-alanine Ethyl Ester

Following General Procedure C and using N-(isovaleryl)-2-amino-2-cyclohexylacetic acid (prepared from isovaleric acid (Aldrich) and D,L- $\alpha$ -

15

20

cyclohexylglycine ethyl ester hydrochloride (prepared from cyclohexylglycine (Advanced Chemtech) and ethanol using General Procedure H) using General Procedure C, followed by removal of the BOC-group using General Procedure P) and L-alanine ethyl ester hydrochloride (Sigma), the title compound was prepared as a solid (mp = 220-224°C). The reaction was monitored by tlc (Rf = 0.2 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

#### Example 129

### Synthesis of N-2-(N,N-Dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-2-(N,N-dimethylamino)ethyl-L-phenylglycinamide dihydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and N,N-dimethylethylenediamine (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 234-236°C). The reaction was monitored by tlc (Rf = 0.3 in 15% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub>, followed by slurrying in acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.22$  (d, 3H); 5.41 (d, 1H). Optical Rotation:  $[\alpha]_{20} = +5.7^{\circ}$  @ 589 nm (c = 1.01, DMSO).  $C_{23}H_{28}N_4O_3F_2$  (MW = 446.50); mass spectroscopy (MH<sup>+</sup>) 446.

25

30

#### Example 130

## Synthesis of N-(2-Pyridylmethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and N-(2-pyridylmethyl)-L-phenylglycinamide
dihydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech)
and 2-(aminomethyl)pyridine (Aldrich) using General Procedure C, followed by
removal of the BOC-group using General Procedure P), the title compound was
prepared as a solid (mp = 272-275°C). The reaction was monitored by tlc (Rf
= 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel
chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by
recrystallization from acetonitrile.

NMR data was as follows:

15 
<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.24$  (d, 3H); 5.50 (d, 1H).

Optical Rotation:  $[\alpha]_{20} = +12.4^{\circ}$  @ 589 nm (c = 1.02, DMSO).  $C_{25}H_{24}N_4O_3F_2$  (MW = 466.49); mass spectroscopy (MH<sup>+</sup>) 467.

#### Example 131

Synthesis of N-[N-(3-Pyridylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure C and using 3-pyridylacetic acid hydrochloride (Aldrich) and N-(L-alaninyl)-L-phenylalanine methyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylalanine methyl ester hydrochloride (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 150-152°C). The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.16$  (d, 3H); 347 (s, 2H). Optical Rotation:  $[\alpha]_{20} = -19.0^{\circ}$  @ 589 nm (c = 1.03, DMSO).  $C_{20}H_{23}N_3O_4$  (MW = 369.42); mass spectroscopy (MH<sup>+</sup>) 369.

#### Example 132

5

10

15

### Synthesis of N-[N-(2-Pyridylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure C and using 2-pyridylacetic acid hydrochloride (Aldrich) and N-(L-alaninyl)-L-phenylalanine methyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylalanine methyl ester hydrochloride (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 137-139°C). The reaction was monitored by tlc (Rf = 0.4 in 8% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 8% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.17$  (d, 3H); 3.65 (s, 2H). Optical Rotation:  $[\alpha]_{20} = -17.48^{\circ}$  @ 589 nm (c = 1.09, DMSO).  $C_{20}H_{23}N_3O_4$  (MW = 369.42); mass spectroscopy (MH<sup>+</sup>) 369.

20

25

#### Example 133

### Synthesis of N-[N-(4-Pyridylacetyl)-L-alaninyl]-L-phenylalanine Methyl Ester

Following General Procedure C and using 4-pyridylacetic acid hydrochloride (Aldrich) and N-(L-alaninyl)-L-phenylalanine methyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and L-phenylalanine methyl ester hydrochloride (Sigma) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 152-154°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel

10

15

25

30

chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.17$  (d, 3H); 3.47 (s, 2H).

Optical Rotation:  $[\alpha]_{20} = -17^{\circ}$  @ 589 nm (c = 1.00, DMSO).

 $C_{20}H_{23}N_3O_4$  (MW = 369.42); mass spectroscopy (MH<sup>+</sup>) 369.

#### Example 134

## Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-2-amino-2-(4-fluorophenyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-(4-fluorophenyl)acetate hydrochloride (prepared from 4-fluorophenylglycine (Fluka) and ethanol using General Procedure H), the title compound was prepared as a solid (mp = 169-183°C). The reaction was monitored by tlc (Rf = 0.3 in 4% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 4% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 3.49$ , 3.53 (singlets, 2H); 5.40 (m, 1H).

 $C_{21}H_{21}N_2O_4F_3$  (MW = 422.4); mass spectroscopy (MH<sup>+</sup>) 422.

#### Example 135

## Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-fluorophenyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-(2-fluorophenyl)acetate hydrochloride (prepared from 2-fluorophenylglycine (Fluka) and ethanol using General Procedure H), the title compound was prepared as a solid (mp = 153-

170°C). The reaction was monitored by tlc (Rf = 0.3 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 3.50$ , 3.54 (singlets, 2H), 5.66 (m, 1H).

 $C_{21}H_{21}N_2O_4F_3$  (MW = 422.40); mass spectroscopy (MH<sup>+</sup>) 422.

#### Example 136

10

15

20

5

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-phenylglycinyl]-L-alanine Ethyl Ester

Following General Procedure C and using 3,5-difluorophenylacetic acid (Oakwood) and N-(L-phenylglycinyl)-L-alanine ethyl ester hydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and L-alanine ethyl ester hydrochloride (Aldrich) using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.3 in 3% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 3.50$  (s, 2H), 5.53 (d, 1H).  $C_{21}H_{22}N_2O_4F_2$  (MW = 404.42); mass spectroscopy (MH<sup>+</sup>) 405.

25

30

#### Example 137

## Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-3-phthalimidopropionate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-3-phthalimidopropionate

hydrochloride (from Example D10 above), the title compound was prepared as a solid (mp = 197-201°C). The reaction was monitored by tlc (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 7.88$  (m, 4H), 8.29 (t, 1H), 8.48, 8.55 (doublets, 1H).

 $C_{24}H_{23}N_3O_6F_2$  (MW = 487.46); mass spectroscopy (MH<sup>+</sup>) 487.

#### Example 138

10

25

5

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine Neopentyl Ester

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and L-phenylglycine neopentyl ester

hydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech)
and 2,2-dimethyl-1-propanol (Aldrich) using General Procedure C (with
catalytic DMAP), followed by removal of the BOC-group using General
Procedure P), the title compound was prepared as a solid (mp = 133-136°C).
The reaction was monitored by tlc (Rf = 0.7 in 10% MeOH/CHCl<sub>3</sub>) and the
product was purified by silica gel chromatography using 4% MeOH/CHCl<sub>3</sub> as
the eluent, followed by recrystallization from 1-chlorobutane/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 3.50$  (s, 2H), 5.42 (d, 1H). Optical Rotation:  $[\alpha]_{20} = +45.9^{\circ}$  @ 589 nm (c = 1.02, DMSO).  $C_{24}H_{28}N_2O_4F_2$  (MW = 446.50); mass spectroscopy (MH<sup>+</sup>) 446.

Example 139

Synthesis of N-tert-Butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

15

20

25

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), S-(+)- $\alpha$ -methylbenzylamine (Aldrich), benzaldehyde (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 233-235°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 8% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ )(1:1 mixture of diastereomers):  $\delta = 3.52$  (s, 2H), 5.40 (m, 1H).

 $C_{23}H_{27}N_3O_3F_2$  (MW = 431.49); mass spectroscopy (MH<sup>+</sup>) 432.

#### Example 140

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylglycine ten-butyl ester hydrochloride (Advanced Chemtech), the title compound was prepared as a solid (mp = 145-147°C). The reaction was monitored by tlc (Rf = 0.5 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 2.5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1-chlorobutane/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.26$  (d, 3H); 5.20 (d, 1H). Optical Rotation: [α]<sub>20</sub> = +14.8° @ 589 nm (c =1.01, MeOH).  $C_{23}H_{26}N_2O_4F_2$  (MW = 432.47); mass spectroscopy (MH+) 433.

#### Example 141

 $Synthesis of \\ N'-[N-(3,5-Difluor ophenylacetyl)-L-alaninyl]-L-phenylglycinamide$ 

15

20

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylglycinamide hydrochloride (prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and ammonia using General Procedure C, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 288-290°C). The reaction was monitored by tlc (Rf = 0.4 in 15% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 15% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from EtOH.

NMR data was as follows:

#### Example 142

### Synthesis of 4-[N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-valinyl]morpholine

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and 4-(L-valinyl)morpholine (prepared from N-BOC-L-valine (Aldrich) and morpholine (Aldrich) using General Procedure M, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.5 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 98:2 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.12$  (d, 2H), 8.08 (dd, 1H), 7.59 (d, 1H, J=7 Hz), 7.42 (t, 1H), 7.32 (d, J=8 Hz, 1H), 7.03(d, J=8 Hz, 1H), 4.78 (m, 1H), 4.68 (m, 1H), 3.61 (m, 10H), 1.90 (m, 1H), 1.96 (d, 3H), 1.31 (d, 3H), 0.88 (d, 3H), 0.80 (d, 3H).

Optical Rotation:  $[\alpha]_{23} = -5^{\circ}$  (c 5, MeOH).

### Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-valine Ethyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and L-valine ethyl ester hydrochloride (Aldrich), the title compound was prepared as a solid. The reaction was monitored by the (Rf = 0.2 in 97:3 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 97:3 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 8.13 (m, 2H), 7.62 (d, J=7 Hz, 1H), 7.47 (t, 1H), 6.52 (m, 2H), 4.57 (m, 1H), 4.46 (m, 1H), 4.19 (m, 2H), 3.65 (s, 2H), 2.13 (m, 1H), 1.38 (d, 3H), 1.22 (t, 3H), 0.82 (d, 3H).

Optical Rotation:  $[\alpha]_{23} = -24.3^{\circ}$  @ 589 nm (c 1, DMSO).  $C_{18}H_{25}N_3O_6$  (MW = 379.42); mass spectroscopy (MH<sup>+</sup>) 380.

15

20

5

#### Example 144

### Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-threonine Methyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and L-threonine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.1 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 95:5 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.08$  (d, 1H), 7.96 (d, 1H), 7.59 (d, 1H), 7.45 (d, 1H), 7.34 (t, 1H), 7.20 (d, 1H), 4.43 (m, 1H), 4.39 (dd, 1H), 4.13 (m, 1H), 3.59 (s, 3H), 3.51 (s, 2H), 1.20 (d, 3H), 1.03 (d, 3H).

Optical Rotation:  $[\alpha]_{23} = -20.8^{\circ}$  (c 5, MeOH).  $C_{16}H_{20}N_2O_7$  (MW = 367.3); mass spectroscopy (MH<sup>+</sup>) 368.

# Synthesis of 4-[N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-(S)-2-amino-3-tert-butoxybutyryl]morpholine

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and 4-[(S)-2-amino-3-tert-butoxybutyryl]-morpholine (prepared from N-BOC-O-tert-butyl-L-threonine (Sigma) and morpholine (Aldrich) using General Procedure M, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.1 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 96:4 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.12$  (m, 2H), 7.66 (d, 1H), 7.47 (t, 1H), 6.88 (d, 1H), 6.32 (d, 1H), 4.78 (m, 1H), 4.50 (m, 1H), 3.90-3.40 (m, 11H), 1.40 (d, 3H), 1.18 (s, 9H), 1.0 (d, 3H).

 $C_{23}H_{33}N_3O_7$  (MW = 478.5); mass spectroscopy (MH<sup>+</sup>) 479.

#### Example 146

#### Synthesis of

4-[N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-isoleucinyl]morpholine

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and 4-(L-isoleucinyl)morpholine (prepared from N-BOC-L-isoleucine (Aldrich) and morpholine (Aldrich) using General Procedure M, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp =  $156-160^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.45 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 98:2 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

20

25

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.16$  (d, 1H), 8.09 (d, 1H), 7.63 (d, 1H), 7.45 (t, 1H), 7.30 (d, 1H), 6.89 (d, 1H), 4.78 (m, 1H), 4.62 (m, 1H), 3.6 (m, 10H), 1.65 (m, 1H), 1.4 (m, 1H), 1.29 (d, 3H), 1.03 (d, 3H), 0.90-0.76 (m, 6H).

10

Optical Rotation:  $[\alpha]_{23} = -55^{\circ}$  @ 589 nm (c 1, MeOH).

#### Example 147

### Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]L-isoleucine Methyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and L-isoleucine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.15 in 97:3 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 97:3 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.12$  (m, 2H), 7.66 (d, 1H), 7.49 (t, 1H), 6.50 (m, 2H), 4.52 (m, 2H), 3.72 (s, 3H), 3.61 (s, 2H), 1.87 (m, 1H), 1.32 (m, 4H), 1.07 (m, 1H), 0.81(d, 6H).

15 Optical Rotation:  $[\alpha]_{23} = -7.3^{\circ}$  (c 5, MeOH).  $C_{18}H_{25}N_2O_6$  (MW = 379); mass spectroscopy (MH<sup>+</sup>) 379.

#### Example 148

### Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-isoleucine

Following General Procedure AF and using N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine methyl ester (from Example 147 above), the title compound was prepared as a solid.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.41$  (d, 1H), 8.15 (s, 1H), 8.07 (d, 1H), 7.91 25 (d, 1H), 7.68 (d, 1H), 7.53 (t, 1H), 4.36 (m, 1H), 4.12 (m, 1H), 3.62 (s, 2H), 1.71 (m, 1H), 1.31 (m, 1H), 1.18 (d, 3H), 1.07 (m, 1H), 0.79 (m, 6H). Optical Rotation:  $[\alpha]_{23} = -42^{\circ}$  (c 5, MeOH).  $C_{17}H_{23}N_2O_6$  (MW = 365.3); mass spectroscopy (MH<sup>+</sup>) 366.

PCT/US97/20804 WO 98/22494

-- 268 --

#### Example 149

#### Synthesis of N-[N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-threoninyl]-L-valine Ethyl Ester

5 Following General Procedure C and using N-[N-(3-nitrophenylacetyl)-Lalaninyl]-L-threonine (prepared from N-[N-(3-nitrophenylacetyl)-L-alaninyl]-Lthreonine methyl ester (from Example 144 above) using General Procedure AF) and L-valine ethyl ester hydrochloride (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.1 in 96:4 10 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 96:4 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

15

20

25

30

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.12$  (m, 1H), 7.60 (d, 1H), 7.48 (t, 1H), 7.05 (d, 1H), 6.98 (d, 1H), 6.48 (d, 1H), 4.60 (m, 1H), 4.47 (m, 3H), 4.22 (m, 2H) 3.65 (s, 2H), 2.19 (m, 1H), 1.38 (d, 3H), 1.28 (t, 3H), 1.09 (d, 3H), 0.87 (m, 6H).

Optical Rotation:  $[\alpha]_{23} = -85^{\circ}$  (c 5, MeOH).

#### Example 150

#### Synthesis of Methyl N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and methyl (S)-2-aminopentanoate hydrochloride (prepared from (S)-2-aminopentanoic acid (Novabiochem) using General Procedure H), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.4 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.39$  (m, 1H), 8.28 (m, 1H), 8.19 (m, 1H), 8.11 (m, 1H), 7.73 (d, 1H), 7.61 (d, 1H), 4.36 (m, 1H), 4.22 (m, 1H), 3.64 (m, 5H), 1.62 (m, 2H), 1.26 (m, 2H), 1.22 (d, 3H), 0.86 (m, 3H). Optical Rotation:  $[\alpha]_{23} = -29^{\circ}$  (c 1, MeOH).

 $C_{17}H_{23}N_3O_6$  (MW = 365); mass spectroscopy (MH<sup>+</sup>) 366.

#### Example 151

### Synthesis of N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-leucine Methyl Ester

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine (from Example D11 above) and L-leucine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.75 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 97:3 CHCl<sub>3</sub>/MeOH as the eluent.

10 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 8.12 (m, 2H), 8.04 (m, 1H), 7.58 (m, 1H), 7.48-7.30 (m, 2H), 7.11 (d, 1H), 4.63 (m, 1H), 4.48 (m, 1H), 3.68 (s, 2H), 3.64 (s, 3H), 1.63 (m, 1H), 1.31 (m, 2H), 0.85 (d, 3H), 0.82 (m, 3H).

Optical Rotation:  $[\alpha]_{23} = -32^{\circ}$  (c 1, MeOH).

15  $C_{18}H_{25}N_3O_6$  (MW = 379); mass spectroscopy (MH<sup>+</sup>) 380.

#### Example 152

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-leucine Methyl Ester

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-leucine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.5 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.78$  (m, 2H), 6.69 (m, 1H), 4.52 (m, 2H), 3.73 (m, 1H), 3.52 (d, 2H), 1.63 (m, 2H), 1.36 (m, 3H), 0.88 (m, 3H). Optical Rotation:  $[\alpha]_{23} = -34^{\circ}$  (c 1, MeOH).  $C_{18}H_{24}N_2O_4F_2$  (MW = 370); mass spectroscopy (MH<sup>+</sup>) 370.

# Synthesis of N-2-Methoxyethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and 2-methoxyethylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.35 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

> Optical Rotation:  $[\alpha]_{23} = -50^{\circ}$  (c 1, MeOH).  $C_{17}H_{23}N_3O_4F_2$  (MW = 371); mass spectroscopy (MH<sup>+</sup>) 372.

15

#### Example 154

### Synthesis of N-2-(N,N-Dimethylamino) ethyl-N'-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-20 L-alaninyl]-L-alanine (from Example D7 above) and N,N-dimethyl-ethylenediamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.05 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.38$  (m, 1H), 8.02 (m, 1H), 7.66 (m, 1H), 7.09 (m, 1H), 6.97 (m, 2H), 4.22 (m, 2H), 3.53 (s, 2H), 3.08 (m, 2H), 2.22 (m, 2H), 2.11 (m, 6H), 1.21 (d, 6H).

Optical Rotation:  $[\alpha]_{23} = -55^{\circ}$  (c 1, MeOH).  $C_{18}H_{26}N_4O_3F_2$  (MW = 384); mass spectroscopy (MH<sup>+</sup>) 384.

### Synthesis of N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and cyclohexylamine (Aldrich), the title compound was prepared as a solid (mp = 239-244 °C). The reaction was monitored by tlc (Rf = 0.25 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

10 ¹H-nmr (DMSO- $d_6$ ): δ = 8.39 (m, 1H), 7.94 (m, 1H), 7.56 (m, 1H), 7.08 (m, 1H), 6.97 (m, 2H), 4.20 (m, 2H), 3.32 (s, 2H), 3.27 (m, 1H), 1.64 (m, 4H), 1.54 (m, 2H), 1.20 (m, 10H).

Optical Rotation:  $[\alpha]_{23} = -58^{\circ}$  (c 1, MeOH).

 $C_{20}H_{27}N_3O_3F_2$  (MW = 395); mass spectroscopy (MH<sup>+</sup>) 395.

15

#### Example 156

### Synthesis of N-Neopentyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)
L-alaninyl]-L-alanine (from Example D7 above) and neopentylamine (Aldrich),
the title compound was prepared as a solid. The reaction was monitored by the
(Rf = 0.25 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.37$  (d, 1H), 8.01 (m, 1H), 7.67 (m, 1H), 7.11 (m, 1H), 6.98 (m, 2H), 4.28 (m, 2H), 3.51 (s, 2H), 2.88 (m, 2H), 1.23 (d, 3H), 0.80 (m, 9H).

Optical Rotation:  $[\alpha]_{23} = -54^{\circ}$  (c 1, MeOH).

 $C_{19}H_{27}N_3O_3F_2$  (MW = 383); mass spectroscopy (MH<sup>+</sup>) 383.

## Synthesis of N-Tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and tetrahydrofurfurylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = <math>0.20 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

Optical Rotation:  $[\alpha]_{23} = -70^{\circ}$  (c 1, MeOH).

 $C_{19}H_{25}N_3O_4F_2$  (MW = 397); mass spectroscopy (MH<sup>+</sup>) 398.

15

#### Example 158

# Synthesis of N-2-Pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)
L-alaninyl]-L-alanine (from Example D7 above) and 2-(aminomethyl)pyridine
(Aldrich), the title compound was prepared as a solid. The reaction was
monitored by tlc (Rf = 0.1 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.49$  (m, 1H), 8.41 (m, 2H), 8.14 (d, 1H), 7.74 (m, 1H), 7.28 (m, 2H), 7.09 (m, 1H), 6.98 (m, 2H), 4.33 (m, 4H), 3.52 (s, 2H), 1.24 (m, 6H).

Optical Rotation:  $[\alpha]_{23} = -68^{\circ}$  (c 5, MeOH).  $C_{20}H_{22}N_4O_3F_2$  (MW = 404); mass spectroscopy (MH<sup>+</sup>) 405.

### Synthesis of 3-[N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-alaninyl]thiazolidine

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and thiazolidine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.25 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

> Optical Rotation:  $[\alpha]_{23} = -67^{\circ}$  (c 1, MeOH).  $C_{17}H_{21}N_3O_3F_2$  (MW = 385); mass spectroscopy (MH<sup>+</sup>) 385.

15

#### Example 160

# Synthesis of Methyl N-[N-(3,5-difluorophenylacetyl)L-alaninyl]-(S)-2-aminobutanoate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and methyl (S)-2-aminobutanoate
hydrochloride (prepared from (S)-(+)-2-aminobutyric acid (Aldrich) using
General Procedure H), the title compound was prepared as a solid (mp = 103106°C).

NMR data was as follows:

25  $^{1}$ H-nmr (CDCl<sub>3</sub>):  $\delta = 6.83$  (m, 2H), 6.72 (m, 1H), 6.49 (d, 1H), 4.55 (m, 1H), 4.48 (m, 1H), 3.72 (s, 3H), 3.49 (s, 2H), 1.85 (m, 1H), 1.69 (m, 1H), 1.39 (d, 3H), 0.86 (t, 3H).

Optical Rotation:  $[\alpha]_{23} = -70^{\circ}$  (c 1, MeOH).  $C_{16}H_{20}N_2O_4F_2$  (MW = 342.35); mass spectroscopy (MH<sup>+</sup>) 342.

### Synthesis of Methyl N-[N-(3,5-difluorophenylacetyl)L-alaninyl]-(S)-2-aminopentanoate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (S)-2-aminopentanoate hydrochloride (prepared from (S)-2-aminopentanoic acid (Novabiochem) using General Procedure H), the title compound was prepared as a solid (mp = 154-155°C).

10 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.80$  (m, 2H), 6.69 (m, 1H), 6.45 (d, 1H), 6.28 (d, 1H), 4.52 (m, 2H), 3.71 (s, 3H), 3.51 (s, 2H), 1.77 (m, 1H), 1.58 (m, 1H), 1.35 (d, 3H), 1.27 (m, 2H), 0.87 (t, 3H).

Optical Rotation:  $[\alpha]_{23} = -69^{\circ}$  (c 1, MeOH).

15

30

#### Example 162

### Synthesis of Methyl N-[N-(3-nitrophenylacetyl)-L-alaninyl](S)-2-aminobutanoate

Following General Procedure C and using N-(3-nitrophenylacetyl)-L-alanine

(from Example D11 above) and methyl (S)-2-aminobutanoate hydrochloride

(prepared from (S)-(+)-2-aminobutyric acid (Aldrich) using General Procedure

H), the title compound was prepared as a solid (mp = 154-157°C).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.13$  (m, 1H), 8.04 (m, 1H), 7.57 (m, 1H), 7.38 (m, 1H), 4.72 (m, 1H), 4.39 (m, 1H), 3.69 (s, 3H), 3.41 (s, 2H), 1.73 (m, 1H), 1.61 (m, 1H), 1.34 (d, 3H), 0.79 (t, 3H).

Optical Rotation:  $[\alpha]_{23} = -75^{\circ}$  (c 1, MeOH).

#### Example 163

Synthesis of N-(R)-sec-Butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

WO 98/22494 PCT/US97/20804

-- 275 --

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and (R)-(-)-sec-butylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.15 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 95:5 CHCl<sub>3</sub> MeOH as the eluent.

NMR data was as follows:

5

15

20

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.39$  (m, 1H), 7.95 (m, 1H), 7.49 (m, 1H), 7.09 (m, 1H), 7.01 (m, 2H), 4.20 (m, 4H), 3.61 (m, 1H), 3.52 (s, 2H), 1.34 (m, 2H), 1.21 (m, 6H), 0.97 (d, 3H), 0.79 (m, 3H).

10 Optical Rotation:  $[\alpha]_{23} = -50^{\circ}$  (c 1, MeOH).  $C_{18}H_{25}N_3O_3F_2$  (MW = 369.41); mass spectroscopy (MH<sup>+</sup>) 370.

#### Example 164

### Synthesis of 1-[N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-alaninyl]pyrrolidine

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and pyrrolidine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.15 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.31$  (m, 1H), 8.08 (m, 1H), 7.09 (m, 1H), 6.99 (m, 2H), 4.48 (m, 1H), 4.29 (m, 1H), 3.51 (s, 2H), 3.44-3.22 (m, 4H), 1.80 (m, 4H), 1.27 (m, 6H).

25  $C_{18}H_{23}N_3O_3F_2$  (MW = 367.40); mass spectroscopy (MH<sup>+</sup>) 367.

#### Example 165

Synthesis of N-(S)-sec-Butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and (S)-(+)-sec-butylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.25 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.38$  (m, 1H), 7.92 (m, 1H), 7.30 (m, 1H), 7.18 (m, 1H), 6.99 (m, 2H), 4.20 (m, 4H), 3.62 (m, 1H), 3.52 (s, 2H), 1.34 (m, 2H), 1.20 (m, 6H), 1.01 (m, 3H), 0.81 (t, 3H).

10 Optical Rotation:  $[\alpha]_{23} = -52^{\circ}$  (c 1, MeOH).  $C_{19}H_{25}N_3O_3F_2$  (MW = 369.41); mass spectroscopy (MH<sup>+</sup>) 370.

#### Example 166

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-valine Methyl Ester

15

5

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-valine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.81$  (m, 2H), 6.73 (m, 1H), 6.48 (d, 1H), 6.22 (d, 1H), 4.48 (m, 2H), 3.70 (s, 3H), 3.51 (s, 2H), 2.16 (m, 1H), 1.37 (m, 1H), 0.87 (t, 3H).

Optical Rotation:  $[\alpha]_{23} = -65^{\circ}$  (c 1, MeOH).  $C_{17}H_{22}N_2O_4F_2$  (MW = 356.37); mass spectroscopy (MH<sup>+</sup>) 360.

25

30

#### Example 167

## Synthesis of N-2-Fluoroethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine (from Example D7 above) and 2-fluoroethylamine

hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 230-235°C).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.38$  (d, 1H), 8.04 (m, 2H), 7.07 (m, 1H), 6.99 (m, 2H), 4.39 (m, 2H), 4.24 (m, 1H), 3.53 (s, 2H), 3.35 (m, 2H), 1.20 (m, 6H).

Optical Rotation:  $[\alpha]_{23} = -33^{\circ}$  (c 1, MeOH).  $C_{16}H_{20}N_3O_3F_3$  (MW = 359.37); mass spectroscopy (MH<sup>+</sup>) 359.

#### Example 168

10

15

5

# Synthesis of N-[(S)-6-Methyl-3-oxohept-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure M and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and (S)-6-methyl-3-oxohept-2-ylamine hydrochloride (prepared by treating N-BOC-L-alanine N-methoxy-N-methyl amide (Weinreb et al., Tetrahedron Lett., 22, 3815 (1981)) with isopropyl magnesium bromide (Aldrich), followed by removal of the BOC group using General Procedure P), the title compound was prepared as a solid. The product was purified by silica gel chromatography using CHCl<sub>3</sub>/MeOH as the eluent.

20 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.84$  (m, 2H), 6.69 (m, 1H), 6.31 (m, 1H), 4.50 (m, 2H), 3.51 (s, 2H), 2.48 (m, 2H), 1.47 (m, 2H), 1.32 (m, 7H), 0.90 (d, 6H).

Optical Rotation:  $[\alpha]_{23} = -42^{\circ}$  (c 1, MeOH).

25  $C_{19}H_{26}N_2O_3F_2$  (MW = 368); mass spectroscopy (MH<sup>+</sup>) 368.

#### Example 169

Synthesis of N-4-Nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutyramide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutyric acid (prepared from methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate (from Example 160 above) using General Procedure AF) and 4-nitrobenzylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.3 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 97:3 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.57$  (t, 1H), 8.40 (d, 1H), 8.21 (d, 2H), 8.02 (d, 1H), 7.50 (d, 2H), 7.08 (m, 1H), 6.98 (m, 2H), 4.42 (d, 2H), 4.37 (m, 1H), 4.17 (m, 1H), 3.53 (s, 2H), 1.64 (m, 2H), 1.21 (m, 3H), 0.83 (t, 3H). Optical Rotation:  $[\alpha]_{23} = -42^{\circ}$  (c 1, MeOH).  $C_{22}H_{24}N_4O_5F_2$  (MW = 462.45); mass spectroscopy (MH<sup>+</sup>) 462.

#### Example 170

15

20

5

# Synthesis of N-4-Nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminopentanamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoic acid (prepared from methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate (from Example 161 above) using General Procedure AF) and 4-nitrobenzylamine (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.3 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.57$  (m, 1H), 8.41 (d, 1H), 8.22 (d, 2H), 8.06 (d, 1H), 7.51 (d, 2H), 7.12 (m, 1H), 7.00 (m, 2H), 4.43 (d, 2H), 4.30 (m, 2H), 3.56 (s, 2H), 1.65 (m, 2H), 1.29 (m, 5H), 0.91 (t, 3H).

Optical Rotation:  $[\alpha]_{23} = +97^{\circ}$  (c 1, MeOH).

30  $C_{23}H_{26}N_4O_5F_2$  (MW = 476.4); mass spectroscopy (MH<sup>+</sup>) 476.

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]2-amino-2-(3-fluorophenyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and methyl 2-amino-2-(3-fluorophenyl)acetate
hydrochloride (from Example D12 above), the title compound was prepared as
a solid. The reaction was monitored by tlc (Rf = 0.2 in 95:5 CHCl<sub>3</sub>/MeOH)
and the product was purified by silica gel chromatography using 95:5

CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.36$  (m, 1H), 7.18 (m, 1H), 7.13 (m, 1H), 7.06 (m, 1H), 6.87 (m, 2H), 6.74 (m, 1H), 6.09 (m, 1H), 5.49 (d, 1H), 4.59 (m, 1H), 3.74 (s, 3H), 3.57 (s, 2H), 1.35 (d, 3H), 0.97 (d, 3H).

15  $C_{20}H_{19}N_2O_4F_3$  (MW = 408.38); mass spectroscopy (MH<sup>+</sup>) 408.

#### Example 172

# Synthesis of N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]- (S)-2-amino-2-(2-thienyl)acetamide

Following General Procedure L and using methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate (from Example 178 below), the title compound was prepared as a solid (mp = decomposition at 190°C). The product was purified by preparative LC 2000 chromatography using 8:2 EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>/DMSO- $d_6$ ):  $\delta = 8.9$ -6.14 (Ar + NH's 10 H), 5.43-5.39 (m, 1H), 4.16-4.10 (m, J=7 Hz, 1H), 3.19 (s, 2H), 1.15 (d, J=7.05 Hz, 3H). C<sub>17</sub>H<sub>17</sub>F<sub>2</sub>N<sub>3</sub>O<sub>3</sub>S (MW = 381.4); mass spectroscopy (MH<sup>+</sup>) 381.

#### Example 173

30

25

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-2-amino-2-(5-chlorobenzothiophen-2-yl)acetate

10

15

20

25

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(5-chlorobenzothiophen-2-yl)acetate (prepared from 5-chlorobenzothiophene-2-acetic acid [CAS No. 23799-65-7] using General Procedure G, followed by amination using a procedure essentially the same as that described in Example D4 above), the title compound was prepared as a solid (mp = 189-190°C). The product was purified by titration using Et20/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.7-7.63$  (m, 2H), 7.33-7.17 (m, 2H), 6.89-6.63 (m, 3H), 6.16-6.03 (m, 1H), 5.85 (dd, 1H), 4.7-4.53 (m, 1H), 3.83 (s, 1.5H), 3.8 (s, 1.5H), 3.59 (s, 1H), 3.5 (s, 1H), 1.4 (dt, 3H).

 $C_{22}H_{19}ClF_2N_2O_4S$  (MW = 481); mass spectroscopy (MH<sup>+</sup>) 480.

#### Example 174

### Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-2-amino-2-(benzothiophen-2-yl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and ethyl 2-amino-2-(benzothiophen-2-yl)acetate [CAS No. 98800-64-7], the title compound was prepared as a solid (mp = 189-190°C). The product was purified by preparative LC 2000 chromatography using 2:8 EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.8-7.75$  (m, 2H), 7.34-7.27 (m, 2H), 7.25-7.09 (m, 3H), 6.81-6.76 (m, 1H), 6.76-6.63 (m, 1H), 6.23 (dd, J=7 Hz, 1H), 5.84(d, J=7.07 Hz, 1H), 4.61-4.59 (m, 1H), 4.33-4.2 (m, 2H), 3.54 (s, 1H), 3.50 (s, 1H), 1.70 (d, J=11.9 Hz, 1.5H), 1.38 (d, J=11.9 Hz, 1.5 H), 1.36-1.23 (dt, 3H).

 $C_{23}H_{22}N_2O_4SF_2$  (MW = 460.49); mass spectroscopy (MH<sup>+</sup>) 460.

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-2-amino-2-(benzothiophen-3-yl)acetate

- Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(benzothiophen-3-yl)acetate (prepared from 2-amino-2-(benzothiophen-3-yl)acetic acid [CAS 95834-94-9] using General Procedure H), the title compound was prepared as a solid (mp = 185-186°C).
- 10 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.86$  (m, 2H), 7.4-7.3 (m, 3H), 7.4-7.2 (m, 2H), 6.9-6.6 (m, 3H), 6.3-6.13 (m, 1H), 5.95-5.85 (m, 1H), 4.55-4.5 (m, 1H), 3.75 (s, 1.5H), 3.65 (s, 1.5H), 3.55 (s, 1H), 3.35 (s, 1H), 1.4 (d, 1.5H), 1.3 (d, 1.5H).

15  $C_{22}H_{20}N_2O_4F_2S$  (MW = 446); mass spectroscopy (MH<sup>+</sup>) 446.

#### Example 176

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-thienyl)acetate

- Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(2-thienyl)acetate (prepared from L- $\alpha$ -2-thienylglycine (Sigma) using General Procedure G), the title compound was prepared as a solid (mp = 161-162°C). The product was purified by preparative LC 2000 chromatography using 1:4 EtOAc/hexanes.
- NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 7.3-6.65 (Ar, 7H), 6.25 (bt, 1H), 5.8 (dd, 1H), 4.68-4.5 (m, 1H), 3.85 (s, 1H), 3.75 (s, 1H), 3.52 (s, 1H), 3.5 (s, 1H), 1.35 (overlaying d, 3H).

 $C_{18}H_{18}N_2O_4F_2S$  (MW = 396); mass spectroscopy (MH<sup>+</sup>) 396.1.

# Synthesis of Ethyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-2-amino-2-(benzothiophen-5-yl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and ethyl 2-amino-2-(benzothiophen-5yl)acetate (prepared as described in S. Kukolja et al., J. Med. Chem., 1985,
28, 1896-1903), the title compound was prepared as a solid (mp = 126.5127.5°C). The product was purified by preparative LC 2000 chromatography
using 1:1 hexanes/EtOAc as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.1$  (s, 1H), 8.05 (s, 1H), 7.6-7.5 (m, 2H), 7.4-7.25 (m, 3H), 7.15 (bd, J=12 Hz, 5H), 7.05 (bd, J=12 Hz, 5H), 6.89-6.675 (m, 2H), 6.225 (bd, J=12 Hz, 5H), 6.075 (bd, J=12 Hz, 5H), 4.55 (q, J=7.5 Hz, 1H), 4.2 (dq, 2H), 3.575 (s, 1H), 3.242 (s, 1H), 1.4 (d, J=7.05 Hz, 1.5H), 1.15 (d, J=7.05 Hz, 1.5H).

 $C_{23}H_{22}N_2O_4F_2S$  (MW = 460); mass spectroscopy (MH<sup>+</sup>) 460.1.

#### Example 178

20

25

30

15

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate

Following General Procedure G and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetic acid (from Example 180 below), the title compound was prepared as a solid (mp = 180-181°C). The product was purified by preparative LC 2000 chromatography using 6:4 EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.3\text{-}6.6$  (Ar + NH, 7H), 6.37 (bd, J=7 Hz, 1H), 5.77 (d, J=7 Hz, 1H), 4.6-4.56 (m, J=7 Hz, 1H), 3.7 (s, 3H), 3.4 (s, 2H), 1.38 (d, J=7 Hz, 3H).

 $C_{18}H_{18}N_2O_4SF_2$  (MW = 396); mass spectroscopy (MH<sup>+</sup>) 396.1.

### Synthesis of tert-Butyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate

Following General Procedure J and using N-[N-(3,5-difluorophenylacety])-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetic acid (from Example 180 below), the title compound was prepared as a solid (mp = 117-118°C). The product was purified by tituraration using ether/hexanes.

NMR data was as follows:

10 ¹H-nmr (CDCl<sub>3</sub>):  $\delta$  = 7.24 (d, J=6.5 Hz, 1H), 7.05-6.63 (m, 6H), 6.19 (bd, J=7.2 Hz, 1H), 5.66 (d, J=7.5 Hz, 1H), 4.6-4.5 (m, 1H), 3.5 (s, 2H), 1.44 (s, 9H), 1.38 (d, J=7.1 Hz, 3H).

 $C_{21}H_{24}N_2O_4SF_2$  (MW = 438.5); mass spectroscopy (MH<sup>+</sup>) 438.

#### Example 180

15

20

25

30

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetic Acid

Following General Procedure M and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L- $\alpha$ -2-thienylglycine (Sigma), the title compound was prepared as a solid. The product was purified by tituration using EtOAc/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.73$  (d, J=7 Hz, 1H), 8.38 (d, J=7 Hz, 1H), 7.56-7.4 (m, 1H), 7.2-6.9 (m, 4H), 5.54 (d, J=8 Hz, 1H), 4.5-4.3 (m, 1H), 3.33 (s, 2H), 1.23 (d, J=7 Hz, 3H).

 $C_{17}H_{16}N_2O_4SF_2$  (MW = 382); mass spectroscopy (MH<sup>+</sup>) 382.

#### Example 181

Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(1H-tetrazol-5-yl)acetate

10

15

20

25

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(1H-tetrazol-5-yl)acetate (prepared from ethyl 1H-tetrazole-5-acetate [CAS 173367-99-2] using procedures essentially the same as those described in S. Kukolja, J. Med. Chem., 1985, 28, 1886-1896), the title compound was prepared as a solid. The reaction was product was purified by tituration using EtOAc/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 9.13$  (d, J=7.6 Hz, 1H), 8.39 (t, J=7 Hz, 1H), 7.1-6.95 (m, 3H), 5.9 (dd, 1H), 4.4-4.3 (m, 1H), 4.14 (q, J=7 Hz, 2H), 3.5 (s, 3H), 1.27-1.11 (m, 6H).

 $C_{16}H_{18}N_6O_4F_2$  (MW = 396.3); mass spectroscopy (MH<sup>+</sup>) 396.3.

#### Example 182

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(6-methoxy-2-naphthyl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (S)-2-amino-2-(6-methoxy-2-naphthyl)acetate (from Example D3 above), the title compound was prepared as a solid (mp = 177-178°C). The product was purified by tituration using hexanes/EtOAc.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.84$  (d, J=9 Hz, 1H), 8.4 (d, J=9 Hz, 1H), 7.90-7.76 (m, 2H), 7.247-6.90 (m, 5H), 5.5 (J=7 Hz, 1H), 4.243 (d, J=3.5 Hz, 1H), 3.86 (s, 3H), 3.6 (s, 3H), 3.29 (s, 2H), 1.26 (d, J=7.5 Hz, 3H).  $C_{25}H_{24}N_2O_5F_2$  (MW = 470.48); mass spectroscopy (MH<sup>+</sup>) 470.

#### Example 183

Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-trifluoromethylphenyl)acetate

10

15

20

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(3-trifluoromethylphenyl)acetate (prepared from 2-(hydroxyimino)-2-(3-trifluoromethylphenyl)acetic acid [CAS 179811-81-5] using General Procedures G and R above), the title compound was prepared as a solid (mp = 133-134°C). The product was purified by tituration from EtOAc/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.57-7.37$  (m, 4H), 6.8-6.6 (m, 3H), 6.05 (BA, 1H), 5.5 (A, J=7.5Hz, 1H), 3.7 (s, 1.5H), 3.675 (s, 1.5H), 3.5 (s, 1H), 3.45 (s, 1H), 1.33 (d, J=7.5 Hz, 1.5H), 1.275 (d, J=7.5 Hz, 1.5H).  $C_{21}H_{19}N_2O_4F_5$  (MW = 458.39); mass spectroscopy (MH<sup>+</sup>) 459.

#### Example 184

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4,5,6,7-tetrahydrobenzothiophen-2-yl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(4,5,6,7-tetrahydrobenzothiophen-2-yl)acetate (prepared from N-Boc-2-amino-2-(4,5,6,7-tetrahydrobenzothiophen-2-yl)acetic acid [CAS 95361-97-0] using General Procedures G above and the Boc removal procedure described in Example D3 above), the title compound was prepared as a solid. The product was purified by tituration using Et<sub>2</sub>0/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.05$  (d, J=5 Hz, 1H), 7.02 (d, J=5 Hz, 1H), 6.82-6.66 (m, 3H), 6.31 (bd, J=8 Hz, 1H), 5.66 (dd, J=7.2 Hz, 1H), 4.63-4.55 (m, 1H), 3.76 (s, 1.5H), 3.75 (s, 1.5H), 3.52 (s, 1H), 3.50 (s, 1H), 2.67-2.65 (m, 2H), 2.54-2.52 (m, 2H), 1.77-1.7 (m, 4H), 1.36 (dd, J=7 Hz, 3H).  $C_{22}H_{24}N_2O_4F_2S$  (MW = 450); mass spectroscopy (MH<sup>+</sup>) 450.

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(thieno[2,3-b]thiophen-2-yl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(thieno[2,3-b]thiophen-2-yl)acetate (from Example D4 above), the title compound was prepared as a solid. The product was purified by titruation from Et<sub>2</sub>0/hexanes.

NMR data was as follows:

10 ¹H-nmr (CDCl<sub>3</sub>): δ = 7.35 (d, J=7.5 Hz, 1H), 7.2-7.0 (m, 3H), 6.9-6.69 (m, 3H), 6.13-6.0 (m, 1H), 5.8 (dd, 1H), 4.63-4.5 (m, 1H), 3.8 (s, 3H), 3.58 (s, 1H), 3.469 (1H), 1.4 (dd, 3H).

 $C_{20}H_{18}N_2O_4F_2S_2$  (MW = 452); mass spectroscopy (MH<sup>+</sup>) 452.

#### Example 186

15

20

25

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-methylthiazol-4-yl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(2-methylthiazol-4-yl)acetate (prepared from N-Boc-2-amino-2-(2-methylthiazol-4-yl)acetic acid [CAS 105381-90-6] using General Procedure H above), the title compound was prepared as a solid. The product was purified by tituration using Et<sub>2</sub>0/hexanes. NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.2$ -6.66 (pr + NH, 5H), 5.69-5.6 (m, 1H), 4.8-4.69 (m, 1H), 3.76 (s, 3H), 3.56 (s, 1H), 3.5 (s, 1H), 2.69 (s, 3H), 1.4 (d, J=14 Hz, 1.5H), 1.35 (s, J=14 Hz, 1.5H).

 $C_{18}H_{19}N_3O_4F_2S$  (MW = 411); mass spectroscopy (MH<sup>+</sup>) 411.

#### Example 187

30

Synthesis of Methyl (3S,4S)-N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-4-amino-3-hydroxy-5-phenylpentanoate

10

15

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl (3S,4S)-4-amino-3-hydroxy-5-phenylpentanoate (Novabiochem), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.2 in 95:5 CHCl<sub>3</sub>/MeOH) and the product was purified by flash column chromatography using 95:5 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.29$  (d, 1H), 7.65 (d, 1H), 7.40-7.20 (m, 5H), 7.10 (m, 1H), 6.99 (m,2H), 5.27 (d, 1H), 4.47 (bs, 2H), 4.09 (m, 2H), 3.57 and 3.51 (m, 3H), 2.72 (m, 2H), 2.31 (m, 2H), 1.19 (m, 2H).

Optical Rotation:  $[\alpha]_{23} = -66^{\circ}$  (c 1, MeOH).  $C_{23}H_{26}N_2O_5F_2$  (MW = 448); mass spectroscopy (MH<sup>+</sup>) 449.

#### Example 188

## Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl](S)-2-aminohex-4-enoate

### Step A -- Synthesis of (S)-3-(Hex-4-enoyl)-4-(phenylmethyl)-2-oxazolidinone

To a mechanically stirred solution of 9.50 g (83.2 mmol, 1.10 equiv.) of 4hexenoic acid (commericially available from Lancaster, Catalog #252-427-6) 20 and 13.9 mL (10.1 g, 99.7 mmol, 1.33 equiv.) of triethylamine in 150 mL of dry THF, cooled to -78°C under dry N2, was added 10.71 mL (10.49 g, 87.0 mmol, 1.15 equiv.) of pivaloyl chloride (Aldrich). The mixture was warmed to 0°C for 60 min, and then recooled to -78°C. A solution of 13.4 g (75.6 mmol, 25 1.00 equiv) of (S)-(-)-(phenylmethyl)-2-oxazolidone (Aldrich) and 22 mg of triphenylmethane (indicator) in 150 mL of dry THF, stirred at -30°C to -45°C under  $N_2$ , was treated dropwise with n-butyllithium ( $\sim 2.5$  M in hexanes) (Aldrich) until an orange color persisted (~30 mL required). The resulting solution was cooled to -78°C and then added, via rapid cannulation, to the 30 above stirred mixture containing the mixed anhydride. The residual lithiated oxazolidone was rinsed in with two 10-mL portions of dry THF and the

10

20

25

30

resulting mixture was stirred at -78°C for 1.5 h, and then at 0°C for 1 h. The mixture was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and pH 7 phosphate buffer. The CH<sub>2</sub>Cl<sub>2</sub> phase was washed with saturated aqueous NaHCO<sub>3</sub> followed by half-saturated aqueous NaCl, dried (MgSO<sub>4</sub>), and evaporated *in vacuo*. The residual cream-colored solid (22.4 g) was chromatographed (Waters Prep 2000, 5.0 cm x 25 cm 10  $\mu$  Kromasil KR60-10SIL-5025 column) in two batches eluting with 85:15 hexanes/EtOAc. The chromatographed product was recrystallized from hexane to yield 14.34 g (first crop, 69%) of the title compound as fine white needles. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.37-7.20 (m, 5H, -C<sub>6</sub>H<sub>5</sub>), 5.60-5.43(m, 2H, CH=CHCH<sub>3</sub>), 4.71-4.63(m, 1H, NCH(Ph) CH<sub>2</sub>O), 4.23-4.14(m, 2H, NCH(Ph)CH<sub>2</sub>O), 3.295(dd, J=13.3, 3.3 Hz, 1H, CHHC<sub>6</sub>H<sub>5</sub>), 3.11-2.90 (m, 2H, CH<sub>2</sub>C=O), 2.758 (dd, J=13.3, 9.6 Hz, 1H, CHHC<sub>6</sub>H<sub>5</sub>), 2.42-2.34 (m, 2H, CH=CHCH<sub>2</sub>), 1.67-1.65 (m, 2H, CH=CHCH<sub>3</sub>).

### 15 Step B -- Synthesis of (4S)-3-[(S)-2-Azidohex-4-enoyl]-4- (phenylmethyl)-2-oxazolidinone

A solution of 5.47g (20.0 mmol, 1.00 equiv) of the product from Step A above in 60 mL of dry THF, stirred at -78°C under dry N2, was added via rapid cannulation to a stirred, cooled (-78°C) solution of 43.6 mL (22.0 mmol, 1.10 equiv) of potassium hexamethyldisilazide (0.505 M in toluene) (Aldrich) and 60 mL of dry THF. The residual imide solution was rinsed in with two 5mL portions of dry THF. The resulting solution was stirred at -78°C for 30 min. To the above solution of the K-enolate, stirred at -78°C under dry N<sub>2</sub>, was added a cooled (-78°C) solution of 7.43 g (24.0 mmol, 1.2 equiv) of trisyl azide (prepared as described in R. E. Harmon et al., J. Org. Chem., 1973, 38, 11-16) in 60 mL of dry THF via rapid cannulation. (Note the reaction exothermed to -68°C during the addition). After 1-2 min, 4.24 mL (4.45g, 74.1 mmol, 3.7 equiv) of glacial acetic acid was added in one portion. The resulting mixture was stirred at -78°C for 15 min, and was then allowed to warm to 25°C on stirring overnight. The mixture was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and pH 7 phosphate buffer. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3X) and the organic extracts were combined, washed with dilute

aqueous NaHCO<sub>3</sub>, dried (MgSO<sub>4</sub>), and evaporated *in vacuo*. The residual oil (9.55 g) was chromatographed (Waters Prep 2000, 5.0cm x 25 cm 10  $\mu$  Kromasil KR60-10SIL-5025 column) eluting with a 3 L linear gradient from 30:70 to 80:20 CH<sub>2</sub>Cl<sub>2</sub>/hexanes. After rechromatographing the mixed fractions (2X), a total of 5.27 g (84% yield) of the title compound (faster eluting, major diastereomer) was isolated as a colorless, viscous oil. <sup>1</sup>H NMR (300 MHz; CDCl<sub>3</sub>)  $\delta$  7.38-7.20 (m, 5H, -C<sub>6</sub>H<sub>5</sub>), 5.73-5.62 (m, 1H, CH=CHCH<sub>3</sub>), 5.52-5.41 (m, 1H, CH=CHCH<sub>3</sub>), 5.011 (dd, J=8.3, 5.5 Hz, 1H, CH(N<sub>3</sub>) C=0), 4.71-4.63 (m, 1H, NCH(Ph)CH<sub>2</sub>O), 4.236 (d, J=5.1 Hz, 2H, NCH(Ph)CH<sub>2</sub>O), 3.338 (dd, J= 13.4, 3.3 Hz, 1H, CHHC<sub>6</sub>H<sub>5</sub>), 2.827 (dd, J= 13.4, 9.5 Hz, 1H, CHHC<sub>6</sub>H<sub>5</sub>), 2.64-2.46 (m, 2H, CH<sub>2</sub>CH=CHCH<sub>3</sub>), 1.694 (dd, J=6.4, 1.1 Hz, 3H, CH=CHCH<sub>3</sub>).

### Step C -- Synthesis of (S)-2-Azidohex-4-enoic Acid

A solution of 5.00 g (15.91 mmol) of the product from Step B above in 240 mL of THF and 80 mL of deionized water, stirred at 0°C under N2, was 15 treated with 762 mg (31.8 mmol, 2.00 equiv) of LiOH (anhydrous powder). After stirring at 0°C for 30 min, 100 mL of 0.5 N aqueous NaHCO<sub>3</sub> was added and the THF was removed by rotary evaporation in vacuo. The residue was diluted to 400-500 mL with H<sub>2</sub>O and extracted with 5 portions of CH<sub>2</sub>Cl<sub>2</sub>. The 20 aqueous phase was acidified to pH 1-2 by the cautious addition of 5 N HCl, and then was extracted with 4 portions of EtOAc. The EtOAc extracts were combined, dried (Na2SO4), and evaporated in vacuo to yield 2.45 g (99%) of the title compound as a light amber oil.  $^{1}H$  NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  11.38 (br s, 1H, CO<sub>2</sub>H), 5.73-5.62 (m, 1H, CH<sub>3</sub>C $\underline{H}$ =CH), 5.48-5.38 (m, 1H,CH=CHCH<sub>2</sub>), 3.928 (dd, J= 7.8, 5.4 Hz, 1H, CH(N<sub>3</sub>)CO<sub>2</sub>H), 2.66-25 2.47(m, 2H, CH=CHC $\underline{H}_2$ ), 1.703 (dd, J=6.4, 1.1 Hz, 3H, CH<sub>3</sub>.

### Step D -- Methyl N-[N-tert-Butoxycarbonyl-L-alaninyl]-(S)-2-aminohex-4-enoate

A solution of 504.7 mg (3.25 mmol) of the product from Step C above in diethyl ether, cooled to 0°C, was treated dropwise with ethereal diazomethane

25

30

(prepared as described in L. F. Fieser et al., "Reagents for Organic Synthesis", Vol. 1, p. 191, Wiley & Sons (1967)) until a yellow color persisted. The excess diazomethane was removed by entraining with N2, and the ether was evaporated under a stream of N2. The residual oil was dissolved in 10 mL of anhydrous methanol. The solution was cooled to  $0^{\circ}\text{C}$  under dry  $N_2$  and 1.24~g(6.54 mmol, 2.0 equiv) of anhydrous SnCl<sub>2</sub> was added. The mixture was stirred at ≤25°C for 4 h, and the solvent was evaporated in vacuo to afford a solid tin-amine complex.

A solution of 1.23 g (6.50 mmol, 2.00 equiv.) of N-Boc-L-alanine (Sigma) and 0.715 mL (0.658 g, 6.50 mmol, 2.0 equiv.) of 4-methylmorpholine 10 (redistilled, 99.5%) (Aldrich) in 15 mL of anhydrous THF, cooled to -15 to -20°C under dry N<sub>2</sub>, was treated dropwise with 0.861 mL (0.907 g, 6.50 mmol, 2.00 equiv.) of iso-butyl chloroformate (Aldrich). After stirring at -15 to -20°C for 20 min, the resulting mixture containing the mixed anhydride was added via cannulation to the solid tin-amine complex (vide supra). The residual 15 mixed anhydride was rinsed in with 7 mL of THF and 1.1 g (13.1 mmol, 4.0 equiv.) of NaHCO3 powder and 5 mL of H2O was added. The mixture was stirred at 0°C for 5 h. An additional 1.1 g (13.1 mmol, 4.0 equiv.) of NaHCO<sub>3</sub> powder and 5 mL of H<sub>2</sub>O was added and the mixture was stirred at 20°C for 1.5 h. The mixture was filtered to remove the gelatinous precipitate, 20 and the filter cake was washed with several portions of EtOAc. The filtrate was washed with saturated aqueous NaHCO3 (the aqueous phase was backextracted with 3 portion of EtOAc), followed by pH 4-5 phosphate buffer (the aqueous phase was back-extracted with 3 portions of EtOAc). The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated in vacuo. The residual straw colored oil (1.21 g) was chromatographed (Waters Prep 2000, 5.0 cm x 25 cm 10  $\mu$ Kromasil KR60-10SIL-5025 column) eluting with a 3-L linear gradient from 80:20 to 40:60 hexane/EtOAc to yield 0.9088 g (89%) of the title compound as a white solid. Tlc R<sub>f</sub> 0.25 [silica, hexane/EtOAc 6:4)]; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  6.61 (d, J= 7.6 Hz, 1H, NH), 5.60-5.48(m, 1H, CH=CHCH<sub>3</sub>),

10

15

20

25

5.33-5.23 (m, 1H,  $C\underline{H}$ =CHCH<sub>3</sub>), 5.08 (d, J=7.3 Hz, 1H, NH), 4.591 (dt,  $J_d$  = 7.8 Hz,  $J_t$  = 5.7 Hz, 1H, HNC $\underline{H}$ (CH<sub>2</sub>CH=CHCH<sub>3</sub>)), 4.19 (br m, 1H, HNC $\underline{H}$ (CH<sub>3</sub>), 3.74(s, 3H, OCH<sub>3</sub>), 2.56-2.39 (sym m, 2H,  $\underline{C}\underline{H}_2$ CH=CHCH<sub>3</sub>), 1,658 (dd, J= 6.4, 1.2 Hz, 3H, CH<sub>2</sub>CH=CHC $\underline{H}_3$ ), 1.454 (s, 9H, OC(CH<sub>3</sub>)<sub>3</sub>), 1.358 (d, J=7.1 Hz, 3H, HNC $\underline{H}$ (CH<sub>3</sub>)).

### Step E -- Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohex-4-enoate

A solution of 0.811 g (2.58 mmol) of the product from Step D above in 5 mL of  $CH_2Cl_2$  was cooled to 0°C under dry  $N_2$  and 5 mL of trifluoroacetic acid was introduced by syringe at ≤4°C. The solution was stirred at 0°C for 40 min. Toluene (15 mL) was added and the mixture was evaporated in vacuo on the rotary evaporator. The addition of toluene and solvent evaporation was repeated. The residue was dissolved in 20 mL of CH2Cl2 and the solution was cooled to 0°C under dry N2. To this was added 1.35 mL (1.00 g, 7.74 mmol, 3.0 equiv) of ethyldiisopropylamine (Aldrich), followed by the dropwise addition at  $\leq$  6°C of 0.728 mL (0.938 g, 5.16 mmol, 2.0 equiv) of 3,5difluorophenylacetyl chloride (prepared from 3,5-difluorophenylacetic acid (Aldrich) using General Procedure H'). The resulting solution was stirred at 0°C for 2 h. Excess saturated aqueous NaHCO3 was added and the two phase mixture was stirred in an ice bath for 30 min. The mixture was diluted with  $CH_2Cl_2$  and washed successively with aqueous NaHCO<sub>3</sub>/Na<sub>2</sub>CO<sub>3</sub> (pH 10), 1 N aqueous NaHSO4, and saturated aqueous NaCl. The CH2Cl2 solution was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated in vacuo to afford 1.17 g of a yellow solid. This was recrystallized from EtOAc to yield 602 mg (63%) of the title compound as a fluffy white solid. This material was found by 300 mHz <sup>1</sup>H NMR analysis to consist of a 92:8 mixture of E and Z isomers, respectively.

NMR data was as follows:

<sup>1</sup>H-nmr (300 MHz, CDCl<sub>3</sub>):  $\delta = 6.85\text{-}6.69$  (m, 3H), 6.335 (br d, J=7.8 Hz, 1H), 6.289 (br d, J=7.0 Hz, 1H), 5.58-5.47 (m, 1H), 5.28-5.18 (m, 1H), 4.58-4.45 (m, 2H) 3.745 (s, 3H), 3.528 (s, 2H), 2.457 (apparent t, J=6.4 Hz,

10

15

20

2H), 1.650 (dd, J=6.5, 1.3Hz, 3H), 1.58 (dm, J=6.5 Hz, 0.08H), 1.375 (d, J=7.0 Hz, 3H).

IR (CHCl<sub>3</sub>) 3421, 1742, 1667, 1626, 1597, 1503, and 1120 cm<sup>-1</sup> Anal. Calcd for  $C_{18}H_{22}F_2N_2O_4$ : C, 58.69; H, 6.02; N, 7.60. Found: C, 58.83, H, 5.89; N, 7.67.

 $C_{18}H_{22}F_2N_2O_5$  (MW = 368); mass spectroscopy (MH<sup>+</sup>) 368.

#### Example 189

## Synthesis of N-[N-(Cyclopropylacetyl)-L-alaninyl]L-phenylglycine tert-Butyl Ester

Following General Procedure U and using cyclopropylacetic acid (Lancaster) and N-(L-alaninyl)-L-phenylglycine tert-butyl ester (General Procedure U of Z-alanine (Bachem) to phenylglycine-t-butyl (Novabio) and then General Procedure O), the title compound was prepared as a solid (mp = 105-107°C). The reaction was monitored by tlc (Rf = 0.33 in 1:1 EtOAc/hexane, 0.13 in 5% MeOH/DCM).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.15$  (m, 2H), 0.56 (m, 2H), 0.91 (m, 1H), 1.38 (m, 12H), 2.09 (d, J = 7.1 Hz, 2H), 4.62 (m, 1H), 5.39 (d, J = 7.2 Hz, 1H), 6.52 (d, J = 7.2 Hz, 1H), 7.31 (m, 6H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 4.53$ , 4.55, 6.9, 18.4, 27.8, 41.2, 48.4, 57.1, 82.6, 127.0, 128.2, 128.7, 136.8, 169.4, 171.4, 172.3.

 $C_{20}H_{28}N_2O_4$  (MW = 360.46); mass spectroscopy (MH<sup>+</sup>) 361.

### Example 190

25

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl](S)-2-amino-2-(4-phenylphenyl)acetamide

Following General Procedure AB and using 3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and 4-biphenylcarboxaldehyde (Aldrich), the

title compound was prepared as a solid (mp = 266-267°C). The product was purified by recrystallization from EtOAc and/or EtOAc/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.42$  (d, 1H, J=7 Hz), 8.31 (d, 1H, J=7 Hz), 7.91 (s, 1H), 7.6-7.56 (m, 4H), 7.42-7.59 (m, 5H), 7.2-7.69 (m, 3H), 5.42 (d, 1H, J=8 Hz), 4.42 (pentet, 1H, J=8 Hz), 3.5 (s, 1H), 1.2 (doublet on top of a singlet, 12H).

 $C_{29}H_{31}N_3O_3F_2$  (MW = 508); mass spectroscopy (MH<sup>+</sup>) 508.4.

### Example 191

10

5

## Synthesis of N-[N-(3,5-Diffuorophenylacetyl)-(S)-2-aminobutanoyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure D and using 3,5-difluorophenylacetic acid (Aldrich) and N-[(S)-2-aminobutanoyl]-L-phenylglycine tert-butyl ester (from Example D13 above), the title compound was prepared as a solid (mp = 138.7-140.0°C). The reaction was monitored by tlc (Rf = 0.24 in 2/1 hexanes:EtOAc) and the product was purified by flash chromromatography and HPLC.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ , 250 MHz): δ = 8.66 (d, J=6.75 Hz, 1H), 8.30 (d, J=8.26 Hz, 1H), 7.37 (bs, 5H), 7.11-6.96 (m, 3H), 5.23 (d, J=7.00 Hz, 1H), 4.36 (td, J=7.88, 5.50 Hz, 1H), 3.53 (AB<sub>q</sub>, J<sub>AB</sub>= 14.05 Hz, ΔV<sub>AB</sub>=7.75 Hz, 2H), 1.85-1.48 (m, 2H), 1.34 (s, 9H), 0.88 (t, J=7.38 Hz, 3H).

Optical Rotation:  $[\alpha]_{20} = 21.8^{\circ}$  (c 1.0, MeOH).

25  $C_{24}H_{28}N_2O_4F_2$  (MW = 446.50); mass spectroscopy (MH<sup>+</sup>, minus CO<sub>2</sub>-tBu) 345.2.

#### Example 192

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-valinyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure D and using 3,5-difluorophenylacetic acid (Aldrich) and N-(L-valinyl)-L-phenylglycine tert-butyl ester (from Example D14 above), the title compound was prepared as a solid (mp =  $170.5-171.8^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.39 in 2:1 hexanes/EtOAc) and the product was purified by flash chromromatography and HPLC.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ , 250 MHz):  $\delta = 8.71$  (d, J=6.75 Hz, 1H), 8.22 (d, J=9.26 Hz, 1H), 7.37 (bs, 5H), 7.11-6.96 (m, 3H), 5.23 (d, J=6.50 Hz, 1H), 4.36 (dd, J=8.88, 6.38 Hz, 1H), 3.55 (AB<sub>q</sub>, J<sub>AB</sub>= 13.88 Hz,  $\Delta V_{AB}$ = 21.56 Hz, 2H), 2.10-1.95 (m, 1H), 1.34 (s, 9H), 0.88 (d, J=6.75 Hz, 3H), 0.86 (d, J=6.50 Hz, 3H).

Optical Rotation:  $[\alpha]_{20} = 20.8^{\circ}$  (c 1.0, MeOH).

 $C_{25}H_{30}N_2O_4F_2$  (MW = 460.53); mass spectroscopy (MH<sup>+</sup>, minus  $CO_2$ -tBu) 359.2.

15

5

10

### Example 193

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-methioninyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid

(Aldrich) and N-(L-methioninyl)-L-phenylglycine methyl ester hydrochloride
(prepared from N-BOC-L-methionine (Sigma) and L-phenylglycine methyl ester
hydrochloride (Aldrich) using General Procedure E, followed by removal of the
BOC-group using General Procedure P), the title compound was prepared as a
solid (mp = 189.3°C). The reaction was monitored by tlc (Rf = 0.53 in 5:95

MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by recrystallization from ethyl
acetate/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.85$  (d, J=6.7 Hz, 1H), 8.41 (d, J=8.1 Hz, 1H), 7.38 (m, 5H), 7.09 (m, 1H), 6.98 (m, 2H), 5.38 (d, J=6.6 Hz, 1H), 4.47 (m, J=8.2 Hz, 1H), 3.62 (s, 3H), 3.51 (d, 2H), 2.46 (t, 2H), 2.04 (s, 3H), 1.89 (m, 2H).

30

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta$  = 172.036, 171.729, 169.883, 164.658, 164.479, 161.406, 161.227, 141.689, 141.557, 141.427, 136.524, 129.512, 129.213, 128.717, 126.274, 113.187, 113.085, 112.961, 112.862, 103.023, 102.684, 102.340, 93.065, 57.205, 53.063, 42.231, 33.075, 30.221, 15.465.  $C_{22}H_{24}N_2O_4F_2S$  (MW = 450.51); mass spectroscopy (MH+) 450.

#### Example 194

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-valinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid

(Aldrich) and N-(L-valinyl)-L-phenylglycine methyl ester hydrochloride

(prepared from N-BOC-L-valine (Sigma) and L-phenylglycine methyl ester

hydrochloride (Aldrich) using General Procedure E, followed by removal of the

BOC-group using General Procedure P), the title compound was prepared as a

solid (mp = 226.5°C). The reaction was monitored by tlc (Rf = 0.49 in 5:95

MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by flash chromotography using

MeOH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ): δ = 8.84 (d, J=6.2 Hz, 1H), 8.23 (d, J=8.8 Hz, 1H), 7.38 (m, 5H), 7.07 (m, 1H), 6.98 (m, 2H), 5.37 (d, J=6.5 Hz, 1H), 4.34 (m, J=8.9 Hz, 1H), 3.55 (m, 5H), 2.01 (m, 1H), 0.87 (m, 6H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ): δ = 171.988, 171.690, 169.861, 164.633, 164.456, 161.382, 161.204, 141.987, 141.859, 141.727, 136.553, 129.470, 129.192, 128.791, 113.128, 113.026, 112.902, 112.803, 102.961, 102.619, 102.281, 57.914, 57.262, 52.935, 42.274, 31.728, 19.845, 18.815.  $C_{22}H_{24}N_2O_4F_2$  (MW = 418.44); mass spectroscopy (MH<sup>+</sup>) 418.1.

#### Example 195

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycine Methyl Ester

20

25

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and N-(2-aminobutanoyl)-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-aminobutyric acid (Sigma) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 215.3°C). The reaction was monitored by tlc (Rf = 0.46 in 5:95 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by recrystallization from ethyl acetate/hexanes.

NMR data was as follows:

10 
<sup>1</sup>H-nmr (DMSO- $d_6$ ): δ = 8.83 (d, J=6.8 Hz, 1H), 8.32 (d, J=8.1 Hz, 1H), 7.38 (m, 5H), 7.08 (m, 1H), 6.98 (m, 2H), 5.38 (d, J=6.8 Hz, 1H), 4.35 (m, J=7.9 Hz, 1H), 3.61 (s, 3H), 3.52 (d, 2H), 1.64 (m, 2H), 0.88 (t, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ): δ = 171.684, 170.934, 168.984, 164.193, 163.980, 160.295, 160.083, 141.059, 140.902, 140.743, 135.857, 128.689, 128.372, 127.892, 112.387, 112.257, 112.131, 111.999, 102.254, 101.845, 101.438, 56.351, 53.441, 52.212, 41.436, 25.675, 10.067.  $C_{21}H_{22}N_2O_4F_2$  (MW = 404.42); mass spectroscopy (MH+) 405.1.

### Example 196

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-leucinyl]L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and N-(L-leucinyl)-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-leucine (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 178.4°C). The reaction was monitored by tlc (Rf = 0.51 in 5:95 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by flash chromotograph using MeOH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

WO 98/22494 PCT/US97/20804

-- 297 --

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.85$  (d, J=6.8 Hz, 1H), 8.33 (d, J=8.3 Hz, 1H), 7.37 (m, 5H), 7.08 (m, 1H), 6.95 (m, 2H), 5.37 (d, J=6.8 Hz, 1H), 4.46(m, J=8.3 Hz, 1H), 3.60 (s, 3H), 3.49 (d, 2H), 1.55 (m, 3H), 0.89 (d, 3H), 0.82 (d, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 172.225$ , 170.899, 168.888, 164.197, 163.984, 160.298, 160.086, 141.029, 140.887, 140.723, 135.875, 128.657, 128.348, 127.944, 112.340, 112.207, 112.084, 111.951, 102.251, 101.842, 101.435, 56.343, 52.214, 50.697, 41.510, 40.982, 24.449, 23.056, 21.575.

 $C_{23}H_{26}N_2O_4F_2$  (MW = 432.47); mass spectroscopy (MH<sup>+</sup>) 432.1.

10

15

20

25

30

5

### Example 197

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-phenylalaninyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and N-(L-phenylalaninyl)-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-phenylalanine (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 188.3°C). The reaction was monitored by tlc (Rf = 0.59 in 5:95 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by flash chromotography using MeOH/CH2Cl2 as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.99$  (d, J=6.9 Hz, 1H), 8.44 (d, J=8.6 Hz, 1H), 7.4 (m, 5H), 7.21 (m, 5H), 7.03 (m, 1H), 6.77 (m, 2H), 5.42 (d, J=6.9Hz, 1H), 4.70 (m, J=8.5 Hz, 1H), 3.63 (s, 3H), 3.40 (m, 2H), 3.08 (m, 1H), 2.76 (m, 1H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 171.428$ , 170.896, 168.853, 164.127, 163.915, 160.222, 160.010, 140.756, 140.601, 140.438, 137.662, 135.918, 130.638, 129.247, 128.737, 128.415, 127.908, 126.281, 112.147, 112.025, 111.892, 102.189, 101.782, 101.373, 56.411, 53.461, 52.306, 41.513, 37.796.  $C_{26}H_{24}N_2O_4F_2$  (MW = 466.49); mass spectroscopy (MH<sup>+</sup>) 466.

20

25

30

### Example 198

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)glycinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid 5 (Aldrich) and N-(glycinyl)-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-glycine (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 142.3 °C). The reaction was monitored by tlc (Rf = 0.33 in 5:95MeOH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by recrystallization from diethyl ether/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.82$  (d, J=7.2 Hz, 1H), 8.39 (t, 1H), 7.37 (m, 5H), 7.05 (m, 3H), 5.44 (d, 7.1 Hz, 1H), 3.83 (d, 2H), 3.62 (s, 3H), 3.53 (s, 15 2H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 170.956$ , 169.427, 168.788, 164.226, 164.013, 160.329, 160.115, 140.817, 140.663, 140.499, 136.222, 128.728, 128.338, 127.687, 112.494, 112.360, 112.238, 112.104, 102.310, 101.900, 101.492, 56.200, 52.321, 41.731, 41.464.

 $C_{19}H_{18}N_2O_4F_2$  (MW = 376.36); mass spectroscopy (MH<sup>+</sup>) 376.0.

### Example 199

### Synthesis of $N\hbox{-}[N\hbox{-}(3,5\hbox{-}{\bf Difluor ophenylacetyl})\hbox{-}{\bf L}\hbox{-}{\bf phenylglycinyl}]\hbox{-}$ L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and N-(L-phenylglycinyl)-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-phenylglycine (Novabiochem) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure AH, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 222.8°C). The reaction was monitored by tlc (Rf = 0.61 in 5:95 MeOH/CH $_2$ Cl $_2$ ) and the product was purified by recrystallization from ethyl acetate.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 9.22$  (d, J=6.8 Hz, 1H), 8.85 (d, J=8.2 Hz, 1H), 7.37 (m, 10H), 7.08 (m, 1H), 6.97 (d, 2H), 5.69 (d, J=8.2 Hz, 1H), 5.43 (d, J=6.8 Hz, 1H), 3.61 (d, 2H), 3.55 (s, 3H).

<sup>13</sup>C-nmr (DMSO- $d_6$ ):  $\delta = 170.606$ , 169.727, 168.777, 164.194, 163.982, 160.296, 160.082, 140.920, 140.757, 140.603, 138.391, 135.900, 128.732, 128.425, 128.233, 127.871, 127.556, 127.222, 112.467, 112.340, 112.209,

112.082, 102.292, 101.884, 101.475, 56.431, 55.621, 52.203, 41.205.  $C_{25}H_{22}N_2O_4F_2$  (MW = 452.46); mass spectroscopy (MH<sup>+</sup>) 452.2.

### Example 200

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-alanine Methyl Ester

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-alanine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 140.5-142°C). The reaction was monitored by tlc (Rf = 0.17 in 50% EtOAc/hexanes).

NMR data was as follows:

20  ${}^{1}\text{H-nmr}$  (CDCl<sub>3</sub>):  $\delta = 1.3\text{-}1.4$  (m, 6H), 3.55 (s, 2H), 3.75 (s, 3H), 4.4-4.6 (m, 2H), 6.1-6.3 (brd, 1H), 6.6-6.7 (brd, 1H), 7.2-7.4 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.4$ , 19.0, 44.1, 48.6, 49.3, 53.0, 127.9, 129.5, 129.8, 135.1, 171.5, 172.4, 173.6.

 $C_{15}H_{20}N_2O_4$  (MW = 292.34); mass spectroscopy (MH<sup>+</sup>) 293.

25

10

### Example 201

### $Synthesis \ of \\ N-[N-(Phenylacetyl)-L-alaninyl]-L-leucine \ Methyl \ Ester$

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-leucine methyl ester hydrochloride (Aldrich), the title

compound was prepared as a solid (mp = 102.5-105°C). The reaction was monitored by tic (Rf = 0.25 in 50% EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.8$ -0.95 (m, 6H), 1.3 (d, J=7 Hz, 3H), 1.4-1.6 (m, 3H), 3.58 (s, 2H), 3.75 (s, 3H), 4.4-4.6 (m, 2H), 6.2 (brd, 1H), 6.7 (brd, 5 1H), 7.2-7.4 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.7, 22.3, 23.4, 25.3, 41.5, 44.1, 49.2, 51.4,$ 52.8, 127.9, 129.5, 129.8, 135.0, 171.5, 172.6, 173.7.

 $C_{18}H_{26}N_2O_4$  (MW = 334.42); mass spectroscopy (MH<sup>+</sup>) 335.

10

15

### Example 202

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-isoleucine Methyl Ester

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-isoleucine methyl ester hydrochloride (Sigma), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.24 in 50% EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.8$ -0.95 (m, 6H), 1.0-1.2 (m, 1H), 1.2-1.4 (m including 1.3 (d, J=7 Hz, 4H)), 1.8-1.9 (m, 1H), 3.58 (s, 2H), 3.75 (s, 3H),

4.4-4.6 (m, 2H), 6.2 (brd, 1H), 6.7 (brd, 1H), 7.2-7.4 (m, 5H). 20

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 12.1, 16.0, 18.5, 25.6, 38.1, 44.1, 49.3, 52.7,$ 57.2, 127.9, 129.6, 129.8, 135.0, 171.5, 172.5, 172.6.

 $C_{18}H_{26}N_2O_4$  (MW = 334.42); mass spectroscopy (MH<sup>+</sup>) 335.

### Example 203

25

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-proline Methyl Ester

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-proline methyl ester hydrochloride (Bachem), the

title compound was prepared as an oil. The reaction was monitored by tlc (Rf = 0.12 in 50% EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 1.33$  (d, J=7 Hz, 3H), 1.9-2.1 (m, 3H), 2.1-2.25 (m, 1H), 3.5-3.8 (m including 3.58 (s) and 3.75 (s), total 7H), 4-4.4 (m, 1H), 4.7-4.8 (m, 1H), 6.5 (brd, 1H), 7.2-7.4 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.5, 25.5, 29.5, 44.1, 47.3, 47.4, 52.8, 59.3, 127.8, 129.42, 129.48, 129.9, 135.2, 170.9, 171.8, 172.8.$ 

 $C_{17}H_{22}N_2O_4$  (MW = 318.38); mass spectroscopy (MH<sup>+</sup>) 319.

10

15

5

### Example 204

### $Synthesis \ of \\ N-[N-(Phenylacetyl)-L-alaninyl]-L-phenylalanine \ Methyl \ Ester$

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-phenylalanine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 148-149.5°C). The reaction was monitored by tlc (Rf = 0.24 in 50% EtOAc/hexanes) and the product was not purified.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 1.25$  (d, J=7 Hz, 3H), 3.02 (dd, J=7, 14Hz, 1H), 3.12 (dd, J=5, 14 Hz, 1H), 3.53 (s, 2H), 3.72 (s, 3H), 4.4-4.5 (m, 1H), 4.75-4.85 (m, 1H), 5.9 (brd, 1H), 6.5 (brd, 1H), 7.0-7.5 (m, 10H). <sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.6$ , 38.3, 44.0, 49.2, 52.9, 53.9, 127.7, 128.0, 129.1, 129.6, 129.8, 129.9, 135.0, 136.3, 171.4, 172.2, 172.3. C<sub>21</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> (MW = 368.44); mass spectroscopy (MH<sup>+</sup>) 369.

25

#### Example 205

# Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]N,-(tert-butoxycarbonyl)-L-lysine Methyl Ester

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and N<sub>4</sub>-(tert-butoxycarbonyl)-L-lysine methyl ester

hydrochloride (Bachem), the title compound was prepared as a solid (mp =  $119-121^{\circ}$ C). The reaction was monitored by tlc (Rf = 0.46 in 90:10:1 CH<sub>2</sub>Cl<sub>2</sub>\MeOH\NH<sub>4</sub>OH).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 1.2-1.9 (m, 18H)(includes 1.3 (d, J=7 Hz) and 1.4 (s)), 3.0-3.15 (m, 2H), 3.12 (dd, J=5, 14 Hz, 1H), 3.57 (s, 2H), 3.72 (s, 3H), 4.4-4.5 (m, 2H), 4.75-4.85 (m, 1H), 6.2 (brd, 1H), 6.75 (brd, 1H), 7.2-7.4 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.6$ , 22.9, 29.0, 29.9, 32.0, 40.5, 44.0, 49.4, 10 52.7, 53.0, 79.8, 127.9, 129.5, 129.8, 135.1, 156.7, 171.6, 172.7, 173.0.  $C_{23}H_{35}N_3O_6$  (MW = 449.55); mass spectroscopy (MH<sup>+</sup>) = 450.

#### Example 206

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]glycine Methyl Ester

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and glycine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 152-153.5°C). The reaction was monitored by tlc (Rf = 0.10 in 50% EtOAc/hexanes).

NMR data was as follows:

20  $^{1}$ H-nmr (CDCl<sub>3</sub>):  $\delta = 1.33$  (d, J=7 Hz, 3H), 3.59 (s, 2H), 3.75 (s, 3H), 3.97 (d, J=6.5 Hz, 2H), 4.5-4.6 (m, 1H), 6.1 (brd, 1H), 6.8 (brs, 1H), 7.2-7.6 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 18.7$ , 41.6, 43.9, 49.2, 52.9, 127.9, 129.5, 129.9, 135.0, 170.6, 171.7, 173.2.

25  $C_{14}H_{18}N_2O_4$  (MW = 278.31); mass spectroscopy (MH<sup>+</sup>) 279.

#### Example 207

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-valine Methyl Ester

15

20

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-valine methyl ester hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 112-115°C). The reaction was monitored by tlc (Rf = 0.33 in 50% EtOAc/hexanes) and the product was not purified.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.8$ -0.9 (overlapping d appearing as t, J=6 Hz, 6H), 2.0-2.2 (m, 1H), 3.57 (s, 2H), 3.72 (s, 3H), 4.4-4.5 (m, 1H), 4.5-4.65 (m, 1H), 6.2 (brd, 1H), 6.75 (brd, 1H), 7.2-7.4 (m, 5H).

10  ${}^{13}\text{C-nmr}$  (CDCl<sub>3</sub>):  $\delta = 18.3$ , 18.5, 19.5, 31.5, 44.1, 49.3, 52.7, 57.9, 127.9, 129.5, 129.8, 135.0, 171.5, 172.7, 172.7.

 $C_{17}H_{24}N_2O_4$  (MW = 320.39); mass spectroscopy (MH<sup>+</sup>) 321.

### Example 208

### Synthesis of

Methyl N-[N-(Phenylacetyl)-L-alaninyl]-2-(S)-aminobutanoate

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and methyl L-2-aminobutanoate hydrochloride (prepared from L-2-aminobutanoic acid (Bachem) using General Procedure H (without extractions)), the title compound was prepared as a solid (mp = 120°C). The reaction was monitored by tlc (Rf = 0.2 in 50% EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.85$  (t, J=6 Hz, 3H), 1.32 (d, J=7 Hz, 3H), 1.6-1.9 (m, 2H), 3.57 (s, 2H), 3.72 (s, 3H), 4.4-4.6 (m, 2H), 6.2 (brd, 1H), 6.75 (brd, 1H), 7.2-7.4 (m, 5H).

25  $^{13}$ C-nmr (CDCl<sub>3</sub>):  $\delta = 10.2, 18.9, 25.8, 44.0, 49.3, 52.8, 54.0, 127.9, 129.5, 129.8, 135.1, 171.5, 172.7, 173.0.$ 

 $C_{16}H_{22}N_2O_4$  (MW = 306.36); mass spectroscopy (MH<sup>+</sup>) 307.

#### Example 209

### $Synthesis of \\ Methyl N-[N-(Phenylacetyl)-L-alaninyl]-2-(S)-aminopentanoate$

15

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and methyl 2-(S)-aminopentanoate hydrochloride (prepared from L-2-aminovaleric acid (Bachem) using General Procedure H (without extractions)), the title compound was prepared as a solid (mp = 135-137°C).

The reaction was monitored by tlc (Rf = 0.30 in 50% EtOAc/hexanes).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 0.87$  (t, J=6 Hz, 3H), 1.2-1.4 (m with d, J=7 Hz, 5H), 1.5-1.8 (m, 2H), 3.57 (s, 2H), 3.72 (s, 3H), 4.4-4.5 (m, 2H), 6.4 (brd, 1H), 7.0 (brd, 1H), 7.2-7.4 (m, 5H).

10  ${}^{13}\text{C-nmr}$  (CDCl<sub>3</sub>):  $\delta = 14.2$ , 19.0, 19.2, 34.5, 44.0, 49.2, 52.7, 52.8, 127.8, 129.4, 129.8, 135.2, 171.5, 172.8, 173.3.

 $C_{17}H_{24}N_2O_4$  (MW = 320.39); mass spectroscopy (MH<sup>+</sup>) 321.

### Example 210

### $Synthesis of \\ N-[N-(3-Nitrophenylacetyl)-L-alaninyl]-L-valine$

Following General Procedure AF and using N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine ethyl ester (from Example 143 above), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.05 in 9:1 CHCl<sub>3</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.41$  (d, 1H), 8.13 (s, 1H), 8.09 (d, 1H), 7.91 (d, 1H), 7.68 (d, 1H), 7.56 (t, 1H), 4.4 (m, 1H), 4.10 (m, 1H), 3.63 (s, 2H), 2.01 (m, 1H), 1.19 (m, 3H), 0.89 (d, 6H).

Optical Rotation:  $[\alpha]_{23} = -49^{\circ}$  (c 1, MeOH).

25  $C_{16}H_{21}N_3O_6$  (MW = 351.3); mass spectroscopy (MH<sup>+</sup>) 352.

### Example 211

### Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-N-methylalanine Methyl Ester

Following General Procedure A and using N-(phenylacetyl)-L-alanine (from Example B1 above) and L-N-methylalanine methyl ester hydrochloride (prepared from L-N-methylalanine hydrochloride (Bachem) using General Procedure H (without extractions)), the title compound was prepared as an oil. The reaction was monitored by tlc (Rf = 0.13 in 50% EtOAc/hexanes) and the product was purified by column chromatography using 60% EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 1.2$ -1.6 (m including 1.32 (d, J=7 Hz, 6H), 2.97 (s (rotomers), 3H), 3.57 (s, 2H), 3.7-3.8 (s (rotomers), 3H), 4.4-5.2 (m, 2H), 6.6 (brd, 1H), 7.2-7.4 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 14.7, 15.0, 18.8, 19.1, 31.6, 32.3, 44.3, 46.2, 46.3, 52.7, 52.88, 52.93, 53.6, 127.81, 127.85, 129.45, 129.48, 129.9, 135.2, 170.60, 170.67, 172.19, 172.4, 173.25. 173.31.

15  $C_{16}H_{22}N_2O_4$  (MW = 306.36); mass spectroscopy (MH<sup>+</sup>) 307.

#### Example 212

### $Synthesis \ of \\ N-[N-(Isovaleryl)-L-phenylglycinyl]-L-alanine Isobutyl \ Ester$

Following General Procedure C and using N-(isovaleryl)-L-phenylglycine

(prepared from isovaleric acid (Aldrich) and L-phenylglycine methyl ester
hydrochloride (Aldrich) using General Procedure C, followed by hydrolysis
using General Procedure AF) and L-alanine isobutyl ester hydrochloride
(prepared from N-BOC-L-alanine (Sigma) and 2-methyl-1-propanol (Aldrich)
using General Procedure C (with catalystic DMAP), followed by removal of the
BOC-group using General Procedure P), the title compound was prepared as a
solid (mp = 181-186°C). The reaction was monitored by tlc (Rf = 0.4 in 1:1
EtOAc/hexanes) and the product was purified by silica gel chromatography
using 1:1 EtOAc/hexanes as the eluent.

NMR data was as follows:

30  ${}^{1}\text{H-nmr}$  (CDCl<sub>3</sub>):  $\delta = 1.31$  (d, 3H), 5.59 (d, 1H).

10

15

Optical Rotation:  $[\alpha]_{20} = +19^{\circ} @ 589 \text{ nm} \text{ (c 1.03, DMSO)}.$  $C_{20}H_{30}N_2O_4 \text{ (MW = 362); mass spectroscopy (MH<sup>+</sup>) 363.}$ 

### Example 213

### Synthesis of N-[N-(Isovaleryl)-L-isoleucinyl]-L-alanine Isobutyl Ester

Following General Procedure C and using N-(isovaleryl)-L-isoleucine (prepared from isovaleric acid (Aldrich) and L-isoleucine methyl ester hydrochloride (Aldrich) using General Procedure C, followed by hydrolysis using General Procedure AF) and L-alanine isobutyl ester hydrochloride (prepared from N-BOC-L-alanine (Sigma) and 2-methyl-1-propanol (Aldrich) using General Procedure C (with catalystic DMAP), followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 142-146°C). The reaction was monitored by tlc (Rf = 0.4 in 1:1 EtOAc/hexanes) and the product was purified by silica gel chromatography using 1:1 EtOAc/hexanes as the eluent to provide a 1:4 mixture of diastereomers.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.26$  (d, 3H), 7.70 (d, 1H), 7.80 (d, 1H), 8.30 (d, 1H), 8.40 (d, 1H).

20  $C_{18}H_{34}N_2O_4$  (MW = 342.48); mass spectroscopy (MH<sup>+</sup>) 343.

### Example 214

## Synthesis of N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

25 Following General Procedure C and using N-(3,5-difluorophenylacetyl)-Lalanine (from Example B2 above) and N-cyclohexyl-L-phenylglycinamide
(prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and
cyclohexylamine (Aldrich) using General Procedure M, followed by removal of
the BOC-group using General Procedure P), the title compound was prepared as

a solid. The reaction was monitored by tlc (Rf = 0.5 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by trituration from ethanol.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.55$  (m, 2H), 8.08 (d, 1H), 7.30 (m, 5H), 7.08 (m, 1H), 6.97 (d, 2H), 5.37 (m, 1H), 3.47 (s, 2H), 1.8-1.6 (m, 6H), 1.23-0.98 (m, 7H).

Optical Rotation:  $[\alpha]_{23} = -32.7^{\circ}$  (c 1, MeOH).  $C_{25}H_{29}F_2N_3O_3$  (MW = 457); mass spectroscopy (MH<sup>+</sup>) 458.

### Example 215

10

15

5

# Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4-hydroxyproline Ethyl Ester

Following General Procedure F and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-4-hydroxyproline ethyl ester hydrochloride (Pfaltz & Bauer), the title compound was prepared as a foam. The reaction was monitored by tlc (Rf = 0.32 in 95:5  $CH_2Cl_2/MeOH$ ) and the product was purified by flash column chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>, 250 Mz):  $\delta = 7.31$  (d, 1H, J= 7.00 Hz), 6.83-6.64 (m, 3H), 4.67 (p, 1H, J=7.09 Hz), 4.58 (t, 1H, J= 8.26 Hz), 4.47 (bs, 1H), 4.25-4.06 (m, 2H), 3.81 (bd, 1H, J=11.01 Hz), 3.62 (dd, 1H, J=10.76, 3.75 Hz), 3.46 (s, 2H), 2.30 (dd, 1H, J= 13.51, 8.26 Hz), 1.96 (ddd, 1H, J= 13.44, 8.82, 4.56 Hz), 1.33 (d, 3H, J= 6.75 Hz), 1.24 (t, 3H, J= 7.13 Hz).  $C_{18}H_{23}F_2N_2O_5$  (MW = 384.38); mass spectroscopy (MH<sup>+</sup>) 385.1.

25

30

### Example 216

### Synthesis of

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-lysine Methyl Ester

Following General Procedure Y and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]- $N_c$ -(tert-butoxycarbonyl)-L-lysine methyl ester (from Example 43 above), the title compound was prepared as an oil. The title compound was

isolated as the trifluoroacetic acid salt (containing about 5% excess trifluoroacetic acid).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub> + 2 drops of CD<sub>3</sub>OD):  $\delta = 6.40$ -6.52 (m, 3H), 4.17 (t, 1H), 4.40 (q, 1H), 3.42 (s, 3H), 3.23 (s, 3H), 2.53 (bs, 2H), 1.60 (m, 1H), 1.32 (m, 3H), 1.02-1.13 (m, 2H), 1.10 (d, 2H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub> + 2 drops of CD<sub>3</sub>OD):  $\delta$  = 174.1, 166.4, 166.2, 163.1, 163.0, 141.3, 113.8, 113.7, 113.5, 103.5, 55.2, 56.3, 43.0, 40.9, 32.2, 28.1, 24.0, 18.2.

10  $C_{21}H_{26}F_5N_3O_6$  (MW = 511.4); mass spectroscopy (MH<sup>+</sup>) 512.

### Example 217

### Synthesis of N'-[N-(3,5-Diffuorophenylacetyl)-L-alaninyl]-L-glutamide

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-glutamide hydrochloride (Bachem), the title compound was prepared as a solid (mp = 260-263°C). The reaction was monitored by tlc (Rf = 0.77 in 10% MeOH/DCM) and the product was purified by silica gel chromatography.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.40$  (d, J=7.1 Hz, 1H), 8.02 (d, J=6.9, 1H), 7.2 (m, 2H), 7.0 (m, 4H), 6.76 (s, 1H), 4.2 (m, 1H), 3.56 (s, 2H), 2.1 (m, 2H), 1.9 (m, 2H), 1.21 (d, J=7.0 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.5$ , 172.4, 169.5, 112.5, 110.4, 102.3, 52.5, 49.0, 41.6, 35.7, 31.8, 28.1, 18.4.

25  $C_{16}H_{20}F_2N_4O_4$  (MW = 370); mass spectroscopy (MH<sup>+</sup>) 371.

### Example 218

## Synthesis of Methyl 1-[N-(3,5-Difluorophenylacetyl)L-alaninyl]piperidine-2-carboxylate

Following General Procedure A and using N-(3,5-difluorophenylacetyl)-

5

15

L-alanine (from Example B2 above) and methyl pipecolinate hydrochloride (Aldrich), the title compound was prepared as a solid (mp = 114-118°C). The reaction was monitored by tlc (Rf = 0.71 in 10% MeOH/DCM) and the product was purified by acid/base washes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.95$  (dd, J=7.1, 7.1, 7.1 Hz; 1H), 6.81 (d, J=6.1 Hz, 2H), 6.7 (m, 1H), 5.28 (dd, J=5.0 Hz, 12.6, 5.4, 1H), 4.93 (q, J=7.0, 6.9, 7.0 Hz, 1H), 3.75 (s, 1H), 3.70 (s, 3H), 3.50 (s, 2H), 3.2 (m, 1H), 2.27 (d, J=3.5 Hz, 1H), 1.5 (m, 5H), 1.31 (d, J=5.2 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.8$ ; 172.6; 171.7; 171.6; 169.2; 169.1; 112.9; 112.8; 112.7; 112.6; 103.2; 102.8; 53.0; 52.9; 52.9; 52.7; 46.2; 46.1; 43.9; 43.9; 27.1; 26.8; 25.6; 21.4; 19.9; 18.5.

 $C_{18}H_{22}F_2N_2O_4$  (MW = 368); mass spectroscopy (MH<sup>+</sup>) 369.

### Example 219

15

20

5

10

# Synthesis of N-[(S)-3-Hydroxy-6-methylhept-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure AA and using N-[(S)-6-methyl-3-oxohept-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (from Example 168 above), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.75 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 99:1 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.81$  (m, 1H), 6.72 (m, 2H), 6.39 (m, 2H), 4.45 (m, 1H), 3.97 (m, 1H), 3.60 (m, 1H), 3.52 (s, 2H), 1.54 (m, 1H), 1.4 (m, 5H), 1.09 (m, 3H), 0.9 (m, 6H).

> Optical Rotation:  $[\alpha]_{23} = -39^{\circ}$  (c 1, MeOH).  $C_{19}H_{28}F_2N_2O_3$  (MW = 448); mass spectroscopy (MH<sup>+</sup>) 449.

### Example 220

# Synthesis of N-[(S)-2-Hydroxy-1-phenyleth-1-yl]-N'-(3,5-Difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)L-alanine (from Example B2 above) and (S)-2-hydroxy-1-phenyleth-1-ylamine
(e.g., (S)-phenylglycinol) (Aldrich), the title compound was prepared as a solid
(mp = 204-206°C). The reaction was monitored by tlc (Rf = 0.5 in 10%
MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using
5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.
NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.24$  (d, 3H), 4.38 (m, 1H), 4.80 (m, 2H). Optical Rotation:  $[\alpha]_{20} = +3.56^{\circ}$  @ 589 nm (c 1.10, DMSO).  $C_{19}H_{20}F_2N_2O_3$  (MW = 362.38); mass spectroscopy (MH<sup>+</sup>) 363.

15

### Example 221

## Synthesis of N-[N-(3,5-Difluorophenyl-α-fluoroacetyl)-L-alaniny]L-phenylglycine tert-Butyl Ester

Following General Procedure M and using 3,5-difluorophenyl-αfluoroacetic acid (from Example D1 above) and N-(L-alaniny)-L-phenylglycine
tert-butyl ester (prepared using N-BOC-L-alanine (Sigma) and L-phenylglycine
tert-butyl ester hydrochloride (Bachem) using General Procedure C, followed
by removal of the BOC group using General Procedure P), the title compound
was prepared as a clear oil. The reaction was monitored by tlc (Rf = 0.44 and
0.51 in 1:1 EtOAc/hexanes) and the product was purified by LC 2000
chromatography using 20% EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>)(1:1 mixture of diasteromers):  $\delta = 1.37$  (s, 9H), 1.39 (s, 9H), 1.42 (d, J = 7.0 Hz, 3H), 1.48 (d, J = 7.0 Hz, 3H), 3.80 (d, J = 7.0 Hz, 1H), 4.62 (pent, J = 7.0 Hz, 2H), 5.36 (d, J = 7.1 Hz, 1H), 5.42 (d, J =

7.2 Hz, 1H), 5.60 (d, J = 4.7 Hz, 1H), 5.73 (d, J = 4.7 Hz, 1H), 6.80 (m, 2H), 6.97 (m, 4H), 7.20-7.33 (m, 12H).

 $C_{23}H_{25}F_3N_2O_4$  (MW = 450.2); mass spectroscopy (MH<sup>+</sup>) 451.

### Example 222

5

# Synthesis of N-[(S)- $\alpha$ -Hydroxy- $\alpha'$ -phenylisopropyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and (S)- $\alpha$ -hydroxy- $\alpha'$ -

phenylisopropylamine (e.g., L-phenylalaninol) (Aldrich), the title compound was prepared as a solid. The reaction was monitored by tlc (Rf = 0.5 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel chromatography using 95:5 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

15  $^{1}$ H-nmr (CDCl<sub>3</sub>):  $\delta = 7.33-7.17$  (m, 5H), 6.72 (m, 3H), 6.62 (d, 1H), 6.32 (d, 1H), 4.43 (m, 1H), 4.10 (m, 1H), 3.61 (m, 2H), 3.45 (s, 2H), 2.84 (m, 2H), 1.32 (d, 3H).

Optical Rotation:  $[\alpha]_{23} = -60^{\circ}$  (c 1, MeOH).  $C_{20}H_{22}F_2N_2O_3$  (MW = 376); mass spectroscopy (MH<sup>+</sup>) 377.

20

### Example 223

# Synthesis of N-[(1S,2R)-1-Hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)
L-alanine (from Example B2 above) and (1S,2R)-1-hydroxy-1-phenylprop-2ylamine hydrochloride (e.g., (1S,2R)-norephedrine hydrochloride) (Aldrich),
the title compound was prepared as a solid. The reaction was monitored by tlc
(Rf = 0.5 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by silica gel
chromatography using 98:2 CHCl<sub>3</sub>/MeOH as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.31$  (m, 5H), 6.84-6.64 (m, 4H), 4.86 (m, 1H), 4.51 (m, 1H), 4.23 (m, 1H), 3.52 (s, 2H), 1.38 (d, 3H), 0.97 (d, 3H). Optical Rotation: [α]<sub>23</sub> = -44° (c 1, MeOH).  $C_{20}H_{22}F_2N_2O_3$  (MW = 376); mass spectroscopy (MH<sup>+</sup>) 377.

**5** .

10

20

25

### Example 224

### Synthesis of

### $N\hbox{-}2\hbox{-}Methoxyethyl\hbox{-}N'\hbox{-}[N\hbox{-}(3,5\hbox{-}difluor ophenylacetyl)\hbox{-}L\hbox{-}alaninyl] glycinamide}$

Following General Procedure K and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine methyl ester (from Example 28 above) and 2-methoxyethylamine (Aldrich), the title compound was prepared as a solid (mp = 148-151°C). The reaction was monitored by tlc (Rf = 0.53 in 10% MeOH/DCM + 1% TEA) and the product was purified by silica gel chromatography.

NMR data was as follows:

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 176.3$ , 173.4, 172.2, 166.5, 163.4, 150.4, 141.6, 114.1, 114.0, 113.9, 113.8, 103.9, 103.5, 72.2, 72.1, 59.4, 51.9, 44.0, 43.0, 40.7, 17.9.

 $C_{16}H_{21}F_2N_3O_4$  (MW = 357); mass spectroscopy (MH<sup>+</sup>) 358.

#### Example 225

#### Synthesis of

### N-[N-(3,5-Difluorophenylacetyl)-2-(S)-aminocyclohexylacetyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and N-[2-(S)-aminocyclohexylacetyl]-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-cyclohexylglycine (Sigma) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title

30

compound was prepared as a solid (mp = 234.4°C). The reaction was monitored by tlc (Rf = 0.65 in 5.95 MeOH/DCM) and the product was purified by recrystallization from ethyl acetate.

NMR data was as follows:

5  $^{1}$ H-nmr (DMSO- $d_3$ ):  $\delta = 8.85$  (d, J=6.5 Hz, 1H), 8.21 (d, J=8.9 Hz, 1H), 7.37 (s, 5H), 7.07 (m, 1H), 6.97 (d, 2H), 5.36 (d, J=6.4 Hz, 1H), 4.35 (t, J=7.7 Hz, 1H), 3.53 (m, 5H), 1.65 (m, 6H), 1.06 (m, 5H).

<sup>13</sup>C-nmr (DMSO- $d_3$ ):  $\delta = 171.065$ , 170.865, 168.953, 164.179, 163.967, 160.282, 160.070, 141.210, 141.058, 135.766, 128.657, 128.374, 128.004,

10 112.371, 112.238, 112.115, 111.981, 102.217, 101.808, 101.399, 56.568, 56.471, 41.467, 40.354, 28.884, 28.025, 25.926, 25.669.

 $C_{25}H_{28}N_2O_4F_2$  (MW = 458.51); mass spectroscopy (MH<sup>+</sup>) 458.1.

### Example 226

15

20

30

# Synthesis of N-[(1R,2S)-1-Hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and (1R,2S)-1-hydroxy-1-phenylprop-2-ylamine hydrochloride (e.g., (1R,2S)-norephedrine hydrochloride) (Aldrich), the title compound was prepared as a foam. The reaction was monitored by the (Rf = 0.5 in 9:1 CHCl<sub>2</sub>/MeOH).

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.35$  (m, 5H), 7.75 (m, 3H), 6.57 (d, 1H), 4.47 (d, 1H), 4.26 (m, 1H), 3.48 (s, 2H), 1.32 (d, 3H), 1.01 (d, 3H).

Optical Rotation:  $[\alpha]_{23} = -64^{\circ}$  (c 1, MeOH).

 $C_{20}H_{22}F_2N_2O_3$  (MW = 376); mass spectroscopy (MH<sup>+</sup>) 377.

### Example 227

Synthesis of

N-[(1R,2S)-1-hydroxy-1,2-diphenyleth-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

10.

15

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and (1R,2S)-2-amino-1,2-diphenylethanol (Aldrich), the title compound was prepared as a solid (mp = 217-219°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 7% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 0.76$  (d, 3H), 5.43 (d, 1H).

Optical Rotation:  $[\alpha]_{20} = -6.9^{\circ}$  @ 589 nm (c 1.10, DMSO).

 $C_{25}H_{24}F_2N_2O_3$  (MW = 438.48); mass spectroscopy (MH<sup>+</sup>) 439.

### Example 228

### Synthesis of

 $N\hbox{-}[(S)\hbox{-}1\hbox{-}Hydroxyhex\hbox{-}2\hbox{-}yl]\hbox{-}N'\hbox{-}(3,5\hbox{-}difluor ophenylacetyl)\hbox{-}L\hbox{-}alanina mide}$ 

Following General Procedure S and using methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate (from Example 1 above), the title compound was prepared as a solid (mp = 157-158.5°C). The reaction was monitored by tlc (Rf = 0.62 in 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>).

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 5.9$  (m, 2H), 5.8 (m, 1H), 4.37 (q, 1H), 3.8 (m,

20 1H), 3.58 (s, 2H), 3.5 (m, 2H), 1.4 (m, 9H), 0.9 (m, 3H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta = 175.4$ , 172.9, 166.7, 166.5, 163.5, 163.2,

141.8, 141.7, 113.9, 113.8, 113.7, 113.6, 103.9, 103.6, 103.2, 65.6, 53.2, 51.2, 43.3, 32.3, 29.7, 24.1, 18.7, 14.9.

 $C_{17}H_{24}F_2N_2O_3$  (MW = 342.39); mass spectroscopy (MH<sup>+</sup>) 343.

25

#### Example 229

Synthesis of  $N-[\alpha-Hydroxy-\alpha'-(4-hydroxyphenyl)isopropyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide$ 

Following General Procedure S and using N-[N-(3,5-

difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester (from Example 15

10

30

above), the title compound was prepared as a solid (mp = 179-183°C). The reaction was monitored by tlc (Rf = 0.42 in 10% MeOH/DCM) and the product was purified by recrystallization from MeOH/diethyl ether.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 6.82$  (d, J=8.3 Hz, 2H), 6.7 (m, 2H), 6.62 (t, J=9.1, 9.1 Hz, 1H), 6.47 (d, J=8.5 Hz, 2H), 4.1 (m, 1H), 3.7 (m, 1H), 3.34 (s, 2H), 3.2 (m, 2H), 2.5 (m, 2H), 1.08-0.94 (dd, J=7.1, 36.0, 7.1 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 175.0$ , 172.8, 157.4, 131.8, 131.8, 130.7, 116.6, 116.5, 113.9, 113.5, 64.1, 54.9, 51.1, 43.3, 37.4, 18.6.

C<sub>20</sub>H<sub>22</sub>F<sub>2</sub>N<sub>2</sub>O<sub>4</sub> (MW = 392); mass spectroscopy (MH<sup>+</sup>) = 393.

### Example 230

# Synthesis of N-2-Pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

Following General Procedure K and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester (from Example 94 above) and 2-(aminomethyl)pyridine (Aldrich), the title compound was prepared.

NMR data was as follows:

### Example 231

Synthesis of N-[ $\alpha$ -Hydroxy- $\alpha'$ -pyrid-2-ylisopropyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure S and using methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-pyridyl)propionate (from Example 19 above), the title compound was prepared as a solid (mp = 225-229°C). The reaction was monitored by tlc (Rf = 0.66 in 10% MeOH/DCM) and the product was purified by recrystallization from MeOH/diethyl ether.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.21$  (d, J=4.5 Hz, 1H), 7.46 (t, J=6.3, 7.6 Hz, 1H), 7.11 (d, J=7.6 Hz, 1H), 7.01 (t, J=5.5, 7.1 Hz, 1H), 6.70 (d, J=6.3 Hz, 2H), 6.62 (t, J=9.6, 9.0 Hz, 1H), 4.1 (m, 1H), 3.4 (m, 1H), 3.33 (s, 2H), 3.3 (m, 2H), 1.06 (d, J=7.0 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 172.754$ , 160.222, 150.134, 139.137, 126.198, 123.680, 113.936, 113.602, 103.578, 64.854, 53.689, 51.191, 43.304, 40.394, 18.769.

 $C_{19}H_{21}F_2N_3O_3$  (MW = 377); mass spectroscopy (MH<sup>+</sup>) 378.

15

5

10

### Example 232

# Synthesis of N-[ $\alpha$ -Hydroxy- $\alpha'$ -pyrid-4-ylisopropyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure S and using methyl N-[N-(3,5-20 difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4-pyridyl)propionate (from Example 23 above), the title compound was prepared as a solid (mp = 189-193°C). The reaction was monitored by tlc (Rf = 0.47 in 10% MeOH/DCM) and the product was purified by silica gel chromatography.

NMR data was as follows:

25  ${}^{1}\text{H-nmr} \text{ (CDCl}_{3}\text{): } \delta = 8.18 \text{ (d, J=5.6 Hz, 2H), 7.27 (d, J=5.6 Hz, 2H), 6.7 (m, 2H), 6.6 (m, 1H), 4.0 (m, 1H), 3.9 (m, 1H), 3.32 (s, 2H), 3.10 (s, 2H), 2.9 (m, 2H), 1.07 (d, J=7.2, 3H).$ 

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 175.8$ , 150.4, 150.2, 126.8, 113.9, 113.6, 103.6, 103.5, 72.0, 59.3, 55.2, 51.6, 42.9, 40.8, 38.3, 17.9.

30  $C_{19}H_{21}F_2N_3O_3$  (MW = 377); mass spectroscopy (MH<sup>+</sup>) 378.

### Example 233

# Synthesis of N-[(S)-1-Hydroxy-4-methylpent-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

### 5 Isomer A:

10

30

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and (S)-1-hydroxy-4-methylpent-2-ylamine (leucinol) (Bachem), the title compound was prepared as a solid (mp = 141-151°C). The reaction was monitored by tlc (Rf = 0.5 in 5% MeOH/methylene chloride) and the product was purified by recrystallization from EtOAc/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 8.15$  (s 1H), 7.5 (t, J=8 Hz, 1H), 6.80-6.55 (m, 3H), 4.15 (m, J=3.5 Hz, 1H), 3.7 (m 1H), 3.35 (s 2H), 3.22 (t, J=3 Hz, 2H), 1.4 (m, 1H), 1.1 (m, 5H), 0.7 (m, 6H).

15  $^{13}$ C-nmr (CD<sub>3</sub>OD):  $\delta = 175.4$ , 175.3, 173.0, 113.9, 113.9, 113.6, 113.5, 103.9, 103.6, 103.2, 66.1, 51.6, 51.4, 51.3, 51.3, 43.4, 41.7, 41.6, 26.5, 26.3, 24.3, 22.8, 22.7, 19.0, 18.7, 18.6.

 $C_{17}H_{24}N_2O_3F_2$  (MW = 342.19); mass spectroscopy (MH<sup>+</sup>) 343.

### Isomer B:

Following General Procedure B and using N-(3,5-difluorophenylacetyl)L-alanine (from Example B2 above) and (S)-1-hydroxy-4-methylpent-2-ylamine (leucinol) (Aldrich), the title compound was prepared as a solid (mp = 151-153°C). The reaction was monitored by tlc (Rf = 0.8 in 10% MeOH/DCM) and the product was purified by recrystallization, followed by flash column chromatography, followed by a preparative tlc using 10% MeOH/DCM as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 8.15$  (s 1H), 7.5 (t, J=8 Hz, 1H), 6.80-6.55 (m, 3H), 4.15 (m, J=3.5 Hz, 1H), 3.7 (m, 1H), 3.35 (s, 2H), 3.22 (t, J=3 Hz, 2H), 1.4 (m, 1H), 1.1(m, 5H), 0.7 (m, 6H).

10

20

25

30

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta$  = 175.2, 172.9, 166.6, 166.5, 141.7, 113.9, 113.9, 113.8, 113.6, 113.6, 103.9, 103.6, 103.2, 66.1, 51.2, 50.4, 50.1, 50.0, 49.8, 49.7, 49.6, 49.4, 49.38, 49.3, 49.0, 48.7, 43.4, 43.3, 41.7, 26.3, 24.3, 22.8, 18.7.

 $C_{17}H_{24}N_2O_3F_2$  (MW = 342.19); mass spectroscopy (MH<sup>+</sup>) 342.

### Example 234

### $Synthesis \ of \\ N-[1-Methoxyprop-2-yl]-N'-(3,5-difluor ophenylacetyl)-L-alanina mide$

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and 2-amino-1-methoxypropane (Aldrich), the title compound was prepared as a solid (mp = 152°C). The reaction was monitored by tlc (Rf = 0.45 in 5% MeOH/DCM) and the product was purified by recrystallization from methanol/water.

NMR data was as follows:

15  $^{1}$ H-nmr (CDCl<sub>3</sub>):  $\delta = 6.9$ -6.7 (m, 3H), 6.6 (d, J=7 Hz, 1H), 6.3 (m, 1H), 4.5 (m, J=7 Hz, 1H), 4.1 (m, 1H), 3.5 (s, 2H), 3.3 (m, 5H), 1.4 (d, J=7 Hz, 3H), 1.15 (t, J=8 Hz, 3H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 172.0, 113.0, 112.9, 112.62, 112.60, 103.7, 103.4, 78.0, 77.6, 77.2, 75.8, 75.7, 59.6, 59.58, 49.6, 49.5, 45.6, 45.6, 43.4, 19.4, 19.38, 18.9, 18.0.

 $C_{17}H_{20}N_2O_3F_2$  (MW = 314.14); mass spectroscopy (MH<sup>+</sup>) 315.

### Example 235

# Synthesis of N-[1-Hydroxy-3-methylbut-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and valinol (Bachem), the title compound was prepared as a solid (mp = 176-179°C). The reaction was monitored by tlc (Rf = 0.4 in 5% MeOH/DCM) and the product was purified by recrystallization from EtOAc/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 7.5$  (d, J=9 Hz, 1H), 6.8-6.5 (m, 3H), 4.15 (m, 1H), 3.45 (m, 2H), 3.35 (m, 3H), 1.65 (m, J=7 Hz, 1H), 1.20 (d, J=5 Hz, 3H), 0.7 (m, 6H).

5  $^{13}$ C-nmr (acetone- $d_6$ ):  $\delta = 113.7, 113.4, 103.0, 63.3, 57.7, 57.69, 50.5, 50.4, 43.2, 31.1, 30.8, 30.6, 30.5, 30.3, 30.2, 30.1, 29.9, 29.9, 29.8, 29.7, 29.6, 20.5, 20.4, 19.5, 19.1, 19.0, 18.8.$ 

 $C_{16}H_{22}N_2O_3F_2$  (MW = 329.19); mass spectroscopy (MH<sup>+</sup>) 329.

### Example 236

10

15

# Synthesis of Methyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-2-amino-2-(6-aminopyrid-2-yl)acetate

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(6-aminopyrid-2-yl)acetate (prepared from 2-(methoxyimino)-2-(6-aminopyrid-2-yl)acetic acid [CAS 71470-33-2] using General Procedures G and AC above), the title compound was prepared. The product was purified by LC 2000 preparative column chromatography using 1:1 EtOAc/hexanes as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.65$ -6.5 (m, 6H), 6.4 (d, J=8.19 Hz, 1H), 5.49-5.33 (m, 1H), 4.8-4.5 (m, 2H), 3.7 (s, 3H), 3.6 (s, 1H), 3.5 (s, 1H), 2.06 (bs, 2H), 1.44 (d, J=7.06 Hz, 1.5 H), 1.35 (d, 7.06 Hz, 1.5 H). C<sub>19</sub>H<sub>20</sub>N<sub>4</sub>O<sub>4</sub>F<sub>2</sub> (MW = 406.39); mass spectroscopy (MH<sup>+</sup>) 406.3.

### Example 237

25

30

# Synthesis of N-[1-Hydroxyprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and alanol (Bachem), the title compound was prepared as a solid (mp = 158-163°C). The reaction was monitored by tlc

(Rf = 0.7 in 10% MeOH/DCM) and the product was purified by recrystallization from ethyl acetate, followed by flash column chromatography using 10% MeOH/DCM.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 8.2$  (m, 1H), 7.6 (m, 1H), 4.1 (m, J=7 Hz, 1H), 3.7 (m, J=5 Hz, 1H), 3.35 (s, 2H), 3.25 (m, 2H), 1.15 (d, J=7 Hz, 3H), 0.9 (d, J=7 Hz, 3H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta = 175.1$ , 175.06, 172.9, 166.6, 166.5, 163.4, 163.2, 141.6, 113.9, 113.8, 113.7, 113.6, 103.9, 103.6, 103.2, 66.5, 51.4, 51.3, 51.3, 51.2, 50.4, 50.1, 49.8, 49.77, 49.6, 49.5, 49.3, 49.1, 49.0, 48.7, 43.3, 18.8, 17.5.

 $C_{14}H_{18}N_2O_3F_2$  (MW = 300); mass spectroscopy (MH+) 301.

### Example 238

15

25

5

10

### Synthesis of N-[(S)-2-methoxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using 3,5-difluorophenylacetic acid (Oakwood) and N-[(S)-2-methoxy-1-phenyleth-1-yl]-L-alaninamide (prepared from N-BOC-L-alanine (Sigma) and (S)-phenylglycinol methyl ether (from Example D15 above) using General Procedure C, followed by removal of the 20 BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 180-182°C). The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from 1chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.22$  (d, 3H), 3.23 (s, 3H). Optical Rotation:  $[\alpha]_{20} = +12.3^{\circ} @ 589 \text{ nm (c 1.04, DMSO)}.$  $C_{20}H_{22}F_2N_2O_3$  (MW = 376.41); mass spectroscopy (MH<sup>+</sup>) 377.

### Example 239

# Synthesis of N-[(S)-1-Methoxy-2-phenylprop-2-yl]- N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure B and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and L-phenylalaninol methyl ether hydrochloride (Fluka), the title compound was prepared as a fluffy solid. The product was purified by recrystallization from MeOH/EtOAc.

NMR data was as follows:

10 
<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta$  = 1.31 (d, J=7 Hz, 3H), 2.8 (d, J=7 Hz, 2H), 3.28 (d, J=3 Hz, 2H), 3.32 (s, 3H), 3.47 (s, 2H), 4.15-4.3 (m, 1H), 4.35-4.5 (m, 1H), 6.3-6.5 (m, 2H), 6.6-6.9 (m, 3H), 7.1-7.35 (m, 5H).

<sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta$  = 19.1, 37.8, 43.4, 49.6, 51.0, 59.6, 72.7, 103.4, 112.6, 113.0, 127.1, 129.0, 129.9, 138.3, 169.8, 172.1.

15

### Example 240

# Synthesis of N-[(S)-1-Acetoxyhex-2-yl]- N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure V and using N-[(S)-1-hydroxyhex-2-yl]-N'(3,5-difluorophenylacetyl)-L-alaninamide (from Example 228 above), the title compound was prepared as a solid (mp = 144-145 °C). The reaction was monitored by tlc (Rf = 0.42 in 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>).

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 6.7$  (m, 2H), 6.6 (m, 1H), 4.09 (q, 1H), 3.9-3.7 (m, 3H), 3.35 (s, 2H), 1.79 (s, 3H), 1.4-1.0 (m, 9H), 0.6 (m, 3H). <sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta = 175.5$ , 173.2, 172.8, 166.6, 166.5, 163.4, 163.2, 141.8, 141.7, 141.5, 113.9, 113.8, 113.7, 113.6, 103.9, 103.5, 103.2, 67.5, 51.2, 43.28, 43.26, 32.2, 29.6, 24.0, 21.3, 18.8, 14.8. C<sub>19</sub>H<sub>26</sub>F<sub>2</sub>N<sub>2</sub>O<sub>4</sub> (MW = 384.43); mass spectroscopy (MH<sup>+</sup>) 385.

10

### Example 241

# Synthesis of N-[(S)-1-(*tert*-Butylcarbonyloxy)-hex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure W and using N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (from Example 228 above) and trimethylacetyl chloride (Aldrich), the title compound was prepared as a solid (mp = 104-107.5°C). The reaction was monitored by tlc (Rf = 0.43 in 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) and the product was purified by preparative thin layer chromatography using 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 7.67$  (bd, 1H), 6.7 (m, 2H), 6.6 (m, 1H), 4.14 (q, 1H), 3.9-3.6 (m, 3H), 3.35 (s, 2H), 1.4-1.0 (m, 9H), 0.98 (s, 9H), 0.6 (m, 3H).

15  $^{13}$ C-nmr (CD<sub>3</sub>OD):  $\delta = 180.3$ , 175.3, 175.2, 172.8, 166.6, 166.5, 163.4, 163.2, 141.8, 141.7, 141.5, 133.9, 113.8, 113.7, 113.6, 103.9, 103.6, 103.2, 67.6, 51.1, 51.0, 43.3, 40.4, 32.4, 32.3, 29.5, 28.2, 24.0, 19.0, 14.9.  $C_{22}H_{32}F_2N_2O_4$  (MW = 426.51); mass spectroscopy (MH<sup>+</sup>) 427.5.

### Example 242

20

# Synthesis of N-[2-Hydroxy-1-(thien-2-yl)ethyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure S and using methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate (from Example 178 above), the title compound was prepared as a solid (mp = 201-202°C). The product was purified by trituration using 1:1 hexanes/EtOAc.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.4-8.25$  (m, 2H), 7.4-7.35 (m, 2H), 7.3-6.91 (m, 4H), 5.1-4.85 (m, 1H), 4.4-4.3 (m, 1H), 3.7-3.5 (m, 2H), 3.51 (s, 1H), 3.50 (s, 1H), 1.23-1.19 (overlaying doublets, 3H).

 $C_{21}H_{23}F_2N_2O_3$  (MW = 368.4); mass spectroscopy (MH<sup>+</sup>) 368.

### Example 243

## Synthesis of N-[(S)-2-hydroxy-2-methyl-1-phenylprop-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and (S)-2-hydroxy-2-methyl-1-phenylprop-1-ylamine (from Example D16 above), the title compound was prepared as a solid. The product was purified by recrystallization from methanol/ethyl acetate.

10 NMR data was as follows:

15

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.32$  (d, 1H), 8.11 (d, 1H), 7.20-7.33 (m, 5H), 7.08 (m, 1H), 6.96 (m, 2H), 4.68 (d, 1H), 4.53 (s, 1H), 4.95 (m, 1H), 3.50 (d, 2H), 1.25 (d, 3H), 1.08 (s, 3H), 0.98 (s, 3H).

Optical Rotation:  $[\alpha]_{23} = -11^{\circ}$  (c 1, MeOH).

 $C_{21}H_{24}F_2N_2O_3$  (MW = 390.42); mass spectroscopy (MH<sup>+</sup>) 391.

### Example 244

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-(thien-2-yl)glycinyl]-L-phenylalanine tert-Butyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-(2-thienyl)glycinyl--L-phenylglycine tert-butyl ester (prepared using N-(9-florenylmethoxycarbonyl)-L-(2-thienyl)glycine (prepared as described below) and L-phenylglycine tert-butyl ester hydrochloride using General Procedure AH, followed by deprotection using dicyclohexylamine in DMF and THF), the title compound was prepared as a solid (mp = 176-177°C). The product was purified by flash chromatography using EtOAc/dichloromethane as the eluent.

 $C_{26}H_{26}N_2O_4F_2$  (MW = 500.56); mass spectroscopy (MH<sup>+</sup>) 500.

Preparation of N-(9-Fluorenylmethoxycarbonyl)-L-(2-Thienyl)glycine:

A round bottom flask containing a magnetic stir bar under an atmosphere of

nitrogen at room temperature was charged with water, dioxane, sodium carbonate (2.5 eq.) and L- $\alpha$ -(2-thienyl)glycine (1.0 eq.) (Sigma). Stirring was initiated and the slurry was cooled in an ice bath. 9-Flurenylmethyl chloroformate was added portionwise to the reaction and stirring was continued in an ice bath for 4 hours followed by 8 hours at room temperature. The reaction mixture was poured onto water and extracted wilth diethyl ether. The aqueous layer was cooled in an ice bath and acidified with vigorous stirring to a pH of 2. The resulting solid was isolated via vacuum filtration, washed with water (3X) and dried under reduced pressure.

10

5

### Example 245

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinol

Following General Procedure E and using 3,5-difluorophenylacetic acid

(Aldrich) and L-phenylglycinyl-L-phenylglycinol (prepared from N-BOC-Lphenylgycine (Novabiochem) and L-phenylglycinol (Novabiochem) using
General Procedure AH, followed by removal of the BOC-group using General
Procedure P), the title compound was prepared as a solid (mp = 231.4°C).

The product was purified by crystallization from ethyl acetate.

20

 $C_{24}H_{22}N_2O_3F_2$  (MW = 424.45); mass spectroscopy (MH<sup>+</sup>) 424.9.

### Example 246

# Synthesis of N-[N-(Cyclopropaneacetyl)-L-phenylglycinyl]-L-phenylglycinol

25

Following General Procedure E and using cyclopropaneacetic acid (Aldrich) and L-phenylglycinyl-L-phenylglycinol (prepared from N-BOC-L-phenylgycine (Novabiochem) and L-phenylglycinol (Novabiochem) using General Procedure AH, followed by removal of the BOC-group using General

Procedure P), the title compound was prepared as a solid (mp = 202.5°C). The product was purified by crystallization from ethyl acetate.

 $C_{21}H_{24}N_2O_3$  (MW = 352.43); mass spectroscopy (MH<sup>+</sup>) 353.2.

### Example 247

5

## Synthesis of N-[N-(Cyclopentaneacetyl)-L-phenylglycinyl]-L-phenylglycinol

Following General Procedure E and using cyclopentaneacetic acid (Aldrich) and L-phenylglycinyl-L-phenylglycinol (prepared from N-BOC-L-phenylgycine (Novabiochem) and L-phenylglycinol (Novabiochem) using General Procedure AH, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 201.4°C). The product was purified by flash chromatography using MeOH/CH<sub>2</sub>CH<sub>2</sub> as the eluent.

15  $C_{23}H_{28}N_2O_3$  (MW = 380.49); mass spectroscopy (MH<sup>+</sup>) 381.4.

### Example 248

## Synthesis of N-[N-(3,5-Diffuorophenylacetyl)-D,L-phenylglycinyl]-D,L-phenylglycinamide

Following General Procedure AO and using N-[N-(3,5-difluorophenylacetyl)-D,L-phenylglycinyl]-D,L-phenylglycine methyl ester (from Example 99 above), the title compound was prepared as a solid (mp = 285.5-288.5°C).

 $C_{24}H_{21}N_3O_3F_2$  (MW = 437.45); mass spectroscopy (MH<sup>+</sup>) 437.1.

25

### Example 249

Synthesis of N-[N-(3,5-Diffuorophenylacetyl)-D,L-valinyl]-D,L-phenylglycinamide

Following General Procedure AO and using N-[N-(3,5-diffuorophenylacetyl)-L-valinyl]-L-phenylglycine methyl ester (from Example 94 above), the title compound was prepared as a solid (mp = 260.3-264.3°C). The product was purified by recrystallization from ethyl acetate/methanol.

5  $C_{21}H_{23}N_3O_3F_2$  (MW = 403.43); mass spectroscopy (MH<sup>+</sup>) 404.

### Example 250

### Synthesis of N-[N-(2-Thienylacetyl)-L-alaninyl]-L-phenylglycinamide

Following the General Procedures described herein, the title compound was prepared.

# Example 251 Synthesis of N-[N-(n-Caproyl)-L-alaninyl]-L-phenylglycinamide

Following the General Procedures described herein, the title compound was prepared.

### Example 252

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-norleucinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-phenylglycinyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-norleucine (Lancaster) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 188-189.5°C). The product was purified by flash chromatography using ethyl acetate/hexanes as the eluant.

 $C_{23}H_{26}N_2O_4F_2$  (MW = 432.47); mass spectroscopy (MH<sup>+</sup>) 432.

-- 327 --

#### Example 253

## Synthesis of N-[N-(3,5-Diffuorophenylacetyl)-L-norvalinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-norvalinyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-norvaline (Lancaster) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 204-205°C). The product was purified by flash chromatography using ethyl acetate/hexanes as the eluent.

 $C_{22}H_{24}N_2O_4F_2$  (MW = 418.44); mass spectroscopy (MH<sup>+</sup>) 418.3.

### Example 254

15

20

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-tent-leucinyl]L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-tert-leucinyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-tert-leucine (Bachem) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 176.4°C). The product was purified by flash chromatography using ethyl acetate/hexanes as the eluent.

 $C_{23}H_{26}N_2O_4F_2$  (MW = 432.47); mass spectroscopy (MH<sup>+</sup>) 432.0.

25

30

### Example 255

## Synthesis of N-[N-(3,5-Diffluorophenylacetyl)-L-isoleucinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-isoleucinyl-L-phenylglycine methyl ester hydrochloride

10

15

(prepared from N-BOC-L-isoleucine (Aldrich) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 228.8°C). The product was purified by flash chromatography using ethyl acetate/hexanes as the eluent.

 $C_{23}H_{26}N_2O_4F_2$  (MW = 432.46); mass spectroscopy (MH<sup>+</sup>) 433.4.

### Example 256

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-cyclohexylalaninyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-cyclohexylalaninyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-cyclohexylalanine (Sigma) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 174.8°C). The product was purified by flash chromatography using ethyl acetate/hexanes as the eluent.

 $C_{26}H_{30}N_2O_4F_2$  (MW = 472.53); mass spectroscopy (MH<sup>+</sup>) 473.2.

### Example 257

20

### Synthesis of N-[N-(3,5-Diffuorophenylacetyl)-(S)-2-amino-2-(cyclopropyl)acetyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and (S)-2-amino-2-(cyclopropyl)acetyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-(S)-2-amino-2-cyclopropylacetic acid (prepared from cyclopropylacetic acid (Lancaster) and (4S)-4-benzyl-2-oxaxolidinone (Aldrich) using the procedures described in Evans et al., J. Am. Chem. Soc., 1990, 112, 4011-4030 and references cited therein) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title

compound was prepared as a solid (mp = 225-226.5°C). The product was purified by flash chromatography using MeOH/CHCl<sub>3</sub> as the eluent.

 $C_{22}H_{22}N_2O_4F_2$  (MW = 416.42); mass spectroscopy (MH<sup>+</sup>) 417.3.

### Example 258

5

10

15

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-(S)-2-amino-2-(thien-3-yl)acetyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-(S)-2-amino-2-(thien-3-yl)acetyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-thien-3-ylglycine (prepared from L- $\alpha$ -2-thienylglycine (Sigma) using General Procedure AJ) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 229.3°C). The product was purified by crystallization from ethyl acetate/hexanes.

 $C_{23}H_{20}N_2O_4SF_2$  (MW = 458.49); mass spectroscopy (MH<sup>+</sup>) 458.

### Example 259

### Synthesis of

20

25

 $N\hbox{-}[N\hbox{-}(3,5\hbox{-}{\rm Difluorophenylacetyl})\hbox{-}(S)\hbox{-}2\hbox{-}amino\hbox{-}2\hbox{-}(thien\hbox{-}2\hbox{-}yl)acetyl]\hbox{-}$ L-phenylglycine Methyl Ester

Following General Procedure AH and using 3,5-difluorophenylacetic acid (Aldrich) and L-(S)-2-amino-2-(thien-2-yl)acetyl-L-phenylglycine methyl ester hydrochloride (prepared from N-BOC-L-thien-2-ylglycine (prepared from L- $\alpha$ -(thien-2-yl)glycine (Sigma) using General Procedure AI) and Lphenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by removal of the BOC-group using General Procedure P), the title compound was prepared as a solid (mp = 230.8°C). The product was purified by flash chromatographyl using MeOH/CH2CH2 as the eluant.

 $C_{23}H_{20}N_2O_4F_2S(MW = 458.49)$ ; mass spectroscopy (MH<sup>+</sup>) 458.

20

25

### Example 260

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-(4-fluorophenyl)glycinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid

(Aldrich) and L-(4-fluorophenyl)glycinyl-L-phenylglycine methyl ester
hydrochloride (prepared from N-Cbz-(4-fluorophenyl)glycine (prepared from (4fluorophenyl)glycine (prepared as described below) using General Procedure
AK) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General

Procedure E, followed by silica gel chromatography using 5% ethyl
acetate/toluene as the eluant and removal of the Cbz-group using General
Procedure AJ), the title compound was prepared as a solid (mp = 213.1°C).
The product was purified by flash chromatography using ethyl acetate/CHCl<sub>3</sub> as
the eluent.

 $C_{25}H_{21}N_2O_4F_2$  (MW = 470.44); mass spectroscopy (MH<sup>+</sup>) 470.1.

### Preparation of (4-fluorophenyl)glycine:

(S)-(-)-4-Benzyl-2-oxazolidinone (15.0 g, 93 mmol) (Aldrich) was dissolved in THF (100 mL). The solution was cooled to -70°C and reaction flask was purged twice with nitrogen. n-Butyl lithium (44.6 mL, 2.0M, 89 mmol) was added to form a solid precipitate which broke up on stirring wo afford a slurry. 4-Fluorophenylacetyl chloride 16.1 g, 93 mmol) (Aldrich) was added to afford a light green solution and stirring was continued for 45 minutes. The reaction mixture was then stirred at room temperature for 1 hour. The reaction mixture was then treated with saturated sodium bisulfate (100 mL) and ethyl acetate (100 mL). The organic phase was washed with water, followed by brine. The organic phase was then dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford an oil. The oil was crystallized to afford 24.4 g of 1-(4-fluorophenylacetyl)-(S)-(-)-4-Benzyl-2-oxazolidinone.

Potassium hexamethyldisilazane (140 mL, 0.5M, 70.0 mmol) was added to THF (80 mL). The solution was cooled to -50°C under nitrogen and a cold solution (-60°C) of 1-(4-fluorophenylacetyl)-(S)-(-)-4-benzyl-2-oxazolidinone (15.0 g, 46 mmol) in THF (100 mL). The resulting mixture was allowed to stir at -70°C for 1 hour and to warm to about -20°C. The mixture was recooled to -70°C and a cold solution (-65°C) of trasyl azide (21.6 g, 70.0 mmol) was added. The mixture was allowed to stir for about 15 min. while warming to -45°C and then glacial acetic acid (18 mL) was added. The mixture was then stirred at about 30°C for 3 hours. A precipitate formed and was removed by filtration. The filtrate was concentrated by 50 and then washed with water, saturated sodium bicarbonate solution and brine. The organic phase was dried over sodium sulfate, concentrated under reduced pressure to afford 37.7 g of crude 1-[2-(4-fluorophenyl)-2-azidoacetyl]-(S)-(-)-4-benzyl-2-oxazolidinone.

15

20

5

10

Crude 1-[2-(4-fluorophenyl)-2-azidoacetyl]-(S)-(-)-4-benzyl-2-oxazolidinone (10.0 g, 28.0 mmol) was dissolved in 100 mL of THF and 100 mL of methanol and trifluoroacetic acid (4.31 mL, 75.3 mmol) was added. Pallidium on carbon (10%, 2.0 g) was added and the mixture was hydrogenated on a Paar shaker at 50 psi overnight at room temperature. The reaction mixture was then filtered through a plug of Celite and the solid cake was rinsed with 100 mL of methanol. The filterate was concentrated to afford 1-[2-(4-fluorophenyl)-2-aminoacetyl]-(S)-(-)-4-benzyl-2-oxazolidinone trifluoroacetate salt as a yellowish oil.

25

To a mixture of THF and de-ionized water (50 mL/50 mL) was added 1-[2-(4-fluorophenyl)-2-aminoacetyl]-(S)-(-)-4-benzyl-2-oxazolidinone trifluoroacetate salt (4.14 g, 9.7 mmol) and lithium hydroxide monohydrate (1.22 g, 29 mmol). The homogenous solution was stirred for 2 hours at room temperature at which time Tlc indicated complete disappearance of starting material. The mixture was extracted with dichloromethane (3 x 100 mL) and

the aqueous phase was acidified to pH 2-3 while cooling in an ice-bath. A precipitate formed. The mixture was then cooled in an ice-bath for 1.5 hours and then filtered. The solid was washed with water followed by pentane to afford 4-fluorophenylglycine hydrochloride.

5

#### Example 261

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-D-(4-fluorophenyl)glycinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid

(Aldrich) and D-(4-fluorophenyl)glycinyl-L-phenylglycine methyl ester hydrochloride (prepared from N-Cbz-(4-fluorophenyl)glycine (prepared from (4-fluorophenyl)glycine (prepared as in Example 260) using General Procedure AK) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General Procedure E, followed by silica gel chromatography using 5% ethyl acetate/toluene as the eluant and removal of the Cbz-group using General Procedure AJ), the title compound was prepared as a solid (mp = 188.0°C). The product was purified by flash chromatography using ethyl acetate/CHCl<sub>3</sub> as the eluent.

 $C_{25}H_{21}N_2O_4F_3$  (MW = 470.44); mass spectroscopy (MH<sup>+</sup>) 470.1.

20

### Example 262

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-(4-methoxyphenyl)glycinyl]-L-phenylglycine Methyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid

(Aldrich) and L-(4-methoxyphenyl)glycinyl-L-phenylglycine methyl ester hydrochloride (prepared from N-Cbz-L-(4-methoxyphenyl)glycine (prepared from (4-methoxyphenyl)glycine (prepared by the Bucherer modification of the Strecker procedure as described in Greenstein et al., "The Chemistry of Amino Acids", Vol. 1, p. 698, Wiley, New York (1961)) using General Procedure

AK) and L-phenylglycine methyl ester hydrochloride (Aldrich) using General

Procedure E, followed by removal of the Cbz-group using General Procedure AJ), the title compound was prepared as a solid (mp = 224.6°C). The product was purified by flash chromatography using MeOH/CHCl<sub>3</sub> as the eluent.

 $C_{26}H_{24}N_2O_5F_2$  (MW = 482.48); mass spectroscopy (MH<sup>+</sup>) 482.1.

5

10

15

### Example 263

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure E and using 3,5-difluorophenylacetic acid (Aldrich) and L-phenylglycinyl-L-phenylglycine *tert*-butyl ester (prepared from N-Cbz-L-phenylglycine (Novabiochem) and L-phenylglycine *tert*-butyl ester hydrochloride (Novabiochem) using General Procedure AH, followed by removal of the Cbz-group using General Procedure AJ), the title compound was prepared as a solid (mp = 185.0°C). The product was purified by flash chromatography using ethyl acetate/CH<sub>2</sub>CH<sub>2</sub> as the eluant.

 $C_{28}H_{28}N_2O_4F_2$  (MW = 494.54); mass spectroscopy (MH<sup>+</sup>, minus  $CO_2$ -t-Bu) 393.

### Example 264

20

25

## Synthesis of N-[N-(Cyclopropylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure E and using cyclopropylacetic acid (Aldrich) and L-phenylglycinyl-L-phenylglycine *tert*-butyl ester hydrochloride (prepared from N-Cbz-L-phenylglycine (Novabiochem) and L-phenylglycine *tert*-butyl ester hydrochloride (Novabiochem) using General Procedure AH, followed by removal of the Cbz-group using General Procedure AJ), the title compound was prepared as a solid (mp = 187.5°C). The product was purified by crystallization from ethyl acetate.

 $C_{25}H_{30}N_2O_4$  (MW = 422.53); mass spectroscopy (MH<sup>+</sup>) 423.4.

### Example 265

## Synthesis of N-[N-(Cyclopentylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure E and using cyclopropylacetic acid (Aldrich) and L-phenylglycinyl-L-phenylglycine tert-butyl ester hydrochloride (prepared from N-Cbz-L-phenylglycine (Novabiochem) and L-phenylglycine tert-butyl ester hydrochloride (Novabiochem) using General Procedure AH, followed by removal of the Cbz-group using General Procedure AJ), the title compound was prepared as a solid (mp = 190.8°C). The product was purified by crystallization from ethyl acetate.

 $C_{27}H_{34}N_2O_4$  (MW = 450.58); mass spectroscopy (MH<sup>+</sup>) 451.

### Example 266

15

5

10

### Synthesis of N-[N-(t-Butylacetyl)-L-alaninyl]-L-phenylglycinamide

Following the General Procedures described herein, the title compound was prepared.

### Example 267

20

25

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(5-bromothien-2-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 5-bromo-2-thiophenecarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 227-228°C). The product was purified by recrystallization from ethyl acetate/hexanes.

 $C_{21}H_{24}N_3O_3BrS$  (MW = 515); mass spectroscopy (MH<sup>+</sup>) 515, 415.

10

15

20

25

30

### Example 268

### Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-(5-bromothien-2-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 5-bromo-2-thiophenecarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 216-217°C). The product was purified by recrystallization from ethyl acetate/hexanes.

 $C_{21}H_{24}N_3O_3BrS$  (MW = 515); mass spectroscopy (MH<sup>+</sup>) 515, 415.

#### Example 269

### Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(4-bromothien-2-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-bromo-2-thiophenecarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 246-247°C). The product was purified by recrystallization from ethyl acetate/hexanes.

 $C_{21}H_{24}N_3O_3BrS$  (MW = 515); mass spectroscopy (MH<sup>+</sup>) 515, 415.

#### Example 270

### Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(thien-2-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 2-thiophenecarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp

= 241-242°C). The product was purified by recrystallization from ethyl acetate/hexanes.

 $C_{21}H_{25}N_3O_3F_2S$  (MW = 438); mass spectroscopy (MH<sup>+</sup>) 438, 338.

#### Example 271

5

### Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide

difluorophenylacetyl)-L-alanine (from Example B2 above), 2thiophenecarboxaldehyde (Aldrich), (S)-(+)-α-methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 235-236°C). The product was purified by recrystallization from ethyl acetate/hexanes.

Following General Procedure AB and using N-(3,5-

 $C_{21}H_{25}N_3O_3F_2S$  (MW = 438); mass spectroscopy (MH<sup>+</sup>) 438, 338.

15

20

### Example 272

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(thien-3-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 3-thiophenecarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 240-241°C). The product was purified by recrystallization from ethyl acetate/hexanes.

25  $C_{21}H_{25}N_3O_3F_2S$  (MW = 438); mass spectroscopy (MH<sup>+</sup>) 438, 338.

#### Example 273

Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-(thien-3-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 3-thiophenecarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 245-246°C). The product was purified by recrystallization from ethyl acetate/hexanes.

 $C_{21}H_{25}N_3O_3F_2S$  (MW = 438); mass spectroscopy (MH<sup>+</sup>) 438, 338.

### Example 274

10

15

5

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), benzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 239-240°C). The reaction was monitored by tlc (Rf = 0.25 in 50% ethyl acetate/hexanes).

 $C_{23}H_{27}N_3O_3F_2$  (MW = 431.53); mass spectroscopy (MH<sup>+</sup>) 432.

### Example 275

20

25

30

# Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure AL and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), benzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 240-241 °C).  $C_{23}H_{27}N_3O_3F_2$  (MW = 431.53); mass spectroscopy (MH<sup>+</sup>) 432.

### Example 276

Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-(5-chlorothien-2-yl)glycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 5-chloro-2-thiophenecarboxaldehyde (from Example D17 above), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 195-198°C). The reaction was monitored by tlc (Rf = 0.15 in 50% ethyl acetate/hexanes).

 $C_{21}H_{24}N_3O_3F_2Cl$  (MW = 472); mass spectroscopy (MH<sup>+</sup>) 472.

### Example 277

10

15

5

## Synthesis of N-Cyclohexyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]- L-4-(phenyl)phenylglycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-biphenylcarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and cyclohexylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 300°C (dec.)). The reaction was monitored by tlc (Rf = 0.23 in 50% ethyl acetate/hexanes).

 $C_{31}H_{33}N_3O_3F_2$  (MW = 533.62); mass spectroscopy (MH<sup>+</sup>, minus cyclohexylamide) 408.2.

20

### Example 278

# Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-3-(phenoxy)phenylglycinamide

Following General Procedure AM and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 3-phenoxybenzaldehyde (Aldrich), (S)-(+)-α-methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.29 in 50% ethyl acetate/hexanes).

 $C_{29}H_{31}N_3O_4F_2$  (MW = 523.63); mass spectroscopy (MH<sup>+</sup>) 524.24.

### Example 279

## Synthesis of N-(S)-(-)- $\alpha$ -Methylbenzyl-N'-[N-(3,5-Difluorophenylacetyl) -L-alaninyl]-D,L-phenylglycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), benzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and (S)-(-)- $\alpha$ -methylbenzylisocyanide (from Example D18 above), the title compound was prepared.

10  $C_{27}H_{27}N_3O_3F_2$  (MW = 479.53); mass spectroscopy (MH<sup>+</sup>) 480.21.

By following the procedures set forth above, N-(R)-(+)- $\alpha$ -Methylbenzyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide was prepared merely by substitution of the appropriate isomer.

 $C_{27}H_{27}N_3O_3F_2$  (MW = 479.53); mass spectroscopy (MH<sup>+</sup>) 480.1.

15

20

5

### Example 280

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-3-(phenyl)phenylglycinamide

Following General Procedure AM and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 3-phenylbenzaldehyde (from Example D20 above), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.25 in 50% ethyl acetate/hexanes).

 $C_{29}H_{31}N_3O_3F_2$  (MW = 507.63); mass spectroscopy (MH<sup>+</sup>) 508.2.

25

#### Example 281

Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4-(ethyl)phenylglycinamide

10

15

Following General Procedure AM and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-ethylbenzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.20 in 50% ethyl acetate/hexanes).

 $C_{25}H_{31}N_3O_3F_2$  (MW = 459.59); mass spectroscopy (MH<sup>+</sup>) 460.2.

### Example 282

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-2-(phenyl)phenylglycinamide

Following General Procedure AM and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 2-phenylbenzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.15 in 50% ethyl acetate/hexanes).

 $C_{29}H_{31}N_3O_3F_2$  (MW = 507.63); mass spectroscopy (MH<sup>+</sup>, minus tert-butylamide) 409.

### Example 283

20

25

### Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-2-(benzyl)phenylglycinamide

Following General Procedure AM and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 2-(benzyl)benzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.19 in 50% ethyl acetate/hexanes).

 $C_{30}H_{33}N_3O_3F_2$  (MW = 521.66); mass spectroscopy (MH+) 522.26.

### Example 284

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-4-bromophenylglycinamide

Following General Procedure AM and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-bromobenzaldehyde (Aldrich), (S)-(+)-α-methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.06 in 50% ethyl acetate/hexanes).

 $C_{23}H_{26}N_3O_3F_2$  (MW = 510.42); mass spectroscopy (MH<sup>+</sup>) 512.1.

### Example 285

## $N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-\\ L-4-(cyclohexyl)phenylglycinamide$

Following General Procedure AL and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-(cyclohexyl)benzaldehyde (from Example D21 above), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 232-235°C).

20  $C_{29}H_{37}N_3O_3F_2$  (MW = 513.69); mass spectroscopy (MH<sup>+</sup>) 514.29.

### Example 286

## Synthesis of N-tent-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-4-(4-ethylphenyl)phenylglycinamide

- Following General Procedure AL and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4,4'-ethylbiphenylcarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 231-233°C).
- 30  $C_{31}H_{35}N_3O_3F_2$  (MW = 513.69); mass spectroscopy (MH<sup>+</sup>) 514.29.

20

25

30

### Example 287

## Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-4-(tert-butyl)phenylglycinamide

Following General Procedure AL and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-(tert-butyl)benzaldehyde (Aldrich), (S)-(+)-α-methylbenzylamine (Aldrich) and tert-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 280°C (dec.)). The reaction was monitored by tlc (Rf = 0.13 in 50% ethyl acetate/hexanes).

 $C_{27}H_{35}N_3O_3F_2$  (MW = 487.65); mass spectroscopy (MH<sup>+</sup>) 488.27.

### Example 288

Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-3-(4-chlorophenoxy)phenylglycinamide

Following General Procedure AL and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 3-(4-chlorophenoxy)benzaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and *tert*-butylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 192-195°C).

 $C_{29}H_{30}N_3O_4F_2Cl$  (MW = 558.07); mass spectroscopy (MH<sup>+</sup>) 558.20.

### Example 289

Synthesis of N-Cyclohexyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-4-(phenyl)phenylglycinamide

Following General Procedure AB and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above), 4-biphenylcarboxaldehyde (Aldrich), (S)-(+)- $\alpha$ -methylbenzylamine (Aldrich) and cyclohexylisocyanide (Aldrich), the title compound was prepared as a solid (mp = 290-291°C).

10

15

20

25

 $C_{31}H_{33}N_3O_3F_2$  (MW = 533.62); mass spectroscopy (MH<sup>+</sup>) 534.3.

### Example 290

## Synthesis of N-[N-(3,5-Difluorophenyl- $\alpha$ -hydroxyacetyl)-L-alaninyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure C and using 3,5-difluoromandelic acid (Fluorochem) and N-(L-alaninyl)-L-phenylglycine tert-butyl ester (prepared using N-BOC-L-alanine (Sigma) and L-phenylglycine tert-butyl ester hydrochloride (Bachem) using General Procedure C, followed by removal of the BOC group using General Procedure P), the title compound was prepared.

 $C_{23}H_{26}N_2O_5F_2$  (MW = 479.53). Elemental analysis: Calc. (%) C, 61.60; H 5.84; N, 6.25. Found (%) C, 61.32; H, 6.02; N, 6.17.

#### Example 291

### Synthesis of N-tert-Butyl-N'-[N-(3,5-Difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using 3,5-difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetic acid (from Example D23 above) and N-(L-alaninyl)-L-phenylglycine *tert*-butyl ester (prepared using N-BOC-L-alanine (Sigma) and L-phenylglycine *tert*-butyl ester hydrochloride (Bachem) using General Procedure C, followed by removal of the BOC group using General Procedure P), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.39 in 30% ethyl acetate/hexanes) and the product was purified by HPLC using 17% ethyl acetate/hexanes as the eluent.

 $C_{23}H_{24}N_2O_4F_4$  (MW = 468.49); mass spectroscopy (MH<sup>+</sup>) 469.17.

Example 292

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycine tert-Butyl Ester

15

20

25

30

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and D-phenylglycine *tert*-butyl ester (prepared from D-phenylglycine (Sigma) using General Procedure J), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.1 in 10% MeOH/CHCl<sub>3</sub>).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.64$  (d, 1H), 8. 38 (d, 1H), 7.34 (m, 5H), 7.09 (m, 1H), 6.99 (m, 2H), 5.27 (d, 1H), 4.45 (m, 1H), 3.32 (s, 2H) 1.28 (s, 9H), 1.18 (d, 3H).

10 Optical Rotation:  $[\alpha]_{20} = -103.58$  (c = 1, MeOH).  $C_{23}H_{26}N_2O_4F_2$  (MW = 432.47); mass spectroscopy (MH<sup>+</sup>) 433.

#### Example 293

Synthesis of N-[(S)-1-Oxo-1-phenylprop-2-yl]- N'-(3,5-difluorophenylacetyl)-L-alaninamide

By oxidation of N-[(1R,2S)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (from Example 226 above) using Jones reagent in acetone, the title compound was prepared. The reaction was monitored by tlc (Rf = 0.7 in 9:1 CHCl<sub>3</sub>/MeOH) and the product was purified by flash chromatography using 97:3 chloroform/methanol as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.98$  (m, 2H), 7.26 (m, 1H), 7.50 (m, 2H), 6.84 (m, 2H), 6.72 (m, 1H), 6.25 (d, 1H), 5.49 (m, 3H), 4.54 (m, 3H), 3.54 (s, 2H), 1.41 (d, 3H), 1.38 (d, 3H).

Optical Rotation:  $[\alpha]_{20} = -106^{\circ}$  @ 589 nm (c = 1, MeOH).  $C_{20}H_{20}F_2N_2O_3$  (MW = 374.39); mass spectroscopy (MH<sup>+</sup>) 374.

#### Example 294

Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-]-D,L-(pyrid-3-yl)glycine tert-Butyl Ester

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and *tert*-butyl 2-amino-2-(3-pyridyl)acetate (prepared as described in Kolar et al., *J. Heterocyclic Chem.*, 28, 171 (1991) and reference cited therein), the title compound was prepared. The reaction was monitored by the (Rf = 0.2 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 8.63$  (m, 1H), 8.54 (m, 1H), 7.62 (m, 1H), 7.45 (t, 1H), 7.26 (m, 1H), 6.82(m, 2H), 6.71 (m, 1H), 6.47 and 6.36 (d, 1H), 5.42 (d, 1H), 4.59 (m,1H), 3.52 and 3.47 (two s, 2H), 1.38 and 1.36 (s, 9H), 1.34 and 1.28 (two d. 3H).

 $C_{22}H_{25}N_3O_4F_2$  (MW = 433.46); mass spectroscopy (MH<sup>+</sup>) 434.

### Example 295

15

20

25

5

10

## Synthesis of [N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]morpholine

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and morpholine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.60$  and 8.49 (two d's, 1H), 8.49 (m, 1H), 7.25 (m, 5H), 7.18 (m, 2H), 6.95 (m, 1H), 5.82 (m, 1H), 4.38 (m, 1H), 3.52 (m, 10H), 1.21 and 1.12 (two d's, 3H).

 $C_{23}H_{25}N_3O_4F_2$  (MW = 445.47); mass spectroscopy (MH<sup>+</sup>) 446.

### Example 296

Synthesis of N-[N-(3,5-Diffuorophenylacetyl)-L-alaninyl]-D,L-(2-methoxy)phenylglycine Methyl Ester

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and methyl 2-amino-2-(2-methoxy)acetate (prepared from 2-methoxybenzaldehyde (Aldrich) using the Bucherer modification of the Strecker procedure as described in J. P. Greenstein et al., "The Chemistry of Amino Acids", Wiley: New York, 1961, Vol. 1, p. 698), the title compound was prepared. The reaction was monitored by tic (Rf = 0.3)in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

10

5

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.28$  (m, 2H), 6.93 (d, 1H), 6.88 (m, 2H), 6.69 (m, 2H), 6.34 (m, 1H), 5.67 (m, 1H), 4.52 (m, 1H), 3.81 (two s, 3H), 3.68 (two s, 3H), 3.59 and 3.45 (two s, 3H) 1.41 and 1.28 (two d, 3H).

 $C_{21}H_{22}N_2O_5F_2$  (MW = 420.42); mass spectroscopy (MH<sup>+</sup>) 420.

### Example 297

15

20

### Synthesis of

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-Butoxycarbonyl(hydroxyl amine) Ester

Following General Procedure C and using N-[N-(3,5difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and N-BOC hydroxyl amine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.35 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 2% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

25

<sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7.79$  (m, 1H), 7.41-7.28 (m, 5H), 6.78-6.59 (m, 3H), 5.52 (m, 1H), 4.69 (m, 1H), 3.38 (two s, 1H), 1.38 (d, 3H), 1.30 (s, 9H).

 $C_{24}H_{27}N_3O_6F_2$  (MW = 491.49); mass spectroscopy (MH<sup>+</sup>) 492.

#### Example 298

### Synthesis of N-Neopentyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and neopentylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.44$  (m, 1H), 7.41 (m, 2H), 7.31 (m, 3H), 7.12 (m, 1H), 6.99 (m, 2H), 5.50 (m, 1H), 4.47 (m, 1H), 3.52 (two s, 2H), 2.84 (m, 2H), 1.22 (m, 3H), 0.71 (s, 9H).

 $C_{24}H_{29}N_3O_3F_2$  (MW = 460); mass spectroscopy (MH<sup>+</sup>) 460.

15

20

10

### Example 299

### Synthesis of N-Tetrahydrofurfuryl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and tetrahydrofurfurylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.4 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.41$  (m, 2H), 7.32 (m, 5H), 7.08 (m, 1H), 6.99 (m, 2H), 5.48 (m, 1H), 4.42 (m, 1H), 3.85-3.54 (m, 3H), 3.48 (two s, 2H), 3.14 (m, 2H), 1.76 (m, 4H), 1.21 (m, 3H).

 $C_{24}H_{27}N_3O_4F_2$  (MW = 459.49); mass spectroscopy (MH<sup>+</sup>) 460.

10

15

25

#### Example 300

### Synthesis of N-Methoxy-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and methoxyamine hydrochloride (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.35 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.63$  (m, 1H), 8.35 (m, 1H), 7.34 (m, 5H), 7.12 (m, 1H), 6.99 (m, 2H), 5.23 (d, 1H), 4.42 (m, 1H), 3.58 (s, 3H), 3.51 (two s, 2H), 1.22 (d, 3H).

 $C_{20}H_{21}N_3O_4F_2$  (MW = 405); mass spectroscopy (MH<sup>+</sup>) 405.

### Example 301

### Synthesis of [N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]azetidine

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and azetidine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.6 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.61$  and 8.46 (two d, 1H), 8.33 (m, 1H), 7.34 (m, 5H), 7.19 (m, 1H), 6.99 (m, 2H), 5.36 (two d, 1H), 4.42 (m, 1H), 4.31 (m, 1H), 3.88 (m, 3H), 3.5 (two s, 2H), 2.36 (m, 2H), 1.18 (two d, 3H).  $C_{22}H_{23}N_3O_3F_2$  (MW = 415.44); mass spectroscopy (MH<sup>+</sup>) 416.

### Example 302

### Synthesis of N-Isobutyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and isobutylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.65 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.41$  (m, 1H), 8.22 (m, 1H), 7.38 (m, 2H), 7.09 (m, 1H), 6.98 (m, 2H), 5.52 (two d, 1H), 4.41 (m, 1H), 3.34 (two s, 2H), 2.85 (s, 2H), 1.61 (m, 1H), 1.20 (m, 3H), 0.92 (m, 6H).

 $C_{23}H_{27}N_3O_3F_2$  (MW = 431.48); mass spectroscopy (MH<sup>+</sup>) 432.

15

30

10

5

### Example 303

### Synthesis of N-Cyclopropanemethyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and (aminomethyl)cyclopropane (Aldrich), the title compound was prepared.

The reaction was monitored by tlc (Rf = 0.25 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

 $C_{23}H_{25}N_3O_3F_2$  (MW = 429.47); mass spectroscopy (MH<sup>+</sup>) 374.

### Example 304

### Synthesis of N-Methoxy-N-methyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

10

15

20

25

30

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and <math>N-methoxy-N-methylamine hydrochloride (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 2% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.65$  and 8.53 (two d, 1H), 8.37 (m, 1H), 7.31 (m, 5H), 7.12 (m, 1H), 6.98 (m, 2H), 5.91 and 5.82 (two d, 1H), 4.49 (m, 1H), 3.60-3.42 (m, 5H), 3.08 (two s, 3H), 1.21 and 1.16 (two d, 3H).  $C_{21}H_{23}N_3O_4F_2$  (MW = 419); mass spectroscopy (MH<sup>+</sup>) 420.

### Example 305

### Synthesis of N-2-Methylprop-2-enyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure M and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 1-amino-2-methylprop-2-ene (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.45 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.43$  (m, 2H), 7.40 (m, 2H), 7.29 (m, 3H), 7.11 (m, 1H), 6.98 (m, 2H), 5.46 (d, 1H), 4.68 (m, 2H), 4.42 (m, 1H), 3.6 (m, 2H), 3.49 (s, 2H), 1.56 (s, 3H), 1.21 (d, 3H).

 $C_{23}H_{25}N_3O_3F_2$  (MW = 429.47); mass spectroscopy (MH<sup>+</sup>) 430.

#### Example 306

Synthesis of N-(Pyrid-3-yl)methyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

10

15

20

25

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 3-(aminomethyl)pyridine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.1 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.82$  (m, 1H), 8.55 (m, 1H), 8.42 (m, 3H), 7.52 (m, 1H), 7.35 (m, 5H), 7.10 (m, 1H), 6.99 (m, 2H), 5.43 (d, 2H), 4.44 (m, 1H), 4.30 (bd, 2H) 3.52 (s, 2H) 1.26 (d, 3H).

 $C_{23}H_{24}N_4O_3F_2$  (MW = 466.49); mass spectroscopy (MH<sup>+</sup>) 467.

### Example 307

### Synthesis of N-(Pyrid-4-yl)methyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylgiycinamide

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 4-(aminomethyl)pyridine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.1 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.88$  (m, 1H), 8.54 (d, 1H), 8.43 (m, 3H), 7.37 (m, 4H), 7.12 (m, 3H), 6.9(m, 1H), 5.44 (d, 1H), 4.45 (m, 1H), 4.31 (d, 2H), 3.51 (s, 2H), 1.25 (d, 3H).

 $C_{23}H_{24}N_4O_3F_2$  (MW = 466.49); mass spectroscopy (MH<sup>+</sup>) 467.

#### Example 308

Synthesis of N-Furfuryl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

10

15

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and furfurylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.5 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.66$  (m, 1H), 8.45 (d, 1H), 8.39 (m, 1H), 7.57 (s, 1H), 7.33 (m, 5H), 7.09 (m, 1H), 6.99 (m, 2H), 6.36 (m, 1H), 6.12 (s, 1H), 5.41 (d, 1H), 4.22 (m, 1H), 3.52 (s, 2H) 1.24 (d, 3H).

 $C_{24}H_{23}N_3O_4F_2$  (MW = 455); mass spectroscopy (MH<sup>+</sup>) 456.

#### Example 309

### Synthesis of N-Cyclopentyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and cyclopentylamine (Aldrich), the title compound was prepared. The product was purified by recrystallization from ethanol.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.32$  (m, 2H), 8.16 (m, 1H), 7.33-7.20 (m, 5H), 7.04 (m, 1H), 6.93 (m, 2H), 5.34 (d, 1H), 4.37 (m, 1H), 3.9 (m, 1H), 3.49 (s, 2H), 1.80-1.29 (m, 8H), 1.19 (d, 3H).

 $C_{24}H_{27}N_3O_3F_2$  (MW = 443.49); mass spectroscopy (MH<sup>+</sup>) 444.

### Example 310

25

30

### Synthesis of

N-1-Benzylpiperidin-4-yl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 4-amino-1-benzylpiperdine (Aldrich), the title compound was prepared.

The reaction was monitored by tlc (Rf = 0.2 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 3% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

5

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.39$  (m, 2H), 8.21 (m, 1H), 7.30 (m, 5H), 7.11 (m, 1H), 6.98 (m, 2H), 5.39 (d, 1H), 4.21 (m, 1H), 3.54 (bm, 3H), 3.42 (bs, 2H), 2.70 (bm, 2H), 1.89 (bm, 2H), 1.71 (bm, 2H), 1.42 (3H), 1.22 (m, 3H).

 $C_{31}H_{34}N_4O_3F_2$  (MW = 548.64); mass spectroscopy (MH<sup>+</sup>) 548.

10

15

#### Example 311

### Synthesis of N,N-Dimethyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and dimethylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.65 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

20

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.13$  and 8.01 (two d, 1H), 7.32 (m, 5H), 6.78 (m, 2H), 6.63 (m, 1H), 5.88 (m, 1H), 4.72 (m, 1H), 3.45 (two s, 2H), 2.94 (two s, 6H), 1.32 and 1.17 (two d, 3H).

 $C_{21}H_{23}N_3O_3F_2$  (MW = 403.43); mass spectroscopy (MH<sup>+</sup>) 404.

### Example 312

25

### Synthesis of N-2,2,6,6-Tetramethylpiperidin-4-yl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-

#### D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-

difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above)

10

30

and 4-amino-2,2,6,6,-tetramethylpiperdine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.2 in 2% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 2% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.46$  (d, 1H), 8.33 (d, 1H), 8.12 (bm, 1H), 7.33 (m, 5H), 7.13 (m, 1H), 6.99 (m, 2H), 5.37 (d, 1H), 4.41 (m, 1H), 3.98 (m,1H), 3.52 (s, 2H), 1.67 (bm, 1H), 1.44 (bm, 1H), 1.22 (d, 3H), 1.01 (bm, 14H).

 $C_{28}H_{36}N_4O_3F_2$  (MW = 514.62); mass spectroscopy (MH<sup>+</sup>) 514.

#### Example 313

### Synthesis of N-2-Methylcyclohexyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylgiycinamide

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 2-methylcyclohexylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.4 in 2% MeOH/CHCl<sub>3</sub>).

NMR data was as follows:

20  $^{1}$ H-nmr (DMSO- $d_6$ ):  $\delta = 8.41$  (m, 2H), 8.0 (m, 1H), 7.33 (m, 5H), 7.11 (m, 1H), 6.99 (m, 2H), 5.35 (m, 1H), 4.41 (m, 1H), 3.52 (s, 2H), 3.18 (m,1H), 1.78-0.82 (m 11H), 0.81 (m, 3H).

 $C_{26}H_{31}N_3O_3F_2$  (MW = 472.5); mass spectroscopy (MH<sup>+</sup>) 472.

#### Example 314

25 Synthesis of

N-4-Methylcyclohexyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 4-methylcyclohexylamine (Aldrich), the title compound was prepared. The

10

15

25

30

reaction was monitored by tlc (Rf = 0.2 in 2% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 2% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.38$  (m, 2H), 8.08 (m, 1H), 7.33 (m, 5H), 7.09 (m, 1H), 7.01 (m, 2H), 5.54 and 5.36 (two d, 1H), 4.43 (m, 2H), 3.76 (m, 1H), 3.52 (s, 2H), 1.79-1.17 (m, 11H), 0.84 (d, 3H).

 $C_{26}H_{31}N_3O_3F_2$  (MW = 472.5); mass spectroscopy (MH<sup>+</sup>) 472.

### Example 315

Synthesis of N-1-Ethoxycarbonylpiperidin-4-yl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and 4-amino-1-ethoxycarbonylpiperdine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.2 in 2% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 2% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ): δ = 8.42 (m, 2H), 8.23 (m, 1H), 7.33 (m, 5H), 7.09 (m, 1H), 6.98 (m, 2H), 5.38 (m, 1H), 4.41 (m, 1H), 4.01 (q, 2H), 3.9-3.64 (m, 3H), 3.49 (s, 2H), 2.88 (bm, 2H), 1.75 (m, 1H), 1.54 (m, 1H), 1.2 (m, 6H).

 $C_{27}H_{32}N_4O_5F_2$  (MW = 530.57); mass spectroscopy (MH<sup>+</sup>) 531.

Example 316

Synthesis of N-Methyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and N-methyl-(S)-2-amino-2-

phenylacetamide [CAS 129213-83-8], the title compound was prepared. The reaction was monitored by tlc (Rf = 0.2 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by flash chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

5  $^{1}$ H-nmr (DMSO- $d_{6}$ ):  $\delta = 8.43$  (m, 2H), 8.21 (m, 1H), 7.36 (m, 5H), 7.09 (m, 1H), 6.95 (m, 2H), 5.36 (m, 1H), 4.40 (m, 1H), 3.41 (s, 2H), 2.56 (d, 3H), 1.22 (d, 3H).

Optical Rotation:  $[\alpha]_{20} = -67$  (c = 1, MeOH).

 $C_{20}H_{21}N_3O_3F_2 \cdot 0.75 H_2O (MW = 403.43);$  mass spectroscopy (MH+)

10 404.

### Example 317

### Synthesis of N-tert-Butoxy-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

Following General Procedure C and using N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and <math>O-(tert-butoxy)hydroxylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.65 in 10% MeOH/CHCl<sub>3</sub>).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 8.72$  and 8.58 (two d, 1H), 8.39 (m, 1H), 7.37 (m, 5H), 7.10 (m, 1H), 6.99 (m, 2H), 5.41 (m, 1H), 4.46 (m, 1H), 3.51 (two s, 3H), 1.22 (m, 3H), 1.09 (s, 9H).

 $C_{23}H_{27}N_3O_4F_2$  (MW = 447.48); mass spectroscopy (MH<sup>+</sup>) 448.

### Example 318

25

### Synthesis of

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-Butyl(hydroxylamine) Ester

Following General Procedure C and using N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above)

and N-(tert-butoxy)hydroxylamine (Aldrich), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.65 in 10% MeOH/CHCl<sub>3</sub>).  $C_{23}H_{27}N_3O_4F_2 \cdot 0.25 H_2O \text{ (MW = 447.48); mass spectroscopy (MH+)}$ 

448.

5

### Example 319

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine Hydrazide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine methyl ester (2.0 g, 5.1 mmol) (from Example 111 above) was stirred in ethanol (40 mL) 10 and anhydrous hydrazine (0.3 mL, 10 mmol) (Aldrich) was added. The solution was heated at reflux for 12 hours and then allowed to cool to ambient temperature with stirring. A title compound was collected as a white solid by filtration, washing with ethanol and dring in a vacuum oven (52% yield).

15

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.20$  (t, 3H), 5.41 (m, 1H).  $C_{19}H_{20}N_3O_4F_2$  (MW = 390.39); mass spectroscopy (MH<sup>+</sup>) 390.

### Example 320

20

25

### Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine Acetohydrazonate

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine hydrazide (0.5 g, 1.3 mmol) (from Example 319 above) was heated at reflux in triethylorthoacetate (40 mL). After 14 hours, the reaction mixture was concentrated under reduced pressure to afford the title compound as a white solid (84% yield). The reaction was monitored by tlc (Rf = 0.65 in 10%MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

30 <sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 4.03$  (q, 2H), 5.54 (m, 1H).

10

15

30

 $C_{23}H_{26}N_4O_4F_2$  (MW = 460.49); mass spectroscopy (MH<sup>+</sup>) 460.

### Example 321

## Synthesis of N-[N-(Phenylacetyl)-L-alaninyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure C and using phenylacetic acid (Aldrich) and L-alaninyl-L-phenylglycine tent-butyl ester (prepared using N-BOC-L-alanine (Sigma) and L-phenylglycine tent-butyl ester hydrochloride (Bachem) using General Procedure C, followed by removal of the BOC group using General Procedure P), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.25 in 3% MeOH/CHCl<sub>3</sub>) and the product was purified by crystallization from chlorobutane/hexanes.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 4.43$  (m, 1H), 5.20 (d, 1H).  $C_{23}H_{28}N_2O_4$  (MW = 396.49); mass spectroscopy (MH<sup>+</sup>) 397.

#### Example 322

### Synthesis of N-4-(phenyl)butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)L-alanine (from Example B2 above) and N-4-(phenyl)butyl-L-phenylglycinamide
(prepared from N-BOC-L-phenylglycine (Advanced Chemtech) and 4phenylbutylamine (Aldrich) using General Procedure C, followed by removal of
the BOC-group using General Procedure P), the title compound was prepared.

The reaction was monitored by tlc (Rf = 0.45 in 5% MeOH/CHCl<sub>3</sub>) and the
product was purified by trituration in water, followed by trituration in
acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 4.42$  (m, 1H), 5.37 (d, 1H). C<sub>29</sub>H<sub>31</sub>N<sub>3</sub>O<sub>3</sub>F<sub>2</sub> (MW = 507.5); mass spectroscopy (MH<sup>+</sup>) 507.

### Example 323

## Synthesis of N-3-(4-Iodophenyl)propyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)L-alanine (from Example B2 above) and N-3-(4-iodophenyl)propyl-Lphenylglycinamide (prepared from N-BOC-L-phenylglycine (Advanced
Chemtech) and 3-(4-iodophenyl)propylamine (from Example D26 above) using
General Procedure C, followed by removal of the BOC-group using General
Procedure P), the title compound was prepared. The product was purified by
trituration in water, followed by trituration in ethanol.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 4.41$  (q, 2H), 5.35 (m, 1H).  $C_{28}H_{28}N_3O_4F_2I$  (MW = 635.45); mass spectroscopy (MH<sup>+</sup>) 635.

15

### Example 324

## Synthesis of N-6-(Amino)hexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]- D,L-phenylglycinamide Hydrochloride

Following General Procedure C and using N-[N-(3,5-20 difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine (from Example D25 above) and N-BOC-1,6-hexanediamine (Fluka), followed by removal of the BOC-group using General Procedure P, the title compound was prepared. The product was isolated as a white solid.

NMR data was as follows:

25  ${}^{1}\text{H-nmr}$  (DMSO- $d_{6}$ ):  $\delta = 4.41$  (m, 1H), 5.40 (t, 1H).  $C_{25}H_{32}N_{4}O_{3}F_{2}$  (MW = 474.56); mass spectroscopy (MH<sup>+</sup>) 475.

### Example 325

 $\label{eq:N-1-Phthalimido} Synthesis of $N$-1-(Phthalimido) pent-2-yl-$N'-(3,5-difluorophenylacetyl)-L-alanina mide$ 

Following General Procedure C and using N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) and 2-amino-1-phthalimidopentane hydrochloride (from Example D27 above), the title compound was prepared. The reaction was monitored by tlc (Rf = 0.3 in 5% MeOH/CHCl<sub>3</sub>) and the product was purified by silica gel chromatography using 5% MeOH/CHCl<sub>3</sub> as the eluent, followed by recrystallization from chlorobutane/acetonitrile.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 4.1$  (m, 2H), 7.83 (bs, 4H).  $C_{24}H_{25}N_3O_4F_2$  (MW = 457.48); mass spectroscopy (MH<sup>+</sup>) 457.

10

15

5

### Example 326

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-(3,5-difluorophenyl)glycine Methyl Ester

Following General Procedure AN and using N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenyl)glycine (from Example D30 above) and L-3,5-difluorophenylglycine methyl ester (from Example D29 above), the title compound was prepared. The product was purified by crystallization.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 9.40$  (m, 1H), 9.0 (m, 1H), 6.80-7.70 (m, 20 9H), 5.45 (d, 1H), 5.25 (m, 1H), 3.55-365 (m, 5H).

## Example 327

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-norleucine

Following General Procedure AF and using THF/H<sub>2</sub>O (1:1) on N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-norleucine methyl ester, the title compound was prepared as a solid (mp = 158.5-160.5°C). The reaction was monitored by tlc (Rf = 0.29 in 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>).

NMR data was as follows:

10

15

<sup>1</sup>H-nmr (CD<sub>3</sub>OD):  $\delta = 8.46$  (bd, J = 6.71, 1H), 8.25 (bd, J = 7.69, 1H), 7.00-6.79 (m, 3H), 4.50-4.35 (m, 2H), 3.61 (d, 2H), 1.94-1.79 (m, 1H), 1.78-1.60 (m, (includes d at 1.40, J = 7.14, 3H), 0.92 (m, 3H).

<sup>13</sup>C-nmr (CD<sub>3</sub>OD):  $\delta = 176.0, 175.5, 172.9, 166.6, 166.5, 163.4,$ 163.2, 141.7, 141.6, 141.5, 113.9, 113.8, 113.7, 113.6, 103.9, 103.6, 103.2, 54.1, 50.9, 43.3, 32.9, 29.4, 23.8, 18.6, 14.8.

 $C_{17}H_{22}N_2O_4F_2$  (MW = 356.37); mass spectroscopy (MH<sup>+</sup>) 357.

### Example 328

## Synthesis of

N-[N-(Cyclopentaneacetyl)-L-alaninyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure D and using cyclopentylacetic acid (Aldrich) and L-alaninyl-L-phenylglycine tert-butyl ester (prepared from N-CBZ-L-alanine (Sigma) and L-phenylglycine tert-butyl ester hydrochloride (Bachem) using General Procedure C, followed by removal of the CBZ-group using General Procedure Y), the title compound was prepared as a solid (mp = 133-138°C). The reaction was monitored by tlc (Rf = 0.48 in 50%EtOAc/hexanes) and the product was purified by flash chromatography using 25-50% EtOAc/hexanes as the eluent.

20 NMR data was as follows:

> <sup>1</sup>H-nmr (CDCl<sub>3</sub>):  $\delta = 7/86$  (bd, J = 7.2 Hz, 1H), 7.30-7.15 (m, 5H), 6.81 (bd, J = 7.82 Hz, 1H), 5.34 (d, J = 7.20 Hz, 1H0, 4.72 (quint, J = 7.2Hz, 1H), 2.04 (m, 3H), 1.75-1.28 (m (includes s at 1.34, 9H) 18H), 1.1-0.9 (m, 2H).

25 <sup>13</sup>C-nmr (CDCl<sub>3</sub>):  $\delta = 173.3$ , 172.8, 170.0, 137.1, 129.2, 128.6, 127.7, 82.7, 57.7, 48.9, 43.0, 37.6, 32.9, 28.3, 25.4, 19.3.

 $C_{22}H_{32}N_2O_4$  (MW = 388.51); mass spectroscopy (MH<sup>+</sup>) 389.5.

## Example 329

## Synthesis of N-[N-(2,5-Dichlorophenylmercaptoacetyl)-L-alaninyl]-L-phenylglycine Methyl Ester

2,5-Dichlorophenylmercaptoacetic acid (TCI America, Portland, OR)

(237 mg) was converted to the acid chloride as described in the General

Procedure A" and utilized to acylate methyl L-alaninyl-L-phenylglycinate as
described in General Procedure B". The title compound (210 mg) was isolated
as crystals from ethyl ether.

10 NMR data was as follows:

<sup>1</sup>H-nmr (DMSO-d<sub>6</sub>)  $\delta = 8.85$  (d, 1H), 8.20 (d; 1H), 6.70-7.45 (m, 8H), 5.45 (d, 1H), 4.45-4.65 (m, 3H), 3.65 (s, 3H), 1.30 (d, 3H).

 $C_{20}H_{20}Cl_2N_2O_4S$  (MW = 455.363) mass spectroscopy (MH<sup>+</sup>) 454.1.

Anal. Calcd. for  $C_{20}H_{20}Cl_2N_2O_4S$ : C, 52.75 H, 4.42 N, 6.15; Found:

15 C, 53.58 H, 5.01, N, 6.34.

### Example 330

## Synthesis of N-[N-(3,4-Dichlorophenylmercptoacetyl)-L-alaninyl]-L-phenylglycine Methyl Ester

3,4-Dichlorophenylmercaptoacetic acid (*J. Med. Chem.*, 15(9), 940-944 (1972)) (237 mg) was converted to the acid chloride as described in the General Procedure A" and utilized to acylate methyl L-alaninyl-L-phenylglycinate as described in General Procedure B". The title compound (182 mg) was isolated as cyrstals from ethyl ether.

NMR data was as follows:

 $^{1}\text{H-nmr}$  (DMSO-d<sub>6</sub>)  $\delta = 8.8$  (d, 1H), 8.40 (d, 1H), 7.25-7.65 (m, 8H), 5.40 (d, 1H),

4.45 (m, 1H), 3.80 (m, 2H), 3.65 (s, 3H), 1.25 (d, 3H).

 $C_{20}H_{20}Cl_2N_2O_4S$  (MW = 455.363); mass spectroscopy (MH+) 454.1

Anal. Calcd. for  $C_{20}H_{20}Cl_1N_2O_4S$ : C, 52.75 H, 4.42 N, 6.15; Found: C, 53.05 H,

4.67 N, 6.26.

25

### Example 331

## Synthesis of N-[N-(3,5-Difluorophenoxyacetyl)-L-alaninyl]-L-phenylglycine Methyl Ester

3,5-Difluorophenoxyacetic acid [prepared by refluxing an aqueous mixture of 3,5-difluorophenol (Aldrich), 2-chloroacetic acid, and NaOH] (188 mg) was converted to the acid chloride as described in the General Procedure A" and utilized to acylate methyl L-alaninyl-L-phenylglycinate as described in General Procedure B". The title compound (210 mg) was isolated as crystals from ethyl ether.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO-d<sub>6</sub>)  $\delta = 8.85$  (d, 1H), 8.20 (d, 1H), 6.70-7.45 (m, 8H), 5.45 (d, 1H), 4.45-4.65 (m, 3H), 3.65 (s, 3H), 1.30 (d, 3H).

 $C_{20}H_{20}F_2N_2O_5$  (MW = 406.39); mass spectroscopy (MH<sup>+</sup>) 406.3.

15 Anal. Calcd. for  $C_{20}H_{20}F_2N_2O_5$ : C, 59.11 H, 4.96 N, 6.89; Found: C, 53.34 H, 4.80 N, 6.94.

### Example 332

#### Synthesis of

## Methyl N-[N-(3,5-Difluorophenoxyacetyl)-L-alaninyl]-L-2,3-dihydroisoindole-1-carboxylate

Following General Procedure AN, L-2,3-Dihydro-1*H*-isoindole-1-carboxylic acid methyl ester hydrochloride (*Gazz. Chim. Ital.*, **106** (1-2) p. 65-75 (1976)) (417 mg) was coupled to N-(3,5-difluorophenylacetyl-L-alanine (from Example B2) to provide the title compound (150 mg).

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO-d<sub>6</sub>)  $\delta = 8.55$  (d, 1H), 6.85-7.45 (m, 7H), 5.50(m, 1H), 4.95(s, 1H),

4.55-4.90(m, 2+H), 3.65 (m, 3H), 1.30 (m, 3H).

 $C_{21}H_{20}F_2N_2O_4$  (MW = 402.40); mass spectroscopy (MH<sup>+</sup>) 402.3.

10

15

### Example 333

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-1-amino-1,3-diphenylpropane-2-one

To a solution of 200 mg of N-methoxy-N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide (from Example 304 above) in THF was added 1.91 mL of a 2M solution of benzyl magnesium bromide in THF (Aldrich) at 0°C. The reaction mixture was stirred at ambient temperature for 72 hours, and was subsequently quenched by addition of water. The reaction mixture was partitioned between ethyl acetate and water and the organic phase was washed with 1N HCl solution. Following removal of solvent under reduced pressure, the crude ketone was purified by chromatography on silica gel, eluting with ethyl acetate, to afford 62 mg of the title compound as a 1:1 mixture of phenyl diastereomers.

NMR data was as follows:

<sup>1</sup>H-Nmr (CDCl<sub>3</sub>) (approx 1:1 mixture of diastereomers)  $\delta = 7.2$ -7.5 (m, 8H), 7.0-7.1 (m, 2H), 6.7-6.9 (m, 4H), 6.2 (m, 1H), 5.5(t, 1H), 3.5-3.6 (m, 2H), 1.28-1.45 (doublets in 1:1 ratio, 3H).

#### Example 334

20

## Synthesis of N-[N-(3,5-Difluorophenylacetyl)L-alaninyl]-D,L-phenylglycine Thiocarboxamide

Step A - <u>Preparation of *t*-butoxycarbonyl-phenylglycine</u> thiocarboxamide:

To a suspension of 500 mg (2.00 mmol) t-butoxycarbonyl-L-phenylglycine carboxamide (prepared as in Example 141) in 50 mL of dry toluene was added 808 mg (2.00 mmol) Lawesson's reagent (Aldrich). The reaction mixture was heated to 95°C for 5 min. Cooling to ambient temperature and dilution with 1:1 ethyl acetate/hexanes resulted in precipitation of insoluble material. Removal of the soluble phase, followed by additional washing of the solids and combination of the soluble phase, and removal of solvent afforded crude thiocarboxamide as a semisolid. Purification by

chromatography on silica gel, eluting with ethyl acetate afforded 364 mg of thiocarboxamide.

## Step B - Preparation of phenylglycine thiocarboxamide hydrobromide:

A solution of 364 mg of t-butoxycarbonyl phenylglycine

thiocarboxamide in 4 mL 30% HBr in acetic acid was stirred for 1 hour. The
volitile materials were removed under reduced pressure and the crude
phenylglycine thiocarboxamide hydrobromide was obtained as a pale solid. The
material was utilized without further purification.

10 (from B2) in 30 mL of dichoromethane was added 383 mg of EDCI, 270 mg of HOBT hydrate, followed by 350 μL of diisopropylethylamine. To this suspension was added phenylglycine thiocarboxamide hydrobromide in dichloromethane. The reaction mixture was stirred at ambient temperature for 72 hours. The reaction mixture was partioned between water and dichloromethane and the organic phase was washed with 1N HCl solution, followed by saturated aqueous sodium bicarbonate solution. Removal of solvent afforded the crude product, which was purified by chromatography on silica gel, eluting with ethyl acetate, to afford 271 mg of the title compound (approximately 3:2 mixture of phenylglycine diastereomers) as a pale solid.

NMR data was as follows:

 $^{1}\text{H-Nmr}$  (CDCl<sub>3</sub>) (approx 3:2 mixture of diastereomers):  $\delta = 7.3-7.7$  (m, 8H), 6.7-6.8 (m, 4H).

### Example 335

25

20

## Synthesis of N-[N-(3,5-Difluorophenyl-2-oxoacetyl)-L-alaninyl]-L-phenylglycine tert-Butyl Ester

Following General Procedure C and using L-alaninyl-L-phenylglycine tert-butyl ester (prepared as described in Example 321) and 3,5-difluorophenylglyoxylate (prepared as described in *J. Org. Chem.*, 45(14),

28883 (1980)), the title compound was prepared as a solid. The product was purified by slurrying with EtOAc/hexanes.

Elemental Anal.: Calc.(%) C, 61.88, H, 5.42, N, 6.27; Found: C, 62.15, H, 5.51, N, 6.18.

5

### Example 336

## N-(2-Hydroxy-1-phenyleth-1-yl)-N'-[N-(3,5-Difluorophenylacetyl)-L-phenylglycinyl]-L-alaninamide

Following General Procedure C and using N-(3,5-difluorophenylacetyl)L-phenylglycinyl-L-alanine (prepared from N-(3,5-difluorophenylacetyl)-Lphenylglycinyl-L-alanine ethyl ester) and (S)-phenylglycinol (Aldrich), the title
compound was prepared (m.p. = 269-272°C). The reaction was monitored by
tlc (Rf = 0.3 in 10% MeOH/CHCl<sub>3</sub>) and the product was purified by
chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.25$  (d, 3H), 8.01 (d, 1H), 8.52 (d, 1H), 8.82 (d, 1H).

Optical Rotation:  $[\alpha]_{20} = -62.7 \ @ 589 \text{ nm} (c = 1.02, DMSO).$  $C_{27}H_{27}N_3O_3F_2 (MW = 495.53); \text{ mass spectroscopy (MH+) 496.}$ 

20

### Example 337

## Synthesis of N-(2-Hydroxyeth-1-yl)-N'-[N-(3,5-Difluorophenylacetyl)-L-alanyl]-L-phenylglycinamide

L-alanine (from Example B2 above) and L-phenylglycine (2-hydroxyethyl)amide hydrochloride (prepared from N-BOC-L-phenylglycine (Bachem) and 2-aminoethanol (Aldrich) using General Procedure C, followed by removal of the BOC group using General Procedure P), the title compound was prepared. The product was purified by chromatography using 10% MeOH/CHCl<sub>3</sub> as the eluent, followed by crystallization from EtOH.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSO- $d_6$ ):  $\delta = 1.22$  (d, 3H), 5.42 (d, 1H).

Optical Rotation:  $[\alpha]_{20} = +8.77 @ 589 \text{ nm (c} = 1.03, DMSO).$ 

 $C_{21}H_{23}N_3O_4F_2$  (MW = 419.43); mass spectroscopy (MH<sup>+</sup>) 420.

5

### Example 338

# Synthesis of N-(4-(4-Azido-2-hydroxybenzamido)but-1-yl)-N'-[N-(3,5-Difluorophenylacetyl)-L-alanyl]-L-phenylglycinamide

Following General Procedure A and using N-(3,5-difluorophenylacetyl)L-alanyl-L-phenylglycine (prepared as described herein) and 4-(4azidosalicylamido)butylamine (Pierce Chemical), the title compound was
prepared as a light sensitive solid. The reaction was conducted under low light
conditions and the reaction vessel was protected from light. The reaction was
monitered by tlc (R<sub>f</sub> = 0.2 in 2.5% MeOH/dichloromethane).

NMR data was as follows:

<sup>1</sup>H-nmr (CD<sub>3</sub>OH/CDCl<sub>3</sub>):  $\delta = 7.72$  (d, 2H), 7.30 (m, 5H), 6.84 (m, 2H), 6.73 (m, 1H), 6.54 (m, 2H), 5.34 (s, 1H), 4.39 (q, 1H), 3.56 (s, 2H), 3.31 (bs, 2H), 3.21 (bs, 2H), 1.57 (bs, 4H), 1.35 (d, 2H).

20

### Example 339

## Synthesis of N-(Methanesulfonyl)-N'-[N-(3,5-Difluorophenylacetyl)-L-alanyl]-L-phenylalanamide

N-Cbz-L-Phenylalanine (Sigma) was coupled to N-hydroxysuccinimide

(Aldrich) using DCC in dicloromethane. The resulting intermediate was reacted with methanesulfonamide in DMF with diisopropylethylamine to provide N-methanesulfonyl-N'-Cbz-L-phenylalanamide amide. The Cbz group was removed using General Procedure O and the resulting intermediate was coupled to N-(3,5-difluorophenylacetyl)-L-alanine (from Example B2 above) using General Procedure B to give the title compound, m.p. = 203-205°C.

## Examples 340-407

By following the procedures set forth above, the following additional compounds were prepared:

	·
5	N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycine methyl ester (Ex 340)
	N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(3- $\alpha$ -phenyl)proline methyl ester (Ex. 341)
	N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-azetidine methyl ester (Ex. 342)
10	methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(5-chlorobenzothiophen-2-yl)acetate (Ex. 343)
	t-butyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-yl)propionate (Ex. 344)
15	t-butyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide (Ex. 345)
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide (Ex. 346)
20	N-[N-(3,4-dichlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 347)
	N-[N-(3-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 348)
	N-[N-(3-bromophenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 349)
25	N-[N-(3-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 350)
	N-[N-(4-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 351)
30	N-[ $N$ -(3-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 352)
	N-[N-(4-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 353)

N-[N-(3-trifluoromethylphenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 354) N-[N-(3-methoxyphenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 355) N-[N-(2-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide 5 (Ex. 356) N-[N-(1-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 357) N-[N-(2-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide 10 (Ex. 358) N-[N-(phenylacetyl)-L-alaninyl]-D-phenylglycinamide (Ex. 359) N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine (Ex. 360)15 N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2furanyl)acetamide (Ex. 361) N'-[N-(3,4-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide (Ex. 362) N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanin-N-20 methylsulfonamide (Ex. 363) N''-methyl-N''-phenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycinamide (Ex. 364) N''-methyl-N''-phenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide (Ex. 365) 25 N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycinamide (Ex. 366) N''-methyl-N''-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycinamide (Ex. 367) N''-4-fluorobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-30 phenylglycinamide (Ex. 368)

- N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-fluoro)phenylglycine neopentyl ester (Ex. 369)
- N-[N-(2,3,4,5,6-pentafluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine methyl ester (Ex. 370)
- 5 N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)serinyl]-L-phenylglycine methyl ester (Ex. 371)
  - N-[N-(3,5-diffuorophenylacetyl)-L-(O-benzyl)threoninyl]-L-phenylglycine methyl ester (Ex. 372)
- N-[N-(3,5-difluorophenylacetyl)-L-threoninyl]-L-phenylglycine methyl ester (Ex. 373)
  - N-[N-(3,5-difluorophenylacetyl)-L-serinyl]-L-phenylglycine methyl ester (Ex. 374)
  - N''-4-methylphenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide (Ex. 375)
- N''-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide (Ex. 376)
  - N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylacetyl
- N'-[N-(3,5-difluorophenylacetyl)-L-methionyl]-L-phenylglycinamide (Ex. 378)
  - N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycinamide (Ex. 379)
  - N'-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinamide (Ex. 380)
- 25 N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycinamide (Ex. 381)
  - N-[(R)- $\alpha$ -methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide (Ex. 382)
- N-[1-phenyl-2-oxo-3-methylbutan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 383)

	N-[1-phenyl-2-oxo-propan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 384)
	N-[1-phenyl-2-oxo-pentan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 385)
5	N-[1-phenyl-2-oxo-2-phenyl-ethan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide (Ex. 386)
	N-[1-phenyl-2-oxo-butan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide (Ex. 387)
10	N-[1-phenyl-2-oxo-4-methylpentan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide (Ex. 388)
	$N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L- $\alpha$ -hydroxyphenylalanine methyl ester (Ex. 389)
	N''-[4-((2-hydroxy-4-azido)-phenyl)-NHC(O)-)butyl] $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide (Ex. 390)
15	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine $t$ -butyl ester (Ex. 391)
	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-phenylphenylglycine $t$ -butyl ester (Ex. 392)
20	[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(2,3-benzo[b]proline) methyl ester (Ex. 393)
·	N''-t-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4- $n$ -butylphenylglycinamide (Ex. 394)
	N''-t-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-(phenylacetenyl)phenylglycinamide (Ex. 395)
25	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide (Ex. 396)
·	N-[1,3-diphenyl-2-oxo-propan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 397)
30	N-[1-phenyl-2-oxo-2-cyclopentylethan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 398)

N-[1-phenyl-2-oxo-hexan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 399)

N-[1-phenyl-2-oxo-3-methylpentan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide (Ex. 400)

5 N''-n-hexyl-6-biotinamidyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide (Ex. 401)

N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-methionine (Ex. 402)

N'-[N-(2-t-BOC-amino)propionyl)-L-alaninyl]-L-phenylglycine methyl ester (Ex. 403)

10 N''-t-butyl N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-fluorophenylglycinamide (Ex. 404)

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-2-phenylglycine methyl ester (Ex. 405)

N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-phenylglycine (Ex. 406)

N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-phenylglycine t-butyl ester (Ex. 407)

### Example 408

Following the procedures set forth above, the following compounds of formula I were or could be prepared:

 $R^1$  is 3,5-difluorophenyl; X' and X'' are hydrogen;  $R^2$  is methyl;  $R^3$  is hydrogen;  $R^4$  is p-fluorophenyl;  $R^5$  is hydrogen; Z is a bond, X is  $-C(O)OCH_2C(CH_3)_3$ ; and n is 1;

R<sup>1</sup> is 3,5-difluorophenyl; X' and X" are hydrogen; R<sup>2</sup> is methyl; R<sup>3</sup> is hydrogen; R<sup>4</sup> is p-(phenyl)phenyl; R<sup>5</sup> is hydrogen; X is -C(O)NHC(CH<sub>3</sub>)<sub>3</sub>; Z is a bond; and n is 1;

 $R^1$  is cyclopentyl; X' and X'' are hydrogen;  $R^2$  is methyl;  $R^3$  is hydrogen;  $R^4$  is phenyl;  $R^5$  is hydrogen; X is  $-C(O)OC(CH_3)_3$ ; X is a bond; and n is 1;

 $R^1$  is cyclopropyl; X' and X" are hydrogen;  $R^2$  is methyl;  $R^3$  is hydrogen;  $R^4$  is phenyl;  $R^5$  is hydrogen; X is  $-C(O)OC(CH_3)_3$ ; Z is a bond; and n is 1; and

 $R^1$  is 3,5-difluorophenyl; X' and X" are hydrogen;  $R^2$  is methyl;  $R^3$  is hydrogen;  $R^4$  is phenyl;  $R^5$  is hydrogen; X is  $-C(O)OCH_2C(CH_3)_3$ ; Z is a bond; and n is 1.

10

15

20

5

### Example 409

## Cellular Screen for the Detection of Inhibitors of $\beta$ -Amyloid Production

Numerous compounds of formula I above were assayed for their ability to inhibit  $\beta$ -amyloid production in a cell line possessing the Swedish mutation. This screening assay employed cells (K293 = human kidney cell line) which were stably transfected with the gene for amyloid precursor protein 751 (APP751) containing the double mutation Lys<sub>651</sub>Met<sub>652</sub> to Asn<sub>651</sub>Leu<sub>652</sub> (APP751 numbering) in the manner described in International Patent Application Publication No. 94/10569<sup>8</sup> and Citron et al. 12. This mutation is commonly called the Swedish mutation and the cells, designated as "293 751 SWE", were plated in Corning 96-well plates at 1.5-2.5 x  $10^4$  cells per well in Dulbecco's minimal essential media plus 10% fetal bovine serum. Cell number is important in order to achieve  $\beta$ -amyloid ELISA results within the linear range of the assay (~0.2 to 2.5 ng per mL).

Following overnight incubation at 37°C in an incubator equilibrated with 10% carbon dioxide, media were removed and replaced with 200 µL of a compound of formula I (drug) containing media per well for a two hour pretreatment period and cells were incubated as above. Drug stocks were prepared in 100% dimethylsulfoxide such that at the final drug concentration

10

15

20

used in the treatment, the concentration of dimethylsulfoxide did not exceed 0.5% and, in fact, usually equaled 0.1%.

At the end of the pretreatment period, the media were again removed and replaced with fresh drug containing media as above and cells were incubated for an additional two hours. After treatment, plates were centrifuged in a Beckman GPR at 1200 rpm for five minutes at room temperature to pellet cellular debris from the conditioned media. From each well,  $100 \mu L$  of conditioned media or appropriate dilutions thereof were transferred into an ELISA plate precoated with antibody  $266^{14}$  against amino acids 13-28 of  $\beta$ -amyloid peptide as described in International Patent Application Publication No.  $94/10569^8$  and stored at  $4^{\circ}C$  overnight. An ELISA assay employing labelled antibody  $6C6^{14}$  against amino acids 1-16 of  $\beta$ -amyloid peptide was run the next day to measure the amount of  $\beta$ -amyloid peptide produced.

Cytotoxic effects of the compounds were measured by a modification of the method of Hansen, et al. 13. To the cells remaining in the tissue culture plate was added 25  $\mu$ L of a 3,(4,5-dimethylthiazol-2-yl)2,5-diphenyltetrazolium bromide (MTT) stock solution (5 mg/mL) to a final concentration of 1 mg/mL. Cells were incubated at 37°C for one hour, and cellular activity was stopped by the addition of an equal volume of MTT lysis buffer (20% w/v sodium dodecylsulfate in 50% dimethylformamide, pH 4.7). Complete extraction was achieved by overnight shaking at room temperature. The difference in the OD<sub>562nun</sub> and the OD<sub>650nm</sub> was measured in a Molecular Device's UV<sub>max</sub> microplate reader as an indicator of the cellular viability.

The results of the  $\beta$ -amyloid peptide ELISA were fit to a standard curve and expressed as ng/mL  $\beta$ -amyloid peptide. In order to normalize for cytotoxicity, these results were divided by the MTT results and expressed as a percentage of the results from a drug free control. All results are the mean and standard deviation of at least six replicate assays.

The test compounds were assayed for  $\beta$ -amyloid peptide production inhibition activity in cells using this assay. The results of this assay demonstrate that, each of the compounds within this invention tested reduced  $\beta$ -amyloid peptide production by at least 30% as compared to control.

5

10

15

20

## Example 410

## In Vivo Suppression of $\beta$ -Amyloid Release and/or Synthesis

This example illustrates how the compounds of this invention could be tested for *in vivo* suppression of  $\beta$ -amyloid release and/or synthesis. For these experiments, 3 to 4 month old PDAPP mice are used [Games et al., (1995) Nature 373:523-527]. Depending upon which compound is being tested, the compound is usually formulated at either 5 or 10 mg/ml. Because of the low solubility factors of the compounds, they may be formulated with various vehicles, such as corn oil (Safeway, South San Francisco, CA); 10% EtOH in corn oil (Safeway); 2-hydroxypropyl- $\beta$ -cyclodextrin (Research Biochemicals International, Natick MA); and carboxy-methyl-cellulose (Sigma Chemical Co., St. Louis MO). Specifically, for example 141 the vehicle was carboxy-methyl-cellulose (Sigma).

The mice are dosed subcutaneously with a 26 gauge needle and 3 hours later the animals are euthanized via CO<sub>2</sub> narcosis and blood is taken by cardiac puncture using a 1 cc 25G 5/8" tuberculin syringe/needle coated with solution of 0.5 M EDTA, pH 8.0. The blood is placed in a Becton-Dickinson vacutainer tube containing EDTA and spun down for 15 minutes at 1500 xg at 5°C. The brains of the mice are then removed and the cortex and hippocampus are dissected out and placed on ice.

## 25 1. Brain Assay

To prepare hippocampal and cortical tissue for enzyme-linked immunosorbent assays (ELISAs) each brain region is homogenized in 10

volumes of ice cold guanidine buffer (5.0 M guanidine-HCl, 50 mM Tris-HCl, pH 8.0) using a Kontes motorized pestle (Fisher, Pittsburgh PA). The homogenates are gently rocked on a rotating platform for three to four hours at room temperature and stored at -20°C prior to quantitation of  $\beta$ -amyloid.

5

The brain homogenates are diluted 1:10 with ice-cold casein buffer [0.25% casein, phosphate buffered saline (PBS), 0.05% sodium azide, 20  $\mu$ g/ml aprotinin, 5 mM EDTA, pH 8.0, 10  $\mu$ g/ml leupeptin], thereby reducing the final concentration of guanidine to 0.5 M, before centrifugation at 16,000 xg for 20 minutes at 4°C. The  $\beta$ -amyloid standards (1-40 or 1-42 amino acids) were prepared such that the final composition equaled 0.5 M guanidine in the presence of 0.1% bovine serum albumin (BSA).

10

15

20

The total  $\beta$ -amyloid sandwich ELISA, quantitating both  $\beta$ -amyloid (aa 1-40) and  $\beta$ -amyloid (aa 1-42) consists of two monoclonal antibodies (mAb) to  $\beta$ -amyloid. The capture antibody,  $266^{14}$ , is specific to amino acids 13 - 28 of  $\beta$ -amyloid. The antibody  $3D6^{15}$ , which is specific to amino acids 1 - 5 of  $\beta$ -amyloid, is biotinylated and served as the reporter antibody in the assay. The 3D6 biotinylation procedure employs the manufacturer's (Pierce, Rockford IL) protocol for NHS-biotin labeling of immunoglobulins except that 100 mM sodium bicarbonate, pH 8.5 buffer is used. The 3D6 antibody does not recognize secreted amyloid precursor protein (APP) or full-length APP but detects only  $\beta$ -amyloid species with an amino terminal aspartic acid. The assay has a lower limit of sensitivity of  $\sim 50$  pg/ml (11 pM) and shows no crossreactivity to the endogenous murine  $\beta$ -amyloid peptide at concentrations up to 1 ng/ml.

25

The configuration of the sandwich ELISA quantitating the level of  $\beta$ -amyloid (aa 1-42) employs the mAb 21F12<sup>15</sup> (which recognizes amino acids 33-42 of  $\beta$ -amyloid) as the capture antibody. Biotinylated 3D6 is also the reporter

10

15

25

antibody in this assay which has a lower limit of sensitivity of  $\sim 125$  pg/ml (28 pM).

The 266 and 21F12 capture mAbs are coated at 10  $\mu$ g/ml into 96 well immunoassay plates (Costar, Cambidge MA) overnight at room temperature. The plates are then aspirated and blocked with 0.25% human serum albumin in PBS buffer for at least 1 hour at room temperature, then stored desiccated at 4°C until use. The plates are rehydrated with wash buffer (Tris-buffered saline, 0.05% Tween 20) prior to use. The samples and standards are added to the plates and incubated overnight at 4°C. The plates are washed ≥ 3 times with wash buffer between each step of the assay. The biotinylated 3D6, diluted to 0.5  $\mu$ g/ml in casein incubation buffer (0.25% casein, PBS, 0.05% Tween 20, pH 7.4) is incubated in the well for 1 hour at room temperature. Avidin-HRP (Vector, Burlingame CA) diluted 1:4000 in casein incubation buffer is added to the wells for 1 hour at room temperature. The colorimetric substrate, Slow TMB-ELISA (Pierce, Cambridge MA), is added and allowed to react for 15 minutes, after which the enzymatic reaction is stopped with addition of 2 N H<sub>2</sub>SO<sub>4</sub>. Reaction product is quantified using a Molecular Devices Vmax (Molecular Devices, Menlo Park CA) measuring the difference in absorbance at 450 nm and 650 nm.

## 20 2. Blood Assay

The EDTA plasma is diluted 1:1 in specimen diluent (0.2 gm/l sodium phosphate• $H_2O$  (monobasic), 2.16 gm/l sodium phosphate• $7H_2O$  (dibasic), 0.5gm/l thimerosal, 8.5 gm/l sodium chloride, 0.5 ml TritonX-405, 6.0 g/l globulin-free bovine serum albumin; and water). The samples and standards in specimen diluent are assayed using the total  $\beta$ -amyloid assay (266 capture/3D6 reporter) described above for the brain assay except the specimen diluent was used instead of the casein diluents described.

From the foregoing description, various modifications and changes in the composition and method will occur to those skilled in the art. All such modifications coming within the scope of the appended claims are intended to be included therein.

15

20

25

30

#### WHAT IS CLAIMED IS:

1. A method for inhibiting  $\beta$ -amyloid peptide release and/or its synthesis in a cell which method comprises administering to such a cell an amount of a compound or a mixture of compounds effective in inhibiting the cellular release and/or synthesis of  $\beta$ -amyloid peptide wherein said compounds are represented by formula I:

10 I 
$$R^1Z$$
  $X'$   $X''$   $X''$ 

wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

each R<sup>3</sup> is independently selected from the group consisting of hydrogen and methyl and R<sup>3</sup> together with R<sup>4</sup> can be fused to form a cyclic structure of from 3 to 8 atoms which is optionally fused with an aryl or heteroaryl group;

each R<sup>4</sup> is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclic, substituted alkyl, substituted alkenyl and substituted alkynyl;

each R<sup>5</sup> is selected from hydrogen and methyl or together with R<sup>4</sup> forms a cycloalkyl group of from 3 to 6 carbon atoms;

X is selected from the group consisting of -C(O)Y and -C(S)Y where Y is selected from the group consisting of

- (a) alkyl or cycloalkyl,
- (b) substituted alkyl with the proviso that the substitution on said substituted alkyl do not include  $\alpha$ -haloalkyl,  $\alpha$ -diazoalkyl,  $\alpha$ -OC(O)alkyl, or

 $\alpha$ -OC(O)aryl groups,

- (c) alkoxy or thioalkoxy,
- (d) substituted alkoxy or substituted thioalkoxy,
- (e) hydroxy,
- 5 (f) aryl,

10

15

20

25

- (g) heteroaryl,
- (h) heterocyclic,
- (i) -NR'R" where R' and R" are independently selected from hydrogen, alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, where one of R' or R" is hydroxy or alkoxy, and where R' and R" are joined to form a cyclic group having from 2 to 8 carbon atoms optionally containing 1 to 2 additional heteroatoms selected from oxygen, sulfur and nitrogen and optionally substituted with one or more alkyl, alkoxy or carboxylalkyl groups,
- (j) -NHSO<sub>2</sub>-R<sup>8</sup> where R<sup>8</sup> is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic,
- (k) -NR<sup>9</sup>NR<sup>10</sup>R<sup>10</sup> where R<sup>9</sup> is hydrogen or alkyl, and each R<sup>10</sup> is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, and
- (1)  $-ONR^9[C(O)O]_zR^{10}$  where z is zero or one,  $R^9$  and  $R^{10}$  are as defined above;

X can also be -CR<sup>6</sup>R<sup>6</sup>Y' where each R<sup>6</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic and Y' is selected from the group consisting of hydroxyl, amino, thiol, alkoxy, substituted alkoxy, thioalkoxy, substituted thioalkoxy, -OC(O)R<sup>7</sup>, -SSR<sup>7</sup>, -SSC(O)R<sup>7</sup> where R<sup>7</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

X' is hydrogen, hydroxy, or fluoro;

X" is hydrogen, hydroxy or fluoro, or X' and X" together form an oxo group,

10

20

30

Z is selected from the group consisting of a bond covalently linking  $R^1$  to -CX'X"-, oxygen and sulfur;

n is an integer equal to 1 or 2; and pharmaceutically acceptable salts thereof with the provisos that:

A. when  $R^1$  is phenyl or 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH;

B. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is  $-CH(OH)CH_3$  derived from D-threonine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

C. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^4$  is benzyl,  $R^5$  is hydrogen, X is methoxycarbonyl, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$  is not methyl;

D. when  $R^1$  is *iso*-propyl,  $R^2$  is  $-CH_2C(O)NH_2$ ,  $R^3$  is hydrogen,  $R^4$  is *iso*-butyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

E. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^5$  is hydrogen, X is -C(O)OCH<sub>3</sub>, X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^3$ , the nitrogen atom attached to  $R^3$ , and  $R^4$  do not form 1,2,3,4-tetrahydroiso-quinolin-2-yl or pyrrolidin-2-yl;

F. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not 4-amino-n-butyl;

G. when  $R^1$  is 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH<sub>2</sub> or -CH<sub>2</sub>OH;

H. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-CH_2OCH_3$ , X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not benzyl or ethyl;

10

15

20

25

30

I

I. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is methyl,  $R^4$  is methyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$ ;

J. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$  or -CH<sub>2</sub>OH;

K. when  $R_1$  is N-(2-pyrrolidinonyl),  $R_2$  is methyl,  $R_3$  is hydrogen,  $R_4$  is benzyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OCH<sub>3</sub>;

L. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl derived from D-alanine,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH-benzyl;

M. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is hydrogen,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CH<sub>2</sub>OH;

N. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is 4-phenylphenyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NHC(CH<sub>3</sub>)<sub>3</sub>; and

O. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NHCH(CH<sub>3</sub>) $\phi$ .

2. A method for preventing the onset of AD in a patient at risk for developing AD which method comprises administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I:

$$R^{1}Z$$
 $X'$ 
 $X''$ 
 $X''$ 
 $R^{2}$ 
 $R^{3}$ 
 $R^{4}$ 
 $R^{5}$ 
 $R^{5}$ 

10

15

20

30

wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

each R<sup>3</sup> is independently selected from the group consisting of hydrogen and methyl and R<sup>3</sup> together with R<sup>4</sup> can be fused to form a cyclic structure of from 3 to 8 atoms which is optionally fused with an aryl or heteroaryl group;

each R<sup>4</sup> is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclic, substituted alkyl, substituted alkyl, substituted alkynyl;

each  $R^5$  is selected from hydrogen and methyl or together with  $R^4$  forms a cycloalkyl group of from 3 to 6 carbon atoms;

X is selected from the group consisting of -C(O)Y and -C(S)Y where Y is selected from the group consisting of

- (a) alkyl or cycloalkyl,
- (b) substituted alkyl with the proviso that the substitution on said substituted alkyl do not include  $\alpha$ -haloalkyl,  $\alpha$ -diazoalkyl,  $\alpha$ -OC(O)alkyl, or  $\alpha$ -OC(O)aryl groups,
  - (c) alkoxy or thioalkoxy,
  - (d) substituted alkoxy or substituted thioalkoxy,
  - (e) hydroxy,
  - (f) aryl,
- 25 (g) heteroaryl,
  - (h) heterocyclic,
  - (i) -NR'R" where R' and R" are independently selected from hydrogen, alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, where one of R' or R" is hydroxy or alkoxy, and where R' and R" are joined to form a cyclic group having from 2 to 8 carbon atoms optionally containing 1 to 2 additional

20

30

heteroatoms selected from oxygen, sulfur and nitrogen and optionally substituted with one or more alkyl, alkoxy or carboxylalkyl groups,

- (j) -NHSO<sub>2</sub>-R<sup>8</sup> where R<sup>8</sup> is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic,
- (k) -NR<sup>9</sup>NR<sup>10</sup>R<sup>10</sup> where R<sup>9</sup> is hydrogen or alkyl, and each R<sup>10</sup> is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, and
- (1)  $-ONR^9[C(O)O]_zR^{10}$  where z is zero or one,  $R^9$  and  $R^{10}$  are as defined above;

X can also be -CR<sup>6</sup>R<sup>6</sup>Y' where each R<sup>6</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic and Y' is selected from the group consisting of hydroxyl, amino, thiol, alkoxy, substituted alkoxy, thioalkoxy, substituted thioalkoxy, -OC(O)R<sup>7</sup>, -SSR<sup>7</sup>, -SSC(O)R<sup>7</sup> where R<sup>7</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

X' is hydrogen, hydroxy, or fluoro;

X'' is hydrogen, hydroxy or fluoro, or X' and X'' together form an oxo group,

Z is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X"-, oxygen and sulfur;

n is an integer equal to 1 or 2; and pharmaceutically acceptable salts thereof with the provisos that:

A. when  $R^1$  is phenyl or 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH;

B. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub> derived from D-threonine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH or -C(O)OCH<sub>3</sub>;

10

15

25

C. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^4$  is benzyl,  $R^5$  is hydrogen, X is methoxycarbonyl, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$  is not methyl;

D. when  $R^1$  is iso-propyl,  $R^2$  is  $-CH_2C(O)NH_2$ ,  $R^3$  is hydrogen,  $R^4$  is iso-butyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

E. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^3$ , the nitrogen atom attached to  $R^3$ , and  $R^4$  do not form 1,2,3,4-tetrahydroiso-quinolin-2-yl or pyrrolidin-2-yl;

F. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not 4-amino-n-butyl;

G. when  $R^1$  is 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH<sub>2</sub> or -CH<sub>2</sub>OH;

H. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-CH_2OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not benzyl or ethyl;

- I. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is methyl,  $R^4$  is methyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$ ;
  - J. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$  or -CH<sub>2</sub>OH;

K. when  $R_1$  is N-(2-pyrrolidinonyl),  $R_2$  is methyl,  $R_3$  is hydrogen,  $R_4$  is benzyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OCH<sub>3</sub>;

L. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl derived from D-alanine, R<sup>3</sup> is hydrogen, R<sup>4</sup> is phenyl derived from D-phenylglycine, R<sup>5</sup> is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH-benzyl;

25

30

M. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is hydrogen,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CH<sub>2</sub>OH;

N. when  $R^1$  is 3,5-diffuorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is 4-phenylphenyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHC(CH_3)_3$ ; and

O. when  $R^1$  is 3,5-diffuorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHCH(CH_3)\phi$ .

3. A method for treating a patient with AD in order to inhibit further deterioration in the condition of that patient which method comprises administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I:

15
$$R^{1}Z$$

$$R^{2}$$

$$R^{3}$$

wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

each R<sup>3</sup> is independently selected from the group consisting of hydrogen and methyl and R<sup>3</sup> together with R<sup>4</sup> can be fused to form a cyclic structure of from 3 to 8 atoms which is optionally fused with an aryl or heteroaryl group;

10

20

25

30

each R<sup>4</sup> is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclic, substituted alkyl, substituted alkenyl and substituted alkynyl;

each R<sup>5</sup> is selected from hydrogen and methyl or together with R<sup>4</sup> forms a cycloalkyl group of from 3 to 6 carbon atoms;

X is selected from the group consisting of -C(O)Y and -C(S)Y where Y is selected from the group consisting of

- (a) alkyl or cycloalkyl,
- (b) substituted alkyl with the proviso that the substitution on said substituted alkyl do not include  $\alpha$ -haloalkyl,  $\alpha$ -diazoalkyl,  $\alpha$ -OC(O)alkyl, or  $\alpha$ -OC(O)aryl groups,
  - (c) alkoxy or thioalkoxy,
  - (d) substituted alkoxy or substituted thioalkoxy,
  - (e) hydroxy,
- 15 (f) aryl,
  - (g) heteroaryl,
  - (h) heterocyclic,
  - (i) -NR'R" where R' and R" are independently selected from hydrogen, alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, where one of R' or R" is hydroxy or alkoxy, and where R' and R" are joined to form a cyclic group having from 2 to 8 carbon atoms optionally containing 1 to 2 additional heteroatoms selected from oxygen, sulfur and nitrogen and optionally substituted with one or more alkyl, alkoxy or carboxylalkyl groups,
  - (j) -NHSO<sub>2</sub>-R<sup>8</sup> where R<sup>8</sup> is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic,
  - (k) -NR<sup>9</sup>NR<sup>10</sup>R<sup>10</sup> where R<sup>9</sup> is hydrogen or alkyl, and each R<sup>10</sup> is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, and
  - (1)  $-ONR^9[C(O)O]_zR^{10}$  where z is zero or one,  $R^9$  and  $R^{10}$  are as defined above;

15

20

30

X can also be -CR<sup>6</sup>R<sup>6</sup>Y' where each R<sup>6</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic and Y' is selected from the group consisting of hydroxyl, amino, thiol, alkoxy, substituted alkoxy, thioalkoxy, substituted thioalkoxy, -OC(O)R<sup>7</sup>, -SSR<sup>7</sup>, -SSC(O)R<sup>7</sup> where R<sup>7</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

X' is hydrogen, hydroxy, or fluoro;

X'' is hydrogen, hydroxy or fluoro, or X' and X'' together form an oxo group,

Z is selected from the group consisting of a bond covalently linking  $R^1$  to -CX'X"-, oxygen and sulfur;

n is an integer equal to 1 or 2; and pharmaceutically acceptable salts thereof with the provisos that:

A. when  $R^1$  is phenyl or 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH;

B. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub> derived from D-threonine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH or -C(O)OCH<sub>3</sub>;

C. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^4$  is benzyl,  $R^5$  is hydrogen, X is methoxycarbonyl, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$  is not methyl;

D. when  $R^1$  is iso-propyl,  $R^2$  is  $-CH_2C(O)NH_2$ ,  $R^3$  is hydrogen,  $R^4$  is iso-butyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

E. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^3$ , the nitrogen atom attached to  $R^3$ , and  $R^4$  do not form 1,2,3,4-tetrahydroiso-quinolin-2-yl or pyrrolidin-2-yl;

15

20

25

30

F. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not 4-amino-n-butyl;

G. when  $R^1$  is 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH<sub>2</sub> or -CH<sub>2</sub>OH;

H. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-CH_2OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not benzyl or ethyl;

I. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is methyl,  $R^4$  is methyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$ ;

J. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$  or -CH<sub>2</sub>OH;

K. when  $R_1$  is N-(2-pyrrolidinonyl),  $R_2$  is methyl,  $R_3$  is hydrogen,  $R_4$  is benzyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OCH<sub>3</sub>;

L. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl derived from D-alanine,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH-benzyl;

M. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is hydrogen,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CH<sub>2</sub>OH;

N. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is 4-phenylphenyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHC(CH_3)_3$ ; and

O. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHCH(CH_3)\phi$ .

- 4. The method according to Claim 1, 2 or 3 wherein  $R^1$  is an unsubstituted aryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 5. The method according to Claim 4 wherein the unsubstituted R<sup>1</sup> aryl group is selected from the group consisting of phenyl, 1-naphthyl and 2-naphthyl.
- 6. The method according to Claim 1, 2 or 3 wherein  $R^1$  is a substituted aryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 7. The method according to Claim 6 wherein said substituted aryl group is a mono-substituted, di-substituted or tri-substituted phenyl group.
- 8. The method according to Claim 7 wherein the substituted phenyl groups are selected from the group consisting of 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 3-methoxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2-hydroxyphenyl, 2-methylphenyl, 2-fluorophenyl, 2-chlorophenyl, 3,4-difluorophenyl, 2,3,4,5,6-pentafluorophenyl, 3,4-dibromophenyl, 3,4-dichlorophenyl, 3,4-methylene-dioxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 2,4-dichlorophenyl, and 2,5-difluorophenyl.
- 9. The method according to Claim 1, 2 or 3 wherein R<sup>1</sup> is an alkaryl group and Z is a bond covalently linking R<sup>1</sup> to -CX'X"-.
  - 10. The method according to Claim 9 wherein the R<sup>1</sup> alkaryl group is selected from the group consisting of benzyl, 2-phenylethyl, and 3-phenyl-n-propyl.

- 11. The method according to Claim 1, 2 or 3 wherein  $R^1$  is selected from the group consisting of alkyl, alkenyl, cycloalkyl and cycloalkenyl groups and Z is a bond covalently linking  $R^1$  to -CX'X''-.
  - 12. The method according to Claim 11 wherein R1 is alkyl.
- 13. The method according to Claim 11 wherein R<sup>1</sup> is cycloalkyl.
  - 14. The method according to Claim 11 wherein R1 is alkenyl.
  - 15. The method according to Claim 11 wherein R1 is cycloalkenyl.
- 16. The method according to Claim 11 wherein the R¹ alkyl, cycloalkyl, alkenyl and cycloalkenyl groups are selected from the group consisting of *iso*-propyl, *n*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, aminomethyl, and N-tert-butoxycarbonylaminomethyl.
  - 17. The method according to Claim 1, 2 or 3 wherein  $R^1$  is selected from the group consisting of heteroaryl and substituted heteroaryl groups and Z is a bond covalently linking  $R^1$  to -CX'X"-.
- substituted heteroaryl groups are selected from the group consisting of pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, fluoropyridyls (including 5-fluoropyrid-3-yl), chloropyridyls (including 5-chloropyrid-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-

10

15

(thiophenyl)thiophen-5-yl, 6-methoxythionaphthen-2-yl, 3-phenyl-1,2,4-thiooxadiazol-5-yl and 2-phenyloxazol-4-yl.

- 19. The method according to Claim 1, 2 or 3 wherein R<sup>2</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic.
- 20. The method according to Claim 9 wherein R<sup>2</sup> is selected from the group consisting of methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, phenyl, 4-fluorophenyl, 3,5-difluoro-phenyl, 4-methoxyphenyl, benzyl, cyclopropyl, cyclohexyl, cyclopentyl, cycloheptyl, thien-2-yl, thien-3-yl, -CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>, -CH<sub>2</sub>OCH<sub>2</sub>φ, -CH(CH<sub>3</sub>)OCH<sub>2</sub>φ, -CH(OH)CH<sub>3</sub> and -CH<sub>2</sub>OH.
- 21. The method according to Claim 1, 2 or 3 wherein X' and X'' are hydrogen and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 22. The method according to Claim 21 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, methyl or together with R<sup>4</sup> and the nitrogen to which R<sup>3</sup> is attached forms pyrrolidin-2-yl, 2,3-dihydroindol-2-yl, piperidin-2-yl, 4-hydroxy-pyrrolidin-2-yl and 1,2,3,4-tetrahydroisoquinolin-3-yl.
- 23. The method according to Claim 1, 2 or 3 wherein R<sup>4</sup> substituents are selected from the group consisting of hydrogen, methyl, ethyl, iso-propyl, n-propyl, n-butyl, sec-butyl, iso-butyl, cyclopentyl, cyclohexyl, allyl, iso-but-2-enyl, 3-methylpentyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-indol-3-yl, phenyl, p-(phenyl)phenyl, m-(phenyl)phenyl o-fluorophenyl, m-fluorophenyl, p-fluorophenyl, p-bromophenyl, m-methoxyphenyl, p-methoxyphenyl, phenethyl, benzyl, m-hydroxybenzyl, p-hydroxybenzyl, p-nitrobenzyl, m-trifluoromethylphenyl, p-(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl,
  25 p-(CH<sub>3</sub>)<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, p-phenylphenyl, 3,5-difluorophenyl,

p-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyrid-6-yl, 4-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)benzyl, -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>, -CH<sub>2</sub>-imidazol-4-yl, -CH<sub>2</sub>-(3-tetrahydrofuranyl), -CH<sub>2</sub>-thien-2-yl, -CH<sub>2</sub>-thiazol-4-yl, -CH<sub>2</sub>(1-methyl)cyclopropyl, -CH<sub>2</sub>-thien-3-yl, thien-3-yl, thien-2-yl,  $-CH_2-C(O)O-t$ -butyl,  $-CH_2-C(CH_3)_3$ ,  $-CH_2CH(CH_2CH_3)_2$ , 5 2-methylcyclopentyl, -cyclohex-2-enyl, -CH[CH(CH<sub>3</sub>)<sub>2</sub>]COOCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>SCH<sub>3</sub>,  $-CH_2CH_2N(CH_3)_2$ ,  $-CH_2C(CH_3)=CH_2$ ,  $-CH_2CH=CHCH_3$  (cis and trans), -CH<sub>2</sub>OH, -CH(OH)CH<sub>3</sub>, -CH(O-t-butyl)CH<sub>3</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>NH-Boc, -(CH<sub>2</sub>)<sub>4</sub>NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>4</sub>N(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-pyridyl, pyridyl, -CH<sub>2</sub>-naphthyl, -CH<sub>2</sub>-(N-morpholino), p-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-10 yl, benzo[b]thiophen-3-yl, 5-chlorobenzo[b]thiophen-2-yl, 4,5,6,7tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, tetrazol-5-yl, 5-chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl, -CH<sub>2</sub>-N-phthalimidyl, 2-methylthiazol-4-yl, and thieno[2,3-b]thiophen-2-yl, 5-bromothien-2-yl, 4-bromothien-2-yl, 5-chlorothien-2-yl, 3-phenoxyphenyl, 15 2-phenoxyphenyl, 4-ethylphenyl, 2-benzylphenyl, (4-ethylphenyl)phenyl, 4-tertbutylphenyl, 4-n-butylphenyl, o-(4-chlorophenoxy)phenyl, furan-2-yl, and 4phenylacetylenylphenyl.

- 24. The method according to Claim 1, 2 or 3 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"- and R<sup>4</sup> and R<sup>5</sup> are fused to form a cycloalkyl group selected from the group consisting of cyclopropyl and cyclobutyl.
- 25. The method according to Claims 1, 2 or 3 wherein Z is a covalent bond linking  $R^1$  to -CX'X''-, X is -C(O)Y and Y is selected from the group consisting of hydroxy, alkoxy or substituted alkoxy.
- 26. The method according to Claim 25 wherein Y is alkoxy or substituted alkoxy selected from the group consisting of methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, tert-butoxy, neo-pentoxy, benzyloxy, 2-phenylethoxy, 3-phenyl-n-propoxy, 3-iodo-n-propoxy, 4-bromo-n-butoxy, -ONHC(O)OC(CH<sub>3</sub>)<sub>3</sub>, -ONHC(CH<sub>3</sub>)<sub>3</sub> and hydroxy.

- 27. The method according to Claims 1, 2 or 3 wherein Z is a covalent bond linking  $R^1$  to -CX'X''-, X is -C(O)Y and Y is -NR'R''.
- The method according to Claim 27 wherein Y is selected from 28. the group consisting of amino (-NH2), -NH(iso-butyl), -NH(sec-butyl), Nmethylamino, N,N-dimethylamino, N-benzylamino, N-morpholino, azetidino, 5 N-thiomorpholino, N-piperidinyl, N-hexamethyleneimino, N-heptamethyleneimino, N-pyrrolidinyl, -NH-methallyl, -NHCH2-(furan-2-yl), -NHCH2cyclopropyl, -NH(tert-butyl), -NH(p-methylphenyl), -NHOCH<sub>3</sub>, -NHCH<sub>2</sub>(pfluorophenyl), -NHCH2CH2OCH3, -NH-cyclopentyl, -NH-cyclohexyl, -NHCH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -NHCH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>, -NHCH<sub>2</sub>-(pyrid-2-yl), -NHCH<sub>2</sub>-(pyrid-10 3-yl), -NHCH<sub>2</sub>-(pyrid-4-yl), N-thiazolindinyl, -N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -N[CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -NHOH, -NH(p-NO<sub>2</sub>- $\phi$ ), -NHCH<sub>2</sub>(p-NO<sub>2</sub>- $\phi$ ), -NHCH<sub>2</sub>(m- $NO_2-\phi$ ),  $-N(CH_3)OCH_3$ ,  $-N(CH_3)CH_2-\phi$ ,  $-NHCH_2-(3,5-di-fluorophenyl)$ , -NHCH<sub>2</sub>CH<sub>2</sub>F, -NHCH<sub>2</sub>(p-CH<sub>3</sub>O- $\phi$ ), -NHCH<sub>2</sub>(m-CH<sub>3</sub>O- $\phi$ ), -NHCH<sub>2</sub>(p-CF<sub>3</sub>- $\phi$ ), -N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>2</sub> $\phi$ , -NHCH(CH<sub>3</sub>) $\phi$ , -NHCH<sub>2</sub>-(p-F- $\phi$ ), 15 -N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -NHCH<sub>2</sub>-(tetrahydrofuran-2-yl), -NHCH<sub>2</sub>(ptrifluoromethylphenyl),  $-NHCH_2C(CH_3) = CH_2$ , -NH-[(p-benzyl)pyrid-4-yl], -NH-[(2,6-dimethyl)pyrid-4-yl], -NH-(2-methylcyclohexyl), -NH-(4methylcyclohexyl), -NH-[N-ethoxycarbonyl]-piperidin-4-yl, -NHOC(CH<sub>3</sub>)<sub>3</sub>,  $-NHCH_2CH_2CH_2CH_2-\phi$ ,  $-C(O)NH(CH_2)_3O-(p-CH_3)\phi$ ,  $-C(O)NH(CH_2)_6NH_2$ , 20 -NH-(tetrahydrofuran-2-yl), -N(CH<sub>3</sub>) $\phi$ , -NH(CH<sub>2</sub>)<sub>4</sub>NHC(O)-(2-hydroxy-4azido)-phenyl and -NH(CH<sub>2</sub>)<sub>6</sub>-(biotinamidyl).
- 29. The method according to Claims 1, 2 or 3 wherein X is -C(O)Y and Y is selected from the group consisting of -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>,
  25 -CH<sub>2</sub>OH, -CH(OH)CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(OH)φ, -CH(OH)CH<sub>2</sub>C(O)OCH<sub>3</sub>,
  -C(OH)(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>OC(O)OCH<sub>3</sub>, and -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>,
  methyl, ethyl, iso-propyl, n-propyl, iso-butyl, n-butyl, sec-butyl, tent-butyl,
  -CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-pyridy-2-yl, -CH<sub>2</sub>-pyridy-3-yl, -CH<sub>2</sub>-pyridy-4-yl,
  -CH<sub>2</sub>-fur-2-yl, benzyl, cyclopentyl, phenyl, and -NH-SO<sub>2</sub>-CH<sub>3</sub>.

- 30. The method according to Claims 1, 2 or 3 wherein Z is a covalent bond linking  $R^1$  to -CX'X''-.
- 31. The method according to Claims 1, 2 or 3 wherein the compound of formula I is selected from the group consisting of:
- 5 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-histidine methyl ester

N-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

10 N-2-(N,N-dimethylamino) ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

N-2-(N,N-dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

15

N-(4-pyridyl) methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

N-(3-pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

20 N-(4-pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate *tert*-butyl ester

N-[N-(pent-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester

25 N-[N-(dec-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[3-(N,N-dimethylamino)propoxy]phenylalanine methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[(tert-butyloxycarbonyl)methoxy]phenylalanine methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(carboxymethoxy)phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-5 morpholinoethoxy)phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-6-(N,Ndimethylamino)hexanoate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2pyridyl)propionate methyl ester 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(3pyridyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-proline methyl ester 1-[N-(3,5-difluorophenylacetyl)-L-alaninyl]piperidine-2-carboxylate methyl ester 15 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4pyridyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3methoxypropionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-20 morpholinopropionate methyl ester N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2morpholinoethoxy)phenylalaninamide N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2amino-3-methoxypropionamide 25 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine methyl ester N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2amino-3-(4-pyridyl)propionamide N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2amino ·3-(2-pyridyl)propionamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4yl)propionate methyl ester 2-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1,2,3,4tetrahydroisoquinoline-3-carboxylate methyl ester N-(3-methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 phenylalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(1naphthyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-10 naphthyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2thienyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine benzyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-bromo-15 propyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3iodopropyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine tert-butyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-20 pyridyl)acetamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3pyridyl)acetamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-N,-(tert-butoxycarbonyl)-Llysine methyl ester 25 methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-4phenylbutanoate N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 2-phenylethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 3-phenylpropyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-30 pyridyl)acetamide

N-[N-(phenylacetyl)-L-alaninyl]-L-threonine methyl ester N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N'-[N-(phenylacetyl)-L-alaninyl]-L-alaninamide N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide 5 N'-[N-(phenylacetyl)-L-alaninyl)-L-valinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetate ethyl ester N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N,N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide 10 N, N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N, N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-15 aminohexanamide N, N-dimethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminohexanamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-20 methoxyphenyl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4methoxyphenyl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetate ethyl ester 25 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetate ethyl ester N-[N-(cyclohexylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(cyclohex-1-enylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1-aminocyclopropane-1carboxylate methyl ester 5 N-2-(N,N-dimethylamino) ethyl-N-methyl-N'-[N-(3,5difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine benzyl ester N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester 10 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-alanine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]glycine ethyl ester N-hydroxy-N'-[N-(3-nitrophenylacetyl)-L-alaninyl]-D,L-threoninamide N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester 15 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-2-amino-3-(3hydroxyphenyl)propionate methyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-tyrosine ethyl ester N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester N-[N-[N-(isovaleryl)-L-valinyl]-L-phenylglycinyl]-L-alanine iso-butyl 20 ester N-[N-(isovaleryl)-L-phenylalaninyl]-L-alanine iso-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester 1-[N-(3-nitrophenylacetyl)-L-alaninyl]-indoline-(S)-2-carboxylate ethyl ester 25 N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-methoxy-N-methyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide

N-iso-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N, N-di-n-propyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valinamide N-(4-nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 alaninamide N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninyl]-L-phenylalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide 10 N-iso-butyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide N-(4-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 15 N-(4-nitrophenyl)-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninyll]-L-alanialaninamide N-(4-nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide N-benzyl-N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 alaninamide N-(3,5-difluorobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-(3-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 25 N-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-(4-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophan methyl ester

N-(4-methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[N-(phenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninyl]-L-5 phenylglycine methyl ester N-[N-(cyclohexylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine methyl ester N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl]-L-alaninyl[-alaninyl[-alaninyl]-L-alaninyl[-alaninyl[-alaninyl]-L-alaninyl[-alaninyl[-alaninyl]-L-alaninyl[-alaninyl[-alaninyl[-alaninyl[-alaninyl[-alaninyl[-alaninyl[-alaninyphenylglycine methyl ester N-(2-phenylethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-10 alaninamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophanamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3cyclohexylpropionate methyl ester N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-15 amino-3-(4-nitrophenyl)propionamide N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-serine ethyl ester N-[(R)- $\alpha$ -methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[(S)- $\alpha$ -methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 alaninamide N-(4-fluorobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-(4-pyridylmethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-25 alaninamide N-(4-trifluoromethylbenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-phenylpropionate ethyl ester

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine tent-butylester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-methylpropionate methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-cyclohexylacetate ethyl ester
	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N-[N-(isovaleryl)-2-amino-2-cyclohexylacetyl]-L-alanine ethyl ester
10	N-2-(N,N-dimethylamino) ethyl- $N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide$
	N-(2-pyridylmethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N-[N-(3-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
15	N-[N-(2-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(4-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-fluorophenyl)acetate ethyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-fluorophenyl)acetate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-phthalimidopropionate ethyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine neopentyl ester
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine <i>tert</i> -butyl ester

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide 4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valinyl]morpholine N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threonine methyl ester5 N-[N-(3,5-difluor ophenylacetyl)-L-alaninyl]-(S)-2-amin open tanoatemethyl ester 4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-amino-3-tertbutoxybutyryl]morpholine  $4\hbox{-}[N\hbox{-}(3\hbox{-nitrophenylacetyl})\hbox{-}L\hbox{-alaninyl}]\hbox{-}L\hbox{-isoleucinyl}] morpholine$ N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine methyl ester 10 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine N-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threoninyl]-L-valine ethylester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl 15 ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-leucine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine methyl ester N-2-methoxyethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-2-(N,N-dimethylamino) ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-20 alaninyl]-L-alaninamide N-cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-neopentyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide  $\label{eq:N-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl]-L-alaninyl[-L-alaninyl[-L-alaninyl]-L-alaninyl[-$ 25 alaninamide N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide

3-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]thiazolidine N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester 5 N-(R)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 1-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]pyrrolidine N-(S)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valine methyl ester N-2-fluoroethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[(S)-6-methyl-3-oxohept-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide 15 N-4-nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminobutyramide N-4-nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminopentanamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-20 fluorophenyl)acetate methyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2thienyl)acetamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(5chlorobenzothiophen-2-yl)acetate methyl ester 25 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-2yl)acetate ethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-3yl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-thienyl)acetate 30 methyl ester

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-5 yl)acetate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate <i>tert</i> -butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetic acid
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(1H-tetrazol-5-yl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(6-methoxy-2 naphthyl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-trifluoromethylphenyl)acetate methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4,5,6,7-tetrahydrobenzothiophen-2-yl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(thieno[2,3-b]thiophen-2-yl)acetate methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-methylthiazol-4-yl)acetate methyl ester
	(3S,4S)-N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-4-amino-3-hydroxy-5-phenylpentanoate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohex-4-enoate methyl ester
25	N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(4-phenylphenyl)acetamide
	N-[N-(3,5-difluorophenylacetyl)-(S)-2-aminobutanoyl]-L-phenylglycine tert-Butyl Ester
30	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine tert-butyl ester

	N-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-leucinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylalaninyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)glycinyl]-L-phenylglycine methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-alanine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-leucine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-isoleucine methyl ester
15	N-[N-(phenylacetyl)-L-alaninyl]-L-proline methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-N-(tert-butoxycarbonyl)-L-lysine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-glycine methyl ester
20	N-[N-(phenylacetyl)-L-alaninyl]-L-valine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine
	N-[N-(phenylacetyl)-L-alaninyl]-L-N-methylalanine methyl ester
25	N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester
	N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester

	N-Cyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-hydroxyproline ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-lysine methyl ester
•	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-glutamide
•	1-[N-(3,5-difluorophenylacetyl)-L-alaninyl]piperidine-2-carboxylate methyl ester
10	N-[(S)-3-hydroxy-6-methylhept-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(S)-2-hydroxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
15	$N$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ -fluoroacetyl)-L-alaniny]-L-phenylglycine <i>tert</i> -butyl ester
	N-[N-(3,5-difluorophenylacetyl)-2-(S)-aminocyclohexylacetyl]-L-phenylglycine methyl ester
	N-[(1R,2S)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
20	N-[(1R,2S)-1-hydroxy-1,2-diphenyleth-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(1S,2R)-1-hydroxy-1-phenylprop-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
25	N-2-methoxyethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-glycinamide
	$N$ -[(S)- $\alpha$ -hydroxy- $\alpha$ -phenyl- $iso$ -propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(S)-2-hydroxy-1,2-diphenylethyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
30	N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

 $N-[\alpha-hydroxy-\alpha'-(4-hydroxyphenyl)-iso-propyl]-N'-(3,5$ difluorophenylacetyl)-L-alaninamide N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide 5  $N-[\alpha-hydroxy-\alpha'-pyrid-2-y]-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L$ alaninamide  $N-[\alpha-hydroxy-\alpha'-pyrid-4-yl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L$ alaninamide N-[(S)-1-hydroxy-4-methylpent-2-yl]-N'-(3,5-difluorophenylacetyl)-L-10 alaninamide  $N-[\alpha-\text{methoxy-prop-2-yl}]-N'-(3,5-\text{difluorophenylacetyl})-L-alaninamide$ N-[1-hydroxy-3-methyl-but-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(6-aminopyrid-2-15 yl)acetate methyl ester N-[1-hydroxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide N-[(S)-2-methoxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[(S)-1-methoxy-2-phenyl-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide 20 N-[(S)-1-acetoxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide N-[(S)-1-(tert-butylcarbonyloxy)-hex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide N-[2-hydroxy-1-(thien-2-yl)ethyl]-N'-(3,5-difluorophenylacetyl)-L-25 alaninamide N-[(S)-2-hydroxy-2-methyl-1-phenylprop-1-yl]-N'-(3,5difluorophenylacetyl)-L-alaninamide N-[N-(3,5-difluorophenylacetyl)-L-(thien-2-yl)glycinyl]-L-phenylalanine tert-butyl ester

N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinol

N-[N-(cyclopropaneacetyl)-L-phenylglycinyl]-L-phenylglycinol N-[N-(cyclopentaneacetyl)-L-phenylglycinyl]-L-phenylglycinol N-[N-(3,5-difluorophenylacetyl)-D,L-phenylglycinyl]-D,Lphenylglycinamide 5 N-[N-(3,5-difluorophenylacetyl)-D,L-valinyl]-D,L-phenylglycinamide N-[N-(2-thienylacetyl)-L-alaninyl]-L-phenylglycinamide N-[N-(n-caprotyl)-L-alaninyl]-L-phenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-norleucinyl]-L-phenylglycine methyl ester 10 N-[N-(3,5-difluorophenylacetyl)-L-norvalinyl]-L-phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-tert-leucinyl]-L-phenylglycine methyl N-[N-(3,5-difluorophenylacetyl)-L-isoleucinyl]-L-phenylglycine methyl 15 ester N-[N-(3,5-difluorophenylacetyl)-L-cyclohexylalaninyl]-L-phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(cyclopropyl)acetyl]-Lphenylglycine methyl ester 20 N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-3-yl)acetyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-2-yl)acetyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-(4-fluorophenyl)glycinyl]-L-25 phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-D-(4-fluorophenyl)glycinyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-(4-methoxyphenyl)glycinyl]-Lphenylglycine methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine tertbutyl ester N-[N-(cyclopropylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester 5 N-[N-(cyclopentylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester N-[N-(tert-butylacetyl)-L-alaninyl]-L-phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(5bromothien-2-yl)glycinamide 10 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(5bromothien-2-yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4bromothien-2-yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-2-15 yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-3yl)glycinamide 20 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Dphenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-25 phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(5chlorothien-2-yl)glycinamide N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-4-(phenyl)phenylglycinamide 30 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenoxy)phenylglycinamide

	$N$ -(S)-(-)- $\alpha$ -methylbenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenyl)phenylglycinamide
5	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(ethyl)phenylglycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-(phenyl)phenylglycinamide
10	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -2-(benzyl)phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -4-bromophenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -4-(cyclohexyl)phenylglycinamide
15	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(4-ethylphenyl)phenylglycinamide
	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-( <i>tert</i> -butyl)phenylglycinamide
20	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-3-(4-chlorophenoxy)phenylglycinamide
	N-cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(phenyl)phenylglycinamide
	$N$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ -hydroxyacetyl)-L-alaninyl]-L-phenylglycine $tert$ -butyl ester
25	$N$ -tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetyl)-L-alaninyl]-L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine <i>tert</i> -butyl ester
30	N-[(S)-1-oxo-1-phenylprop-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(pyrid-3-yl)glycine tert-butyl ester
	[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]morpholine
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(2-methoxy)phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-butoxycarbonyl(hydroxyl amine) ester
10	N-neopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-tetrahydrofurfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-methoxy- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
15	[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]azetidine
	N-iso-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
20	N-cyclopropanemethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-methoxy- $N$ -methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-2-methylprop-2-enyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
25	N-(pyrid-3-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-(pyrid-4-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
30	N-furfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide

N-cyclopentyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide N-1-benzylpiperidin-4-yl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide N,N-dimethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-5 phenylglycinamide N-2,2,6,6-tetramethylpiperidin-4-yl-N'-[N-(3,5-difluorophenylacetyl)-Lalaninyl]-D,L-phenylglycinamide N-2-methylcyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-10 phenylglycinamide N-4-methylcyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide N-1-ethoxycarbonylpiperidin-4-yl-N'-[N-(3,5-difluorophenylacetyl)-Lalaninyl]-D,L-phenylglycinamide 15 N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-tert-butoxy-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-20 butyl(hydroxylamine) ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine hydrazide N-(1-ethoxyethen-1-yl)-[N'-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycine hydrazide N-[N-(phenylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester 25 N-4-(phenyl)butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide

N-6-(amino)hexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide Hydrochloride

L-phenylglycinamide

N-3-(4-iodophenoxy)propyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-

N-1-(phthalimido)pent-2-yl-N'-(3,5-difluorophenylacetyl)-L-alaninamide N-[N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenyl)glycinyl]-L-(3,5difluorophenyl)glycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-norleucine 5 N-[N-(cyclopentaneacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine isopropyl ester N-(isopropyl) N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide 10 N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine iso-butyl N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycine methyl 15  $N-[N-(3.5-Diffuorophenylacetyl)-L-alaninyl]-L-(3-\alpha-phenyl)proline$ methyl ester N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-azetidine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(5-20 chlorobenzothiophen-2-yl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4yl)propionate tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide tertbutyl ester 25 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide N-[N-(3,4-dichlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(3-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(3-bromophenylacetyl)-L-alaninyl]-D-phenylglycinamide

	N-[N-(3-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
•	N-[N-(4-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(4-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide
5	N-[N-(3-trifluoromethylphenylacetyl)-L-alaninyl]-D-phenylglycinamid
	N-[N-(3-methoxyphenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(2-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(1-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(2-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide
10	N-[N-(phenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-furanyl)acetamide
15	N'-[ $N$ -(3,5-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide
	N'-[ $N$ -(3,4-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanin-N-methylsulfonamide
20	N''-methyl- $N''$ -phenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-glycinamide
	N''-methyl- $N''$ -phenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycinamide
25	N''-methyl- $N''$ -benzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-glycinamide

N''-4-fluorobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-fluoro)phenylglycine neopentyl ester 5 N-[N-(2,3,4,5,6-pentafluorophenylacetyl)-L-alaninyl]-L-(pyrid-3yl)glycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine tertbutyl ester N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)serinyl]-L-phenylglycine 10 methyl ester N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)threoninyl]-L-phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-threoninyl]-L-phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-serinyl]-L-phenylglycine methyl ester 15 N"-4-methylphenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N"-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenyl-20 glycinamide N'-[N-(3,5-difluorophenylacetyl)-L-methionyl]-L-phenylglycinamide N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinamide 25 N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycinamide  $N-[(R)-\alpha-methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L$ phenylglycinamide N-[1-phenyl-2-oxo-3-methylbutan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-

alaninamide

N-[1-phenyl-2-oxo-propan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[1-phenyl-2-oxo-pentan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[1-phenyl-2-oxo-2-phenyl-ethan-1-yl]-N'-(3,5-difluorophenyl-acetyl)-L-5 alaninamide N-[1-phenyl-2-oxo-butan-1-yl]-N'-(3,5-difluorophenyl-acetyl)-Lalaninamide N-[1-phenyl-2-oxo-4-methylpentan-1-yl]-N'-(3,5-difluorophenyl-acetyl)-10 L-alaninamide  $N'-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-\alpha-hydroxyphenylalanine$ methyl ester N''-[4-((2-hydroxy-4-azido)-phenyl)-NHC(O)-)butyl] N'-[N-(3,5difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide N-[(S)-1-phenyl-2-oxo-2-phenyl-ethan-1-yl]-N'-(3,5-difluorophenyl-15 acetyl)-L-alaninamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine tert-butyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-phenylphenylglycine 20 tert-butyl ester [N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(2,3-benzo[b]proline) methyl ester N''-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-nbutylphenylglycinamide N"-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-25 (phenylacetenyl)phenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide N-[1,3-diphenyl-2-oxo-propan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide 30 N-[1-phenyl-2-oxo-2-cyclopentylethan-1-yl]-N'-(3,5-

difluorophenylacetyl)-L-alaninamide

N-[1-phenyl-2-oxo-hexan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[1-phenyl-2-oxo-3-methylpentan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide 5 N''-n-hexyl-6-biotinamidyl-N'-[N-(3.5-difluorophenylacetyl)-L-alaninyl]-D.L-phenylglycinthioamide N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-methionine N'-[N-(2-tert-BOC-amino)propionyl)-L-alaninyl]-L-phenylglycine methyl ester 10 N"-tert-butyl N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2fluorophenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-2-phenylglycine methylester N-[(S)-1-phenyl-2-oxo-3-phenylpropan-1-yl]-N'-(3,5-15 difluorophenylacetyl)-L-alaninamide N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2phenylglycine N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2phenylglycine tert-butyl ester 20 N'-[N-(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine N'-[N-(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine tert-butyl ester N-[2-hydroxy-1-(S)phenyleth-1-yl]-N'-[(3,5-difluorophenylacetyl)-Lphenylglycinyl]-L-alaninamide 25 N-[2-hydroxyeth-1-yl]-N'-[(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N'-[N-(3,5-difluorophenyl-2-oxo-acetyl)-L-alaninyl]-L-2-phenylglycinetert-butyl ester [N-(2,5-dichlorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester 30 [N-(3,5-difluorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

20

25

30

[N-(3,4-dichlorothiophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3-aminoproprionyl)-L-alaninyl]-L-phenylglycine *tert*-butyl ester; and [N-(3-tert-butoxycarbonylamino)propionyl)-L-alaninyl]-L-phenylglycine *tert*-butyl ester.

32. A pharmaceutical composition comprising a pharmaceutically inert carrier and pharmaceutically effective amount of a compound of formula I:

10
$$R^{1}Z$$

$$R^{2}$$

$$R^{3}$$

$$R^{3}$$

wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

each R<sup>3</sup> is independently selected from the group consisting of hydrogen and methyl and R<sup>3</sup> together with R<sup>4</sup> can be fused to form a cyclic structure of from 3 to 8 atoms which is optionally fused with an aryl or heteroaryl group;

each R<sup>4</sup> is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclic, substituted alkyl, substituted alkenyl and substituted alkynyl;

each R<sup>5</sup> is selected from hydrogen and methyl or together with R<sup>4</sup> forms a cycloalkyl group of from 3 to 6 carbon atoms;

X is selected from the group consisting of -C(O)Y and -C(S)Y where Y is selected from the group consisting of

(a) alkyl or cycloalkyl,

- (b) substituted alkyl with the proviso that the substitution on said substituted alkyl do not include  $\alpha$ -haloalkyl,  $\alpha$ -diazoalkyl,  $\alpha$ -OC(O)alkyl, or  $\alpha$ -OC(O)aryl groups,
  - (c) alkoxy or thioalkoxy,
  - (d) substituted alkoxy or substituted thioalkoxy,
  - (e) hydroxy,
  - (f) aryl,
  - (g) heteroaryl,
  - (h) heterocyclic,
- 10

20

5

- (i) -NR'R" where R' and R" are independently selected from hydrogen, alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl, substituted alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, where one of R' or R" is hydroxy or alkoxy, and where R' and R" are joined to form a cyclic group having from 2 to 8 carbon atoms optionally containing 1 to 2 additional heteroatoms selected from oxygen, sulfur and nitrogen and optionally substituted with one or more alkyl, alkoxy or carboxylalkyl groups,
- (j) -NHSO<sub>2</sub>-R<sup>8</sup> where R<sup>8</sup> is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic,
- (k) -NR<sup>9</sup>NR<sup>10</sup>R<sup>10</sup> where R<sup>9</sup> is hydrogen or alkyl, and each R<sup>10</sup> is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, and
- (1)  $-ONR^9[C(O)O]_zR^{10}$  where z is zero or one,  $R^9$  and  $R^{10}$  are as defined above;

25 the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic and Y' is selected from the group consisting of hydroxyl, amino, thiol, alkoxy, substituted alkoxy, thioalkoxy, substituted thioalkoxy, -OC(O)R<sup>7</sup>, -SSR<sup>7</sup>, -SSC(O)R<sup>7</sup> where R<sup>7</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

X' is hydrogen, hydroxy, or fluoro;

10

15

20

25

X'' is hydrogen, hydroxy or fluoro, or X' and X'' together form an oxo group,

Z is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X"-, oxygen and sulfur;

n is an integer equal to 1 or 2; and pharmaceutically acceptable salts thereof with the provisos that:

A. when  $R^1$  is phenyl or 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH;

B. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub> derived from D-threonine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH or -C(O)OCH<sub>3</sub>;

C. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^4$  is benzyl,  $R^5$  is hydrogen, X is methoxycarbonyl, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$  is not methyl;

D. when  $R^1$  is iso-propyl,  $R^2$  is  $-CH_2C(O)NH_2$ ,  $R^3$  is hydrogen,  $R^4$  is iso-butyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

E. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^5$  is hydrogen, X is -C(O)OCH<sub>3</sub>, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$ , the nitrogen atom attached to  $R^3$ , and  $R^4$  do not form 1,2,3,4-tetrahydroiso-quinolin-2-yl or pyrrolidin-2-yl;

F. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen,  $X^4$  is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and  $R^4$  is not 4-amino-R-butyl;

G. when  $R^1$  is 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH<sub>2</sub> or -CH<sub>2</sub>OH;

- H. when R<sup>1</sup> is phenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is hydrogen, R<sup>5</sup> is hydrogen, X is -CH<sub>2</sub>OCH<sub>3</sub>, X' and X" are hydrogen, Z is a bond, and n is 1, then R<sup>4</sup> is not benzyl or ethyl;
- I. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is methyl, R<sup>4</sup> is methyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$ ;

10

15

20

- J. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is hydrogen, R<sup>4</sup> is phenyl derived from D-phenylglycine, R5 is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$  or -CH<sub>2</sub>OH;
- K. when  $R_1$  is N-(2-pyrrolidinonyl),  $R_2$  is methyl,  $R_3$  is hydrogen,  $R_4$  is benzyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;
  - L. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl derived from D-alanine, R<sup>3</sup> is hydrogen, R<sup>4</sup> is phenyl derived from D-phenylglycine, R<sup>5</sup> is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH-benzyl;
  - M. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is hydrogen, R<sup>4</sup> is hydrogen,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CH<sub>2</sub>OH;
  - N. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is hydrogen, R<sup>4</sup> is 4-phenylphenyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NHC(CH<sub>3</sub>)<sub>3</sub>; and
  - O. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is hydrogen, R<sup>4</sup> is phenyl derived from D-phenylglycine, R<sup>5</sup> is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)NHCH(CH_3)\phi$ .
- 25 33. The pharmaceutical composition according to Claim 32 wherein R<sup>1</sup> is an unsubstituted aryl group and Z is a bond covalently linking R1 to -CX'X"-.
  - 34. The pharmaceutical composition according to Claim 33 wherein the unsubstituted R<sup>1</sup> aryl group is selected from the group consisting of phenyl,

1-naphthyl and 2-naphthyl.

- 35. The pharmaceutical composition according to Claim 32 wherein  $R^1$  is a substituted aryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 36. The pharmaceutical composition according to Claim 35 wherein said substituted aryl group is a mono-substituted, di-substituted or tri-substituted phenyl group.
- 37. The pharmaceutical composition according to Claim 36 wherein the substituted phenyl groups are selected from the group consisting of 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 3-methoxy-phenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2-hydroxy-phenyl, 2-methylphenyl, 2-fluorophenyl, 2-chlorophenyl, 3,4-difluorophenyl, 2,3,4,5,6-pentafluorophenyl, 3,4-dibromophenyl, 3,4-dichlorophenyl, 3,4-methylene-dioxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 2,4-dichlorophenyl, and 2,5-difluorophenyl.
  - 38. The pharmaceutical composition according to Claim 32 wherein  $R^1$  is an alkaryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
  - 39. The pharmaceutical composition according to Claim 38 wherein the R<sup>1</sup> alkaryl group is selected from the group consisting of benzyl, 2-phenylethyl, and 3-phenyl-*n*-propyl.
  - 40. The pharmaceutical composition according to Claim 32 wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl and cycloalkenyl groups and Z is a bond covalently linking R<sup>1</sup> to -CX'X"-.

- 41. The pharmaceutical composition according to Claim 40 wherein R<sup>1</sup> is alkyl.
- 42. The pharmaceutical composition according to Claim 40 wherein R<sup>1</sup> is cycloalkyl.
- 5 43. The pharmaceutical composition according to Claim 40 wherein R<sup>1</sup> is alkenyl.
  - 44. The pharmaceutical composition according to Claim 40 wherein R<sup>1</sup> is cycloalkenyl.
- 45. The pharmaceutical composition according to Claim 40 wherein the R¹ alkyl, cycloalkyl, alkenyl and cycloalkenyl groups are selected from the group consisting of *iso*-propyl, *n*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclobutyl, aminomethyl, and N-*tert*-butoxycarbonylaminomethyl.
  - 46. The pharmaceutical composition according to Claim 32 wherein R<sup>1</sup> is selected from the group consisting of heteroaryl and substituted heteroaryl groups and Z is a bond covalently linking R<sup>1</sup> to -CX'X"-.
- 47. The pharmaceutical composition according to Claim 46 wherein the R<sup>1</sup> heteroaryl and substituted heteroaryl groups are selected from the group consisting of pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, fluoropyridyls (including 5-fluoropyrid-3-yl), chloropyridyls (including 5-chloropyrid-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-

10

(thiophenyl)thiophen-5-yl, 6-methoxythionaphthen-2-yl, 3-phenyl-1,2,4-thiooxadiazol-5-yl and 2-phenyloxazol-4-yl.

- 48. The pharmaceutical composition according to Claim 32 wherein R<sup>2</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic.
- 49. The pharmaceutical composition according to Claim 48 wherein  $R^2$  is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, phenyl, 4-fluorophenyl, 3,5-difluoro-phenyl, 4-methoxyphenyl, benzyl, cyclopropyl, cyclohexyl, cyclopentyl, cycloheptyl, thien-2-yl, thien-3-yl, -CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>, -CH<sub>2</sub>OCH<sub>2</sub> $\phi$ , -CH(CH<sub>3</sub>)OCH<sub>2</sub> $\phi$ , -CH(OH)CH<sub>3</sub> and -CH<sub>2</sub>OH.
- 50. The pharmaceutical composition according to Claim 32 wherein X' and X" are hydrogen and Z is a bond covalently linking R<sup>1</sup> to -CX'X"-.
- 15 51. The pharmaceutical composition according to Claim 50 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, methyl or together with R<sup>4</sup> and the nitrogen to which R<sup>3</sup> is attached forms pyrrolidin-2-yl, 2,3-dihydroindol-2-yl, piperidin-2-yl, 4-hydroxy-pyrrolidin-2-yl and 1,2,3,4-tetrahydroisoquinolin-3-yl.
- 52. The pharmaceutical composition according to Claim 32 wherein R<sup>4</sup> substituents are selected from the group consisting of hydrogen, methyl, ethyl, iso-propyl, n-propyl, n-butyl, sec-butyl, iso-butyl, cyclopentyl, cyclohexyl, allyl, iso-but-2-enyl, 3-methylpentyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-indol-3-yl, phenyl, p-(phenyl)phenyl, m-(phenyl)phenyl, o-fluorophenyl, m-fluorophenyl, p-fluorophenyl, p-bromophenyl, m-methoxyphenyl, p-methoxyphenyl, phenethyl, benzyl, m-hydroxybenzyl,

p-hydroxybenzyl, p-nitrobenzyl, m-trifluoromethylphenyl. p-(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, p-(CH<sub>3</sub>)<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl. p-phenylphenyl, 3,5-difluorophenyl, p-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyrid-6yl, 4-(N-morpholino-CH2CH2O)-benzyl, -CH2CH2C(O)NH2, -CH2-imidazol-4-5 yl, -CH<sub>2</sub>-(3-tetrahydrofuranyl), -CH<sub>2</sub>-thien-2-yl, -CH<sub>2</sub>-thiazol-4-yl, -CH<sub>2</sub>(1methyl)cyclopropyl, -CH<sub>2</sub>-thien-3-yl, thien-3-yl, thien-2-yl, -CH<sub>2</sub>-C(O)O-tbutyl, -CH<sub>2</sub>-C(CH<sub>3</sub>)<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, 2-methylcyclopentyl, -cyclohex-2enyl,  $-CH[CH(CH_3)_2]COOCH_3$ ,  $-(CH_2)_2SCH_3$ ,  $-CH_2CH_2N(CH_3)_2$ , -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=CHCH<sub>3</sub> (cis and trans), -CH<sub>2</sub>OH, -CH(OH)CH<sub>3</sub>, 10  $-CH(O-t-butyl)CH_3$ ,  $-CH_2OCH_3$ ,  $-(CH_2)_4NH-Boc$ ,  $-(CH_2)_4NH_2$ ,  $-(CH_2)_4N(CH_3)_2$ , -CH<sub>2</sub>-pyridyl, pyridyl, -CH<sub>2</sub>-naphthyl, -CH<sub>2</sub>-(N-morpholino), p-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5chlorobenzo[b]thiophen-2-yl, 4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, tetrazol-5-yl, 5-chlorobenzo[b]thiophen-3-yl, 15 benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl, -CH<sub>2</sub>-N-phthalimidyl, 2-methylthiazol-4-yl, and thieno[2,3-b]thiophen-2-yl, 5-bromothien-2-yl, 4-bromothien-2-yl, 5-chlorothien-2-yl, 3-phenoxyphenyl, 2-phenoxyphenyl, 4-ethylphenyl, 2-benzylphenyl, (4-ethylphenyl)phenyl, 4-tertbutylphenyl, 4-n-butylphenyl, o-(4-chlorophenoxy)phenyl, furan-2-yl, and 4-20 phenylacetylenylphenyl.

- 53. The pharmaceutical composition according to Claim 32 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"- and R<sup>4</sup> and R<sup>5</sup> are fused to form a cycloalkyl group selected from the group consisting of cyclopropyl and cyclobutyl.
- 25 54. The pharmaceutical composition according to Claim 32 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"-, X is -C(O)Y and Y is selected from the group consisting of hydroxy, alkoxy or substituted alkoxy.

- 55. The pharmaceutical composition according to Claim 54 wherein Y is alkoxy or substituted alkoxy selected from the group consisting of methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *iso*-butoxy, *tert*-butoxy, *neo*-pentoxy, benzyloxy, 2-phenylethoxy, 3-phenyl-*n*-propoxy, 3-iodo-*n*-propoxy, 4-bromo-*n*-butoxy, -ONHC(O)OC(CH<sub>3</sub>)<sub>3</sub>, -ONHC(CH<sub>3</sub>)<sub>3</sub> and hydroxy.
- 56. The pharmaceutical composition according to Claim 32 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"-, X is -C(O)Y and Y is -NR'R".
- 57. The pharmaceutical composition according to Claim 56 wherein Y 10 is selected from the group consisting of amino (-NH<sub>2</sub>), -NH(iso-butyl), -NH(sec-butyl), N-methylamino, N,N-dimethylamino, N-benzylamino, N-morpholino, azetidino, N-thiomorpholino, N-piperidinyl, N-hexamethyleneimino, N-heptamethylene-imino, N-pyrrolidinyl, -NHmethallyl, -NHCH2-(furan-2-yl), -NHCH2-cyclopropyl, -NH(tert-butyl), -NH(p-15 methylphenyl), -NHOCH<sub>3</sub>, -NHCH<sub>2</sub>(p-fluorophenyl), -NHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, -NHcyclopentyl, -NH-cyclohexyl, -NHCH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -NHCH<sub>2</sub>C(CH<sub>4</sub>)<sub>3</sub>, -NHCH<sub>2</sub>-(pyrid-2-yl), -NHCH<sub>2</sub>-(pyrid-3-yl), -NHCH<sub>2</sub>-(pyrid-4-yl), N-thiazolindinyl,  $-N(CH_2CH_3)_2$ ,  $-N[CH_2CH(CH_3)_2]_2$ , -NHOH,  $-NH(p-NO_2-\phi)$ ,  $-NHCH_2(p-NO_2-\phi)$  $NO_2-\phi$ ),  $-NHCH_2(m-NO_2-\phi)$ ,  $-N(CH_3)OCH_3$ ,  $-N(CH_3)CH_2-\phi$ ,  $-NHCH_2-(3,5-di-1)$ 20 fluorophenyl), -NHCH<sub>2</sub>CH<sub>2</sub>F, -NHCH<sub>2</sub>(p-CH<sub>3</sub>O- $\phi$ ), -NHCH<sub>2</sub>(m-CH<sub>3</sub>O- $\phi$ ),  $-NHCH<sub>2</sub>(p-CF<sub>3</sub>-\phi)$ , -N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>,  $-NHCH<sub>2</sub>CH<sub>2</sub>\phi$ ,  $-NHCH(CH<sub>3</sub>)\phi$ ,  $-NHCH_2-(p-F-\phi)$ ,  $-N(CH_3)CH_2CH_2N(CH_3)_2$ ,  $-NHCH_2-(tetrahydrofuran-2-yl)$ ,  $-NHCH_2(p-trifluoromethylphenyl), -NHCH_2C(CH_3) = CH_2, -NH-I(p-trifluoromethylphenyl), -NHCH_2C(CH_3) = CH_3, -NH-I(p-trifluoromethylphenyl$ benzyl)pyrid-4-yl], -NH-[(2,6-dimethyl)pyrid-4-yl], -NH-(2-methylcyclohexyl), 25 -NH-(4-methylcyclohexyl), -NH-[N-ethoxycarbonyl]-piperidin-4-yl, -NHOC(CH<sub>3</sub>)<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- $\phi$ , -C(O)NH(CH<sub>2</sub>)<sub>3</sub>O-(p-CH<sub>3</sub>) $\phi$ , -C(O)NH(CH<sub>2</sub>)<sub>6</sub>NH<sub>2</sub>, -NH-(tetrahydrofuran-2-yl), -N(CH<sub>3</sub>) $\phi$ , -NH(CH<sub>2</sub>)<sub>4</sub>NHC(O)-(2-hydroxy-4-azido)-phenyl and -NH(CH<sub>2</sub>)<sub>6</sub>-(biotinamidyl).

25

- 58. The pharmaceutical composition according to Claim 32 wherein X is -C(O)Y and Y is selected from the group consisting of -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OH, -CH(OH)CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(OH)φ, -CH(OH)CH<sub>2</sub>C(O)OCH<sub>3</sub>, -C(OH)(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>OC(O)OCH<sub>3</sub>, and -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>, methyl, ethyl, iso-propyl, n-propyl, iso-butyl, n-butyl, sec-butyl, tert-butyl, -CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-pyridy-2-yl, -CH<sub>2</sub>-pyridy-3-yl, -CH<sub>2</sub>-pyridy-4-yl, -CH<sub>2</sub>-fur-2-yl, benzyl, cyclopentyl, phenyl, and -NH-SO<sub>2</sub>-CH<sub>3</sub>.
- 59. The pharmaceutical composition according to Claim 32 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"-.
  - 60. The pharmaceutical composition according to Claim 32 wherein the compound of formula I is selected from the group consisting of:

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-histidine methyl ester

N-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

N-2-(N,N-dimethylamino) ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

20 N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

N-2-(N,N-dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

N-(4-pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

*N*-(3-pyridyl)methyl-*N'*-[*N*-(3,5-difluorophenylacetyl)-**L**-alaninyl]-**L**-phenylalaninamide

	N-(4-pyridyl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-( $S$ )-2-aminohexanamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate tert-butyl ester
5	N-[N-(pent-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(dec-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[3-(N,N-dimethylamino)propoxy]phenylalanine methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[(tert-butyloxycarbonyl)methoxy]phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4- (carboxymethoxy)phenylalanine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-6-(N,N-dimethylamino)hexanoate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-pyridyl)propionate methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(3-pyridyl)propionate methyl ester
•	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-proline methyl ester
	1-[N-(3,5-difluorophenylacetyl)-L-alaninyl]piperidine-2-carboxylate methyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4-pyridyl)propionate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionate methyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-morpholinopropionate methyl ester

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)phenylalaninamide

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionamide

5 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine methyl ester

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(4-pyridyl)propionamide

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(2-pyridyl)propionamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-yl)propionate methyl ester

2-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1,2,3,4-tetrahydroisoquinoline-3-carboxylate methyl ester

N-(3-methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(1-naphthyl)propionate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-naphthyl)propionate methyl ester

20 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-thienyl)propionate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine benzyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-bromo-propyl ester

25 *N*-[*N*-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-iodopropyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine tert-butyl ester

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetamide

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3pyridyl)acetamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-N-(tert-butoxycarbonyl)-Llysine methyl ester 5 methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-4phenylbutanoate N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 2-phenylethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 3-phenylpropyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-10 pyridyl)acetamide N-[N-(phenylacetyl)-L-alaninyl]-L-threonine methyl ester N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N'-[N-(phenylacetyl)-L-alaninyl]-L-alaninamide N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide 15 N'-[N-(phenylacetyl)-L-alaninyl)-L-valinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetate ethyl ester N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N,N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide N,N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide 20 N, N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-25 aminohexanamide N, N-dimethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminohexanamide

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3methoxyphenyl)acetate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4methoxyphenyl)acetate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetate ethyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetate ethyl ester

10 N-[N-(cyclohexylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(cyclohex-1-enylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1-aminocyclopropane-1carboxylate methyl ester

15 N-2-(N,N-dimethylamino)ethyl-N-methyl-N'-[N-(3,5difluorophenylacetyl)-L-alaninyl]-L-alaninamide

5

N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine benzyl ester

N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine ethyl ester

20 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-alanine ethyl ester

N-[N-(3-nitrophenylacetyl)-L-alaninyl]glycine ethyl ester

N-hydroxy-N'-[N-(3-nitrophenylacetyl)-L-alaninyl]-D,L-threoninamide

N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester

N-[N-(3-nitrophenylacetyl)-L-alaninyl]-2-amino-3-(3-25 hydroxyphenyl)propionate methyl ester

N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-tyrosine ethyl ester

N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester N-[N-(isovaleryl)-L-valinyl]-L-phenylglycinyl]-L-alanine iso-butyl ester N-[N-(isovaleryl)-L-phenylalaninyl]-L-alanine iso-butyl ester 5 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester 1-[N-(3-nitrophenylacetyl)-L-alaninyl]-indoline-(S)-2-carboxylate ethyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-methoxy-N-methyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide 10 N-iso-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N,N-di-n-propyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N'-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-valinamideN-(4-nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-15 alaninamide N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninyl]-L-phenylalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamideN-iso-butyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 phenylalaninamide N-(4-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-(4-nitrophenyl)-N'-[N-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninyl]-L-25 alaninamide N-(4-nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide

N-benzyl-N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-(3,5-difluorobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-(3-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 alaninamide N-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-(4-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophan methyl ester N-(4-methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[N-(phenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninyl]-L-15 phenylglycine methyl ester N-[N-(cyclohexylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine methyl ester N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]-Lphenylglycine methyl ester 20 N-(2-phenylethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophanamideN-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3cyclohexylpropionate methyl ester 25 N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2amino-3-(4-nitrophenyl)propionamide N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-serine ethyl ester  $N-[(R)-\alpha-methylbenzyl]-N'-[N-(3.5-difluorophenylacetyl)-L-alaninyl]-L$ alaninamide

	$N$ -[(S)- $\alpha$ -methylbenzyl]- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide
	N-(4-fluorobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
5	N-(4-pyridylmethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
. •	N-(4-trifluoromethylbenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-phenylpropionate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-methylpropionate methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-cyclohexylacetate ethyl ester
	N-(2-methoxyethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N-[N-(isovaleryl)-2-amino-2-cyclohexylacetyl]-L-alanine ethyl ester
20	N-2-(N,N-dimethylamino) ethyl- $N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl-L-phenylglycinamide$
	N-(2-pyridylmethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N-[N-(3-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
25	N-[N-(2-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(4-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-fluorophenyl)acetate ethyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-fluorophenyl)acetate ethyl ester

N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3phthalimidopropionate ethyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine neopentyl 5 ester N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-phenylglycine tert-butylN'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide 10 4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valinyl]morpholine N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threonine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate15 methyl ester 4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-amino-3-tertbutoxybutyryl]morpholine 4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucinyl]morpholine N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine methyl ester 20 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine N-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threoninyl]-L-valine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester 25 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-leucine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine methyl ester N-2-methoxyethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide

N-2-(N,N-dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-Lalaninyl]-L-alaninamide N-cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-neopentyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 alaninamide N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 3-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]thiazolidine 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methylester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester N-(R)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 15 1-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]pyrrolidine N-(S)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valine methyl ester N-2-fluoroethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 alaninamide N-[(S)-6-methyl-3-oxohept-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-4-nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminobutyramide 25 N-4-nitrobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminopentanamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3fluorophenyl)acetate methyl ester

	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(5-chlorobenzothiophen-2-yl)acetate methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-2 yl)acetate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-3 yl)acetate methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-thienyl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-5-yl)acetate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate <i>tert</i> -butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetic acid
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(1H-tetrazol-5-yl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(6-methoxy-2-naphthyl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-trifluoromethylphenyl)acetate methyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4,5,6,7-tetrahydrobenzothiophen-2-yl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(thieno[2,3-b]thiophen-2-yl)acetate methyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-methylthiazol-4-yl)acetate methyl ester

	(3S,4S)-N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-4-amino-3-hydroxy 5-phenylpentanoate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohex-4-enoate methyl ester
5	N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
	N-tert-butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2 (4-phenylphenyl)acetamide
	N-[N-(3,5-difluorophenylacetyl)-(S)-2-aminobutanoyl]-L-phenylglycine tert-Butyl Ester
10	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine tert-butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-leucinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylalaninyl]-L-phenylglycine methyl ester
20	N-[N-(3,5-difluorophenylacetyl)glycinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-alanine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-leucine methyl ester
25	N-[N-(phenylacetyl)-L-alaninyl]-L-isoleucine methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-proline methyl ester
	N-[N-(phenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester

 $N-[N-(phenylacetyl)-L-alaninyl]-N_{\epsilon}-(tert-butoxycarbonyl)-L-lysine methyl$ ester N-[N-(phenylacetyl)-L-alaninyl]-glycine methyl ester N-[N-(phenylacetyl)-L-alaninyl]-L-valine methyl ester 5 N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine N-[N-(phenylacetyl)-L-alaninyl]-L-N-methylalanine methyl ester N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester 10 N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-hydroxyproline ethyl 15 ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-lysine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-glutamide 1-[N-(3,5-difluorophenylacetyl)-L-alaninyl]piperidine-2-carboxylate methyl ester 20 N-[(S)-3-hydroxy-6-methylhept-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[(S)-2-hydroxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide  $N-[N-(3,5-diffuorophenyl-\alpha-fluoroacetyl)-L-alaniny]-L-phenylglycine$ 25 tent-butyl ester N-[N-(3,5-difluorophenylacetyl)-2-(S)-aminocyclohexylacetyl]-Lphenylglycine methyl ester N-[(1R,2S)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-[(1R,2S)-1-hydroxy-1,2-diphenyleth-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

*N*-[(1S,2R)-1-hydroxy-1-phenylprop-2-yl]-*N*'-(3,5-difluorophenylacetyl)-L-alaninamide

5 N-2-methoxyethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-glycinamide

N-[(S)- $\alpha$ -hydroxy- $\alpha$ -phenyl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-[(S)-2-hydroxy-1,2-diphenylethyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-[ $\alpha$ -hydroxy- $\alpha'$ -(4-hydroxyphenyl)-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

 $N-[\alpha-hydroxy-\alpha'-pyrid-2-yl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide$ 

 $N-[\alpha-hydroxy-\alpha'-pyrid-4-yl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide$ 

20 N-[(S)-1-hydroxy-4-methylpent-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

 $N-[\alpha-methoxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide$ 

N-[1-hydroxy-3-methyl-but-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

25 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(6-aminopyrid-2-yl)acetate methyl ester

N-[1-hydroxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-[(S)-2-methoxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

N-[(S)-1-methoxy-2-phenyl-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[(S)-1-acetoxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamideN-[(S)-1-(tert-butylcarbonyloxy)-hex-2-yl]-N'-(3,5-difluorophenylacetyl)-5 L-alaninamide N-[2-hydroxy-1-(thien-2-yl)ethyl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[(S)-2-hydroxy-2-methyl-1-phenylprop-1-yl]-N'-(3,5difluorophenylacetyl)-L-alaninamide N-[N-(3,5-difluor ophenylacetyl)-L-(thien-2-yl)glycinyl]-L-phenylalanine10 tert-butyl ester N-[N-(3,5-diffuor ophenylacetyl)-L-phenylglycinyl]-L-phenylglycinolN-[N-(cyclopropaneacetyl)-L-phenylglycinyl]-L-phenylglycinol N-[N-(cyclopentaneacetyl)-L-phenylglycinyl]-L-phenylglycinolN-[N-(3,5-difluorophenylacetyl)-D,L-phenylglycinyl]-D,L-15 phenylglycinamide N-[N-(3,5-difluor ophenylacetyl)-D,L-valinyl]-D,L-phenylglycinamideN-[N-(2-thienylacetyl)-L-alaninyl]-L-phenylglycinamide N-[N-(n-caprotyl)-L-alaninyl]-L-phenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-norleucinyl]-L-phenylglycine methyl 20 ester N-[N-(3,5-difluorophenylacetyl)-L-norvalinyl]-L-phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-tert-leucinyl]-L-phenylglycine methyl 25 ester N-[N-(3,5-difluorophenylacetyl)-L-isoleucinyl]-L-phenylglycine methylester N-[N-(3,5-difluorophenylacetyl)-L-cyclohexylalaninyl]-L-phenylglycinemethyl ester

N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(cyclopropyl)acetyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-3-yl)acetyl]-Lphenylglycine methyl ester 5 N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-2-yl)acetyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-(4-fluorophenyl)glycinyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-D-(4-fluorophenyl)glycinyl]-L-10 phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-(4-methoxyphenyl)glycinyl]-Lphenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine tertbutyl ester 15 N-[N-(cyclopropylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester N-[N-(cyclopentylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester N-[N-(tert-butylacetyl)-L-alaninyl]-L-phenylglycinamide 20 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(5bromothien-2-yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(5bromothien-2-yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-25 bromothien-2-yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-2yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2yl)glycinamide 30 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-3yl)glycinamide

N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2yl)glycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Dphenylglycinamide 5 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(5chlorothien-2-yl)glycinamide N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-4-10 (phenyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenoxy)phenylglycinamide N-(S)-(-)- $\alpha$ -methylbenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D, L-phenylglycinamide 15 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(ethyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-20 (phenyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-(benzyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4bromophenylglycinamide 25 N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(cyclohexyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(4ethylphenyl)phenylglycinamide N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-(tert-30 butyl)phenylglycinamide

N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-3-(4chlorophenoxy)phenylglycinamide N-cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(phenyl)phenylglycinamide  $N-[N-(3,5-difluorophenyl-\alpha-hydroxyacetyl)-L-alaninyl]-L-phenylglycine$ 5 tert-butyl ester N-tert-butyl-N'-[N-(3,5-difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine tent-butyl10 ester N-[(S)-1-oxo-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(pyrid-3-yl)glycine tertbutyl ester 15 [N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinyl]morpholine N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(2methoxy)phenylglycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-20 butoxycarbonyl(hydroxyl amine) ester N-neopentyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide N-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide 25 N-methoxy-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide [N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinyllazetidine N-iso-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-30 phenylglycinamide

	N-cyclopropanemethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-methoxy- $N$ -methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
5	N-2-methylprop-2-enyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-(pyrid-3-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
10	N-(pyrid-4-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-furfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-cyclopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
15	N-1-benzylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N,N-dimethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
20	N-2,2,6,6-tetramethylpiperidin-4-yl- $N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide$
	N-2-methylcyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-4-methylcyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
25	N-1-ethoxycarbonylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
30	N-tert-butoxy- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide

butyl(hydroxylamine) ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine hydrazide N-(1-ethoxyethen-1-yl)-[N'-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 phenylglycine hydrazide N-[N-(phenylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester N-4-(phenyl)butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-3-(4-iodophenoxy)propyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-10 L-phenylglycinamide N-6-(amino)hexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide Hydrochloride N-1-(phthalimido)pent-2-yl-N'-(3,5-difluorophenylacetyl)-L-alaninamide N-[N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenyl)glycinyl]-L-(3,5-difluorophenylacetyl)15 difluorophenyl)glycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-norleucine N-[N-(cyclopentaneacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine isopropyl ester N-(isopropyl) N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 phenylglycinamide N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine iso-butyl25 ester N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycine methyl $N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(3-\alpha-phenyl)$ proline methyl ester

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-azetidine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(5-difluorophenylacetyl)chlorobenzothiophen-2-yl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-5 yl)propionate tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide tertbutyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide10 N-[N-(3,4-dichlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(3-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(3-bromophenylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(3-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(4-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide15 N-[N-(3-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(4-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamide $N\hbox{-}[N\hbox{-}(3\hbox{-}trifluoromethylphenylacetyl})\hbox{-}L\hbox{-}alaninyl]\hbox{-}D\hbox{-}phenylglycinamide}$ N-[N-(3-methoxyphenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(2-chlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide20 N-[N-(1-naphthylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(2-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(phenylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycineN-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide 25 N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2furanyl)acetamide

N'-[N-(3,5-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide N'-[N-(3,4-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanin-Nmethylsulfonamide 5 N''-methyl-N''-phenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycinamide N''-methyl-N''-phenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycinamide 10 N''-methyl-N''-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycinamide N''-4-fluorobenzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-fluoro)phenylglycine 15 neopentyl ester N-[N-(2,3,4,5,6-pentafluorophenylacetyl)-L-alaninyl]-L-(pyrid-3yl)glycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine tertbutyl ester 20 N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl) serinyl]-L-phenylglycine methyl ester N-[N-(3,5-difluor ophenylacetyl)-L-(O-benzyl) threoninyl]-L-phenylglycinemethyl ester N-[N-(3,5-difluorophenylacetyl)-L-threoninyl]-L-phenylglycine methyl25 ester N-[N-(3,5-difluorophenylacetyl)-L-serinyl]-L-phenylglycine methyl ester N''-4-methylphenyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N''-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-30 phenylglycinamide

	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenyl-glycinamide
	N'-[N-(3,5-difluorophenylacetyl)-L-methionyl]-L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycinamide
. 5	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycinamide
	$N$ -[(R)- $\alpha$ -methylbenzyl]- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
10	N-[1-phenyl-2-oxo-3-methylbutan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-propan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-pentan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
15	N-[1-phenyl-2-oxo-2-phenyl-ethan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-butan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)- $L$ -alaninamide
20	N-[1-phenyl-2-oxo-4-methylpentan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide
	$N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L- $\alpha$ -hydroxyphenylalanine methyl ester
	N''-[4-((2-hydroxy-4-azido)-phenyl)-NHC(O)-)butyl] $N'$ -[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
25	N-[(S)-1-phenyl-2-oxo-2-phenyl-ethan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine tert-butyl ester
30	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-phenylphenylglycine tent-butyl ester

[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(2,3-benzo[b]proline) methyl ester N"-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-nbutylphenylglycinamide 5 N"-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-(phenylacetenyl)phenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide N-[1,3-diphenyl-2-oxo-propan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide 10 N-[1-phenyl-2-oxo-2-cyclopentylethan-1-yl]-N'-(3.5difluorophenylacetyl)-L-alaninamide N-[1-phenyl-2-oxo-hexan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[1-phenyl-2-oxo-3-methylpentan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-15 alaninamide N''-n-hexyl-6-biotinamidyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-methionine N'-[N-(2-tert-BOC-amino)propionyl)-L-alaninyl]-L-phenylglycine methyl 20 ester N"-tert-butyl N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2fluorophenylglycinamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-2-phenylglycine methyl ester 25 N-[(S)-1-phenyl-2-oxo-3-phenylpropan-1-yl]-N'-(3,5difluorophenylacetyl)-L-alaninamide N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2phenylglycine N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-30 phenylglycine tert-butyl ester

15

30

N'-[N-(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine

N'-[N-(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine tert-butyl ester

N-[2-hydroxy-1-(S)phenyleth-1-yl]-N'-[(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-alaninamide

N-[2-hydroxyeth-1-yl]-N'-[(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

N'-[N-(3,5-difluorophenyl-2-oxo-acetyl)-L-alaninyl]-L-2-phenylglycine tert-butyl ester

[N-(2,5-dichlorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester
[N-(3,5-difluorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester
[N-(3,4-dichlorothiophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3-aminoproprionyl)-L-alaninyl]-L-phenylglycine *tert*-butyl ester; and [N-(3-*tert*-butoxycarbonylamino)propionyl)-L-alaninyl]-L-phenylglycine *tert*-butyl ester.

## 61. A compound of formula I:

20 I 
$$R^1Z$$
  $X''$   $X''$   $R^2$   $R^3$ 

wherein R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic;

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

10

25

each R<sup>3</sup> is independently selected from the group consisting of hydrogen and methyl and R<sup>3</sup> together with R<sup>4</sup> can be fused to form a cyclic structure of from 3 to 8 atoms which is optionally fused with an aryl or heteroaryl group;

each R<sup>4</sup> is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, cycloalkenyl, heteroaryl, heterocyclic, substituted alkyl, substituted alkenyl and substituted alkynyl;

each R<sup>5</sup> is selected from hydrogen and methyl or together with R<sup>4</sup> forms a cycloalkyl group of from 3 to 6 carbon atoms;

X is selected from the group consisting of -C(O)Y and -C(S)Y where Y is selected from the group consisting of

- (a) alkyl or cycloalkyl,
- (b) substituted alkyl with the proviso that the substitution on said substituted alkyl do not include  $\alpha$ -haloalkyl,  $\alpha$ -diazoalkyl,  $\alpha$ -OC(O)alkyl, or  $\alpha$ -OC(O)aryl groups,
- 15 (c) alkoxy or thioalkoxy,
  - (d) substituted alkoxy or substituted thioalkoxy,
  - (e) hydroxy,
  - (f) aryl,
  - (g) heteroaryl,
- 20 (h) heterocyclic,
  - (i) -NR'R" where R' and R" are independently selected from hydrogen, alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, where one of R' or R" is hydroxy or alkoxy, and where R' and R" are joined to form a cyclic group having from 2 to 8 carbon atoms optionally containing 1 to 2 additional heteroatoms selected from oxygen, sulfur and nitrogen and optionally substituted with one or more alkyl, alkoxy or carboxylalkyl groups,
  - (j) -NHSO<sub>2</sub>-R<sup>8</sup> where R<sup>8</sup> is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

10

15

20

25

- (k) -NR<sup>9</sup>NR<sup>10</sup>R<sup>10</sup> where R<sup>9</sup> is hydrogen or alkyl, and each R<sup>10</sup> is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, aryl, heteroaryl, heterocyclic, and
- (1)  $-ONR^9[C(O)O]_zR^{10}$  where z is zero or one,  $R^9$  and  $R^{10}$  are as defined above;

X can also be -CR<sup>6</sup>R<sup>6</sup>Y' where each R<sup>6</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic and Y' is selected from the group consisting of hydroxyl, amino, thiol, alkoxy, substituted alkoxy, thioalkoxy, substituted thioalkoxy, -OC(O)R<sup>7</sup>, -SSR<sup>7</sup>, -SSC(O)R<sup>7</sup> where R<sup>7</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic,

X' is hydrogen, hydroxy, or fluoro;

X'' is hydrogen, hydroxy or fluoro, or X' and X'' together form an oxo group,

Z is selected from the group consisting of a bond covalently linking  $R^1$  to -CX'X''-, oxygen and sulfur;

n is an integer equal to 1 or 2; and pharmaceutically acceptable salts thereof with the provisos that:

A. when  $R^1$  is phenyl or 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH;

B. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is  $-CH(OH)CH_3$  derived from D-threonine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OH or  $-C(O)OCH_3$ ;

C. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^4$  is benzyl,  $R^5$  is hydrogen, X is methoxycarbonyl, X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^3$  is not methyl;

10

15

20

25

D. when  $R^1$  is iso-propyl,  $R^2$  is  $-CH_2C(O)NH_2$ ,  $R^3$  is hydrogen,  $R^4$  is iso-butyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not  $-C(O)OCH_3$ ;

E. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^3$ , the nitrogen atom attached to  $R^3$ , and  $R^4$  do not form 1,2,3,4-tetrahydroiso-quinolin-2-yl or pyrrolidin-2-yl;

F. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-C(O)OCH_3$ , X' and X'' are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not 4-amino-n-butyl;

G. when  $R^1$  is 3-nitrophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is -CH(OH)CH<sub>3</sub>,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH<sub>2</sub> or -CH<sub>2</sub>OH;

H. when  $R^1$  is phenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^5$  is hydrogen, X is  $-CH_2OCH_3$ , X' and X" are hydrogen, Z is a bond, and n is 1, then  $R^4$  is not benzyl or ethyl;

I. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is methyl,  $R^4$  is methyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$ ;

J. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -CHOH $\phi$  or -CH<sub>2</sub>OH;

K. when  $R_1$  is N-(2-pyrrolidinonyl),  $R_2$  is methyl,  $R_3$  is hydrogen,  $R_4$  is benzyl,  $R^5$  is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -C(O)OCH<sub>3</sub>;

L. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl derived from D-alanine,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NH-benzyl;

M. when R<sup>1</sup> is 3,5-difluorophenyl, R<sup>2</sup> is methyl, R<sup>3</sup> is hydrogen, R<sup>4</sup> is hydrogen, R<sup>5</sup> is hydrogen, X' and X" are hydrogen, Z is a bond, and n is 1, then X is not -CH<sub>2</sub>OH;

15

20

- N. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is 4-phenylphenyl,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NHC(CH<sub>3</sub>)<sub>3</sub>; and
- O. when  $R^1$  is 3,5-difluorophenyl,  $R^2$  is methyl,  $R^3$  is hydrogen,  $R^4$  is phenyl derived from D-phenylglycine,  $R^5$  is hydrogen, X' and X'' are hydrogen, Z is a bond, and n is 1, then X is not -C(O)NHCH(CH<sub>3</sub>) $\phi$ .
- 62. The compound according to Claim 61 wherein  $R^1$  is an unsubstituted aryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 63. The compound according to Claim 62 wherein the unsubstituted R<sup>1</sup> aryl group is selected from the group consisting of phenyl, 1-naphthyl and 2-naphthyl.
  - 64. The compound according to Claim 61 wherein  $R^1$  is a substituted aryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
  - 65. The compound according to Claim 64 wherein said substituted aryl group is a mono-substituted, di-substituted or tri-substituted phenyl group.
    - 66. The compound according to Claim 65 wherein the substituted phenyl groups are selected from the group consisting of 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 3-methoxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2-hydroxyphenyl, 2-methylphenyl, 2-fluorophenyl, 2-chlorophenyl, 3,4-difluorophenyl, 2,3,4,5,6-pentafluorophenyl, 3,4-dibromophenyl, 3,4-dichlorophenyl, 3,4-methylene-dioxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 2,4-dichlorophenyl, and 2,5-difluorophenyl.

10

- 67. The compound according to Claim 61 wherein  $R^1$  is an alkaryl group and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 68. The compound according to Claim 67 wherein the R<sup>1</sup> alkaryl group is selected from the group consisting of benzyl, 2-phenylethyl, and 3-phenyl-n-propyl.
- 69. The compound according to Claim 61 wherein  $R^1$  is selected from the group consisting of alkyl, alkenyl, cycloalkyl and cycloalkenyl groups and Z is a bond covalently linking  $R^1$  to -CX'X''-.
  - 70. The compound according to Claim 69 wherein R<sup>1</sup> is alkyl.
- 71. The compound according to Claim 69 wherein R<sup>1</sup> is cycloalkyl.
  - 72. The compound according to Claim 69 wherein R1 is alkenyl.
  - 73. The compound according to Claim 69 wherein R<sup>1</sup> is cycloalkenyl.
- 74. The compound according to Claim 69 wherein the R¹ alkyl, cycloalkyl, alkenyl and cycloalkenyl groups are selected from the group consisting of *iso*-propyl, *n*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tent*-butyl, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, aminomethyl, and N-*tent*-butoxycarbonylaminomethyl.
  - 75. The compound according to Claim 61 wherein  $R^1$  is selected from the group consisting of heteroaryl and substituted heteroaryl groups and Z is a bond covalently linking  $R^1$  to -CX'X''-.

15

- 76. The compound according to Claim 75 wherein the R<sup>1</sup> heteroaryl and substituted heteroaryl groups are selected from the group consisting of pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, fluoropyridyls (including 5-fluoropyrid-3-yl), chloropyridyls (including 5-chloropyrid-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-(thiophenyl)thiophen-5-yl, 6-methoxythionaphthen-2-yl, 3-phenyl-1,2,4-thiooxadiazol-5-yl and 2-phenyloxazol-4-yl.
- 77. The compound according to Claim 61 wherein R<sup>2</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl and heterocyclic.
  - 78. The compound according to Claim 77 wherein  $R^2$  is selected from the group consisting of methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, phenyl, 4-fluorophenyl, 3,5-difluoro-phenyl, 4-methoxyphenyl, benzyl, cyclopropyl, cyclohexyl, cyclopentyl, cycloheptyl, thien-2-yl, thien-3-yl,  $-CH_2CH_2SCH_3$ ,  $-CH_2OCH_2\phi$ ,  $-CH(CH_3)OCH_2\phi$ ,  $-CH(OH)CH_3$  and  $-CH_2OH$ .
    - 79. The compound according to Claim 61 wherein X' and X'' are hydrogen and Z is a bond covalently linking  $R^1$  to -CX'X''-.
- 80. The compound according to Claim 79 wherein R³ is selected from the group consisting of hydrogen, methyl or together with R⁴ and the nitrogen to which R³ is attached forms pyrrolidin-2-yl, 2,3-dihydroindol-2-yl, piperidin-2-yl, 4-hydroxy-pyrrolidin-2-yl and 1,2,3,4-tetrahydroisoquinolin-3-yl.
- 81. The compound according to Claim 61 wherein R<sup>4</sup> substituents are selected from the group consisting of hydrogen, methyl, ethyl, iso-propyl,

n-propyl, n-butyl, sec-butyl, iso-butyl, cyclopentyl, cyclohexyl, allyl, iso-but-2enyl, 3-methylpentyl, -CH2-cyclopropyl, -CH2-cyclohexyl, -CH2-indol-3-yl, phenyl, p-(phenyl)phenyl, m-(phenyl)phenyl, o-fluorophenyl, m-fluorophenyl, p-fluorophenyl, p-bromophenyl, m-methoxyphenyl, p-methoxyphenyl, phenethyl, benzyl, m-hydroxybenzyl, p-hydroxybenzyl, p-nitrobenzyl, 5 m-trifluoromethylphenyl, p-(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, p-(CH<sub>3</sub>)<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, p-phenylphenyl, 3,5-difluorophenyl, p-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyrid-6-yl, 4-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)benzyl, -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>, -CH<sub>2</sub>-imidazol-4-yl, -CH<sub>2</sub>-(3-tetrahydrofuranyl), -10 CH2-thien-2-yl, -CH2-thiazol-4-yl, -CH2(1-methyl)cyclopropyl, -CH2-thien-3-yl, thien-3-yl, thien-2-yl, -CH<sub>2</sub>-C(O)O-t-butyl, -CH<sub>2</sub>-C(CH<sub>3</sub>)<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, 2-methylcyclopentyl, -cyclohex-2-enyl, -CH[CH(CH $_3$ ) $_2$ ]COOCH $_3$ , -(CH $_2$ ) $_2$ SCH $_3$ , -CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=CHCH<sub>3</sub> (cis and trans), -CH<sub>2</sub>OH, -CH(OH)CH<sub>3</sub>, -CH(O-t-butyl)CH<sub>3</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>NH-Boc, 15 - $(CH_2)_4NH_2$ , - $(CH_2)_4N(CH_3)_2$ , - $CH_2$ -pyridyl, pyridyl, - $CH_2$ -naphthyl, - $CH_2$ - $(N-CH_2)_4NH_2$ , - $(CH_2)_4NH_2$ morpholino), p-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5-chlorobenzo[b]thiophen-2-yl, 4,5,6,7tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, tetrazol-5-yl, 5chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl, -CH<sub>2</sub>-N-phthalimidyl, 2-methylthiazol-4-yl, and thieno[2,3-b]thiophen-2-yl, 20 5-bromothien-2-yl, 4-bromothien-2-yl, 5-chlorothien-2-yl, 3-phenoxyphenyl, 2-phenoxyphenyl, 4-ethylphenyl, 2-benzylphenyl, (4-ethylphenyl)phenyl, 4-tertbutylphenyl, 4-n-butylphenyl, o-(4-chlorophenoxy)phenyl, furan-2-yl, and 4phenylacetylenylphenyl.

25 82. The compound according to Claim 61 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"- and R<sup>4</sup> and R<sup>5</sup> are fused to form a cycloalkyl group selected from the group consisting of cyclopropyl and cyclobutyl.

- 83. The compound according to Claim 61 wherein Z is a covalent bond linking  $R^1$  to -CX'X''-, X is -C(O)Y and Y is selected from the group consisting of hydroxy, alkoxy or substituted alkoxy.
- 84. The compound according to Claim 83 wherein Y is alkoxy or substituted alkoxy selected from the group consisting of methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, tent-butoxy, neo-pentoxy, benzyloxy, 2-phenylethoxy, 3-phenyl-n-propoxy, 3-iodo-n-propoxy, 4-bromo-n-butoxy, -ONHC(O)OC(CH<sub>3</sub>)<sub>3</sub>, -ONHC(CH<sub>3</sub>)<sub>3</sub> and hydroxy.
- 85. The compound according to Claim 61 wherein Z is a covalent bond linking R<sup>1</sup> to -CX'X"-, X is -C(O)Y and Y is -NR'R".
  - 86. The compound according to Claim 85 wherein Y is selected from the group consisting of amino (-NH<sub>2</sub>), -NH(iso-butyl), -NH(sec-butyl), N-methylamino, N,N-dimethylamino, N-benzylamino, N-morpholino, azetidino, N-thiomorpholino, N-piperidinyl,
- N-hexamethyleneimino, N-heptamethylene-imino, N-pyrrolidinyl, -NH-methallyl, -NHCH<sub>2</sub>-(furan-2-yl), -NHCH<sub>2</sub>-cyclopropyl, -NH(*tert*-butyl), -NH(*p*-methylphenyl), -NHOCH<sub>3</sub>, -NHCH<sub>2</sub>(*p*-fluorophenyl), -NHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, -NH-cyclopentyl, -NH-cyclohexyl, -NHCH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -NHCH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>, -NHCH<sub>2</sub>-(pyrid-2-yl), -NHCH<sub>2</sub>-(pyrid-3-yl), -NHCH<sub>2</sub>-(pyrid-4-yl), N-thiazolindinyl,
- 20 -N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -N[CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -NHOH, -NH(p-NO<sub>2</sub>- $\phi$ ), -NHCH<sub>2</sub>(p-NO<sub>2</sub>- $\phi$ ), -NHCH<sub>2</sub>(m-NO<sub>2</sub>- $\phi$ ), -N(CH<sub>3</sub>)OCH<sub>3</sub>, -N(CH<sub>3</sub>)CH<sub>2</sub>- $\phi$ , -NHCH<sub>2</sub>-(3,5-difluorophenyl), -NHCH<sub>2</sub>CH<sub>2</sub>F, -NHCH<sub>2</sub>(p-CH<sub>3</sub>O- $\phi$ ), -NHCH<sub>2</sub>(m-CH<sub>3</sub>O- $\phi$ ), -NHCH<sub>2</sub>(p-CF<sub>3</sub>- $\phi$ ), -N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>2</sub> $\phi$ , -NHCH(CH<sub>3</sub>) $\phi$ , -NHCH<sub>2</sub>-(p-F- $\phi$ ), -N(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -NHCH<sub>2</sub>-(tetrahydrofuran-2-yl),
- -NHCH<sub>2</sub>(p-trifluoromethylphenyl), -NHCH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -NH-[(p-benzyl)pyrid-4-yl], -NH-[(2,6-dimethyl)pyrid-4-yl], -NH-(2-methylcyclohexyl), -NH-(4-methylcyclohexyl), -NH-[N-ethoxycarbonyl]-piperidin-4-yl, -NHOC(CH<sub>3</sub>)<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-φ, -C(O)NH(CH<sub>2</sub>)<sub>3</sub>O-(p-CH<sub>3</sub>)φ,

- -C(O)NH(CH<sub>2</sub>)<sub>6</sub>NH<sub>2</sub>, -NH-(tetrahydrofuran-2-yl), -N(CH<sub>3</sub>) $\phi$ , -NH(CH<sub>2</sub>)<sub>4</sub>NHC(O)-(2-hydroxy-4-azido)-phenyl and -NH(CH<sub>2</sub>)<sub>6</sub>-(biotinamidyl).
- 87. The compound according to Claim 61 wherein X is -C(O)Y and Y is selected from the group consisting of -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OH,

  -CH(OH)CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(OH)φ, -CH(OH)CH<sub>2</sub>C(O)OCH<sub>3</sub>,

  -C(OH)(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>OC(O)OCH<sub>3</sub>, and -CH<sub>2</sub>OC(O)C(CH<sub>3</sub>)<sub>3</sub>,

  methyl, ethyl, iso-propyl, n-propyl, iso-butyl, n-butyl, sec-butyl, tent-butyl, 
  CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-pyridy-2-yl, -CH<sub>2</sub>-pyridy-3-yl,

  -CH<sub>2</sub>-pyridy-4-yl, -CH<sub>2</sub>-fur-2-yl, benzyl, cyclopentyl, phenyl, and

  -NH-SO<sub>2</sub>-CH<sub>3</sub>.
  - 88. The compound according to Claim 61 wherein Z is a covalent bond linking  $R^1$  to -CX'X''-.
  - 89. The compound according to Claim 61 wherein the compound of formula I is selected from the group consisting of:
- N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-histidine methyl ester

N-benzyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide

- 20 N-2-(N,N-dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide
  - N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide
- N-2-(N,N-dimethylamino)ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide
  - N-(4-pyridyl) methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

N-(3-pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide N-(4-pyridyl)methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminohexanamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanoate tert-5 butyl ester N-[N-(pent-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(dec-4-enoyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[3-(N,N-10 dimethylamino)propoxy]phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-[(tertbutyloxycarbonyl)methoxy]phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tyrosine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-15 (carboxymethoxy)phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2morpholinoethoxy)phenylalanine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-6-(N,N-1)dimethylamino)hexanoate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-ami20 pyridyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(3-b)-2-amino-3-(3pyridyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-proline methyl ester  $1\hbox{-}[\emph{N}\hbox{-}(3,5\hbox{-}difluor ophenylacetyl})\hbox{-}L\hbox{-}alaninyl] piperidine-2\hbox{-}carboxylate$ 25 methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(4pyridyl)propionate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-30 methoxypropionate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-morpholinopropionate methyl ester

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(2-morpholinoethoxy)phenylalaninamide

5 N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-methoxypropionamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine methyl ester

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(4-pyridyl)propionamide

10 N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(2-pyridyl)propionamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4-yl)propionate methyl ester

2-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1,2,3,4tetrahydroisoquinoline-3-carboxylate methyl ester

N-(3-methoxybenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(1-naphthyl)propionate methyl ester

20 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-naphthyl)propionate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(2-thienyl)propionate methyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine benzyl ester

25 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-bromopropyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine 3-iodopropyl ester

N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine tert-butyl ester

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2pyridyl)acetamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3pyridyl)acetamide  $N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-N_{\epsilon}-(tert-butoxycarbonyl)-L-alaninyl]$ 5 lysine methyl ester methyl N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-4phenylbutanoate N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 2-phenylethyl ester 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine 3-phenylpropyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4pyridyl)acetamide N-[N-(phenylacetyl)-L-alaninyl]-L-threonine methyl ester N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide 15 N'-[N-(phenylacetyl)-L-alaninyl]-L-alaninamide N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide N'-[N-(phenylacetyl)-L-alaninyl)-L-valinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-pyridyl)acetateethyl ester 20 N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N,N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide N,N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-leucinamide N, N-dimethyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-phenylalaninamide N-methyl-N'-[N-(phenylacetyl)-L-alaninyl]-L-valinamide 25 N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminohexanamide

N, N-dimethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2aminohexanamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohexanamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-5 methoxyphenyl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4methoxyphenyl)acetate methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-pyridyl)acetate ethyl ester 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-pyridyl)acetate ethyl ester N-[N-(cyclohexylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(cyclohex-1-enylacetyl)-L-alaninyl]-L-phenylalanine methyl ester 15 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-1-aminocyclopropane-1carboxylate methyl ester N-2-(N,N-dimethylamino) ethyl-N-methyl-N'-[N-(3,5difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine methyl ester 20 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycine benzyl ester N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-alanine ethyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]glycine ethyl ester 25 N-hydroxy-N'-[N-(3-nitrophenylacetyl)-L-alaninyl]-D,L-threoninamide N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester

alaninamide

N-[N-(3-nitrophenylacetyl)-L-alaninyl]-2-amino-3-(3hydroxyphenyl)propionate methyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-tyrosine ethyl ester N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester N-[N-[N-(isovaleryl)-L-valinyl]-L-phenylglycinyl]-L-alanine iso-butyl 5 ester N-[N-(isovaleryl)-L-phenylalaninyl]-L-alanine iso-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alanine ethyl ester 1-[N-(3-nitrophenylacetyl)-L-alaninyl]-indoline-(S)-2-carboxylate ethyl 10 ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide  $N\hbox{-methoxy-}N\hbox{-methyl-}N'\hbox{-}[N\hbox{-}(isovaleryl)\hbox{-}L\hbox{-phenylglycinyl}]\hbox{-}L\hbox{-alanina mide}$ N-iso-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N, N-di-n-propyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-15 alaninamide N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valinamide N-(4-nitrophenyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester 20 N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninamide N-iso-butyl-N'-[N-(isovaleryl)-L-phenylglycinyl]-L-alaninamide N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylalaninamide

N-(4-nitrobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-

	N-(4-nitrophenyl)- $N'$ -[ $N$ -(isovaleryl)- $L$ -phenylglycinyl]- $L$ -alaninyl]- $L$ -alaninyl]- $L$ -
	N-(4-nitrophenyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
5	N-benzyl- $N$ -methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N-(3,5-difluorobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
10	N-(3-nitrobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N-benzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N-(4-nitrobenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylalaninamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophan methyl ester
15	N-(4-methoxybenzyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N-[N-(phenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
	N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalaninyl]-L-phenylglycine methyl ester
20	N-[N-(cyclohexylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine methyl ester
	N-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]-L-phenylglycine methyl ester
25	N-(2-phenylethyl)- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -alaninamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-tryptophanamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-cyclohexylpropionate methyl ester

N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2amino-3-(4-nitrophenyl)propionamide N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-serine ethyl ester  $N-[(R)-\alpha-methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-$ 5 alaninamide  $N-[(S)-\alpha-methylbenzyl]-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L$ alaninamide N-(4-fluorobenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 10 N-(4-pyridylmethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-(4-trifluoromethylbenzyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-phenylpropionate15 ethyl ester N-[N-(3,5-diffuorophenylacetyl)-L-alaninyl]-L-phenylalanine tert-butylester N-[N-(3,5-diffuor ophenylacetyl)-L-alaninyl]-2-amino-2-methylpropionatemethyl ester 20 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-cyclohexylacetate ethyl ester N-(2-methoxyethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(isovaleryl)-2-amino-2-cyclohexylacetyl]-L-alanine ethyl ester 25 N-2-(N,N-dimethylamino) ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenylacetyl)alaninyl]-L-phenylglycinamide N-(2-pyridylmethyl)-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(3-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester N-[N-(2-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester30

	N-[N-(4-pyridylacetyl)-L-alaninyl]-L-phenylalanine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4-fluorophenyl)acetate ethyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-fluorophenyl)acetate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-alanine ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-phthalimidopropionate ethyl ester
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine neopentyl ester
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
15	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valinyl]morpholine
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine ethyl ester
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threonine methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester
	4-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-amino-3-tert-butoxybutyryl]morpholine
	4-[N-(N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucinyl]morpholine
	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine methyl ester
25	N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-isoleucine
	N-[N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-threoninyl]-L-valine ethylester

N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-leucine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-leucine methyl ester N-2-methoxyethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-5 alaninamide N-2-(N,N-dimethylamino) ethyl-N'-[N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenylacetyl)alaninyl]-L-alaninamide N-cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide 10 N-neopentyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide N-tetrahydrofurfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide 3-[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninyl]thiazolidine 15 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methylester N-[N-(3-nitrophenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester N-(R)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 alaninamide  $1\hbox{-}[N\hbox{-}(3,5\hbox{-}difluor ophenylacetyl)\hbox{-}L\hbox{-}alaninyl]\hbox{-}L\hbox{-}alaninyl]pyrrolidine$ N-(S)-sec-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lalaninamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-valine methyl ester N-2-fluoroethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-25 alaninamide N-[(S)-6-methyl-3-oxohept-2-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide

	N-4-nitrobenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminobutyramide
	N-4-nitrobenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-( $S$ )-2-aminopentanamide
5	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-fluorophenyl)acetate methyl ester
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetamide
10	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(5-chlorobenzothiophen-2-yl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-2yl)acetate ethyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-3 yl)acetate methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-thienyl)acetate methyl ester
·	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(benzothiophen-5 yl)acetate ethyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetate <i>tert</i> -butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-thienyl)acetic acid
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(1H-tetrazol-5-yl)acetate methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(6-methoxy-2-naphthyl)acetate methyl ester
30	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(3-trifluoromethylphenyl)acetate methyl ester

- N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(4,5,6,7-tetrahydrobenzothiophen-2-yl)acetate methyl ester
- N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(thieno[2,3-b]thiophen-2-yl)acetate methyl ester
- 5 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(2-methylthiazol-4-yl)acetate methyl ester
  - (3S,4S)-N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-4-amino-3-hydroxy-5-phenylpentanoate methyl ester
- N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-aminohex-4-enoate methyl ester
  - N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
  - N-tent-butyl-N'-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(4-phenylphenyl)acetamide
- N-[N-(3,5-difluorophenylacetyl)-(S)-2-aminobutanoyl]-L-phenylglycine tert-Butyl Ester
  - N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine tert-butyl ester
  - N-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycine methyl ester
- 20 N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycine methyl ester
  - N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycine methyl ester
  - N-[N-(3,5-difluorophenylacetyl)-L-leucinyl]-L-phenylglycine methyl ester
- N-[N-(3,5-difluorophenylacetyl)-L-phenylalaninyl]-L-phenylglycine methyl ester
  - N-[N-(3,5-difluorophenylacetyl)]-L-phenylglycine methyl ester
  - N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine methyl ester
  - N-[N-(phenylacetyl)-L-alaninyl]-L-alanine methyl ester

N-[N-(phenylacetyl)-L-alaninyl]-L-leucine methyl ester N-[N-(phenylacetyl)-L-alaninyl]-L-isoleucine methyl ester N-[N-(phenylacetyl)-L-alaninyl]-L-proline methyl ester N-[N-(phenylacetyl)-L-alaninyl]-L-phenylalanine methyl ester  $N-[N-(phenylacetyl)-L-alaninyl]-N_{\epsilon}-(tent-butoxycarbonyl)-L-lysine methyl$ 5 ester N-[N-(phenylacetyl)-L-alaninyl]-glycine methyl ester N-[N-(phenylacetyl)-L-alaninyl]-L-valine methyl ester N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminobutanoate methyl ester N-[N-(phenylacetyl)-L-alaninyl]-(S)-2-aminopentanoate methyl ester 10 N-[N-(3-nitrophenylacetyl)-L-alaninyl]-L-valine N-[N-(phenylacetyl)-L-alaninyl]-L-N-methylalanine methyl ester N-[N-(isovaleryl)-L-phenylglycinyl]-L-alanine iso-butyl ester N-[N-(isovaleryl)-L-isoleucinyl]-L-alanine iso-butyl ester 15 N-Cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-hydroxyproline ethyl ester 20 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-lysine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-glutamide  $1\hbox{-}[\emph{N-}(3,5\hbox{-}difluor ophenylacetyl)\hbox{-}L\hbox{-}alaninyl] piperidine\hbox{-}2\hbox{-}carboxylate}$ methyl ester N-[(S)-3-hydroxy-6-methylhept-2-yl]-N'-(3,5-difluorophenylacetyl)-L-25 alaninamide N-[(S)-2-hydroxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide

 $N-[N-(3,5-difluorophenyl-\alpha-fluoroacetyl)-L-alaniny]-L-phenylglycine$ tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-2-(S)-aminocyclohexylacetyl]-Lphenylglycine methyl ester N-[(1R,2S)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-1-hydroxy5 L-alaninamide N-[(1R,2S)-1-hydroxy-1,2-diphenyleth-2-yl]-N'-(3,5difluorophenylacetyl)-L-alaninamide N-[(1S,2R)-I-hydroxy-1-phenylprop-2-yl]-N'-(3,5-difluorophenylacetyl)-10 L-alaninamide N-2-methoxyethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]glycinamide N-[(S)- $\alpha$ -hydroxy- $\alpha$ -phenyl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[(S)-2-hydroxy-1,2-diphenylethyl]-N'-(3,5-difluorophenylacetyl)-L-15 alaninamide N-[(S)-1-hydroxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide $N-[\alpha-hydroxy-\alpha'-(4-hydroxyphenyl)-iso-propyl]-N'-(3,5$ difluorophenylacetyl)-L-alaninamide N-2-pyridylmethyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-20 phenylalaninamide  $N-[\alpha-hydroxy-\alpha'-pyrid-2-yl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L$ alaninamide  $N-[\alpha-hydroxy-\alpha'-pyrid-4-yl-iso-propyl]-N'-(3,5-difluorophenylacetyl)-L-$ 25 alaninamide N-[(S)-1-hydroxy-4-methylpent-2-yl]-N'-(3,5-difluorophenylacetyl)-L-methylpent-2-yl]-N'-(3,5-difluorophenylacetyl)-N'-(3,5-difluoalaninamide N-[ $\alpha$ -methoxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide N-[1-hydroxy-3-methyl-but-2-yl]-N'-(3,5-difluorophenylacetyl)-L-30 alaninamide

	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-2-(6-aminopyrid-2 yl)acetate methyl ester
	N-[1-hydroxy-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
5	N-[(S)-2-methoxy-1-phenyleth-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(S)-1-methoxy-2-phenyl-prop-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
	N-[(S)-1-acetoxyhex-2-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
10	N-[(S)-1-( $tert$ -butylcarbonyloxy)-hex-2-yl]-N'-(3,5-difluorophenylacetyl) L-alaninamide
	N-[2-hydroxy-1-(thien-2-yl)ethyl]- $N'$ -(3,5-difluorophenylacetyl)- $L$ -alaninamide
·	N-[(S)-2-hydroxy-2-methyl-1-phenylprop-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide
15	N-[N-(3,5-difluorophenylacetyl)-L-(thien-2-yl)glycinyl]-L-phenylalanine tert-butyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinol
	N-[N-(cyclopropaneacetyl)-L-phenylglycinyl]-L-phenylglycinol
	N-[N-(cyclopentaneacetyl)-L-phenylglycinyl]-L-phenylglycinol
20	N-[N-(3,5-difluorophenylacetyl)-D,L-phenylglycinyl]-D,L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-D,L-valinyl]-D,L-phenylglycinamide
	N-[N-(2-thienylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-[N-(n-caprotyl)-L-alaninyl]-L-phenylglycinamide
25	N-[N-(3,5-difluorophenylacetyl)-L-norleucinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-norvalinyl]-L-phenylglycine methyl

	N-[N-(3,5-difluorophenylacetyl)-L-tert-leucinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-isoleucinyl]-L-phenylglycine methyl ester
5	N-[N-(3,5-difluorophenylacetyl)-L-cyclohexylalaninyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(cyclopropyl)acetyl]-L-phenylglycine methyl ester
10	N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-3-yl)acetyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-(S)-2-amino-2-(thien-2-yl)acetyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-(4-fluorophenyl)glycinyl]-L-phenylglycine methyl ester
15	N-[N-(3,5-difluorophenylacetyl)-D-(4-fluorophenyl)glycinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-(4-methoxyphenyl)glycinyl]-L-phenylglycine methyl ester
20	N-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester
	N-[N-(cyclopropylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester
	N-[N-(cyclopentylacetyl)-L-phenylglycinyl]-L-phenylglycine tert-butyl ester
25	N-[N-(tert-butylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -(5-bromothien-2-yl)glycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ -(5-bromothien-2-yl)glycinamide
30	N-tert-butyl- $N'$ - $[N$ - $(3,5$ -difluorophenylacetyl)- $L$ -alaninyl]- $L$ - $(4$ -bromothien- $2$ -yl)glycinamide

	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -(thien-2-yl)glycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide
5	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-(thien-3-yl)glycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-(thien-2-yl)glycinamide
10	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -(5-chlorothien-2-yl)glycinamide
15	N-Cyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ -4- $($ phenyl)phenylglycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenoxy)phenylglycinamide
20	$N$ -(S)-(-)- $\alpha$ -methylbenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-3-(phenyl)phenylglycinamide
	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(ethyl)phenylglycinamide
25	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-(phenyl)phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -2-(benzyl)phenylglycinamide
30	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -4-bromophenylglycinamide

	N-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(cyclohexyl)phenylglycinamide
:	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(4-ethylphenyl)phenylglycinamide
5	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -4-(tert-butyl)phenylglycinamide
	N-tert-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-3-(4-chlorophenoxy)phenylglycinamide
10	N-cyclohexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-(phenyl)phenylglycinamide
	$N$ -[ $N$ -(3,5-difluorophenyl- $\alpha$ -hydroxyacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester
	<i>N-tert</i> -butyl- <i>N'</i> -[ <i>N</i> -(3,5-difluorophenyl- $\alpha$ , $\alpha$ -difluoroacetyl)-L-alaninyl]-L-phenylglycinamide
15	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine <i>tert</i> -butyl ester
	N-[(S)-1-oxo-1-phenylprop-2-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
20	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(pyrid-3-yl)glycine tert-butyl ester
	[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]morpholine
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-(2-methoxy)phenylglycine methyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tert-butoxycarbonyl(hydroxyl amine) ester
	N-neopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
30	N-tetrahydrofurfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide

	N-methoxy- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
·	[N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinyl]azetidine
5	N-iso-butyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-cyclopropanemethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
10	N-methoxy- $N$ -methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
	N-2-methylprop-2-enyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L phenylglycinamide
	N-(pyrid-3-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
15	N-(pyrid-4-yl)methyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-furfuryl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
20	N-cyclopentyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-1-benzylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N,N-dimethyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D,L$ -phenylglycinamide
 25	N-2,2,6,6-tetramethylpiperidin-4-yl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinamide
	N-2-methylcyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide
30	N-4-methylcyclohexyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $D$ , $L$ -phenylglycinamide

N-1-ethoxycarbonylpiperidin-4-yl-N'-[N-(3,5-difluorophenylacetyl)-Lalaninyl]-D,L-phenylglycinamide N-methyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide 5 N-tert-butoxy-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycine N-tertbutyl(hydroxylamine) ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycine hydrazide 10 N-(1-ethoxyethen-1-yl)-[N'-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycine hydrazide N-[N-(phenylacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester N-4-(phenyl)butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide 15 N-3-(4-iodophenoxy)propyl-N'-[N-(3.5-difluorophenylacetyl)-L-alaninyll-L-phenylglycinamide N-6-(amino)hexyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,Lphenylglycinamide Hydrochloride N-1-(phthalimido)pent-2-yl-N'-(3,5-difluorophenylacetyl)-L-alaninamide 20 N-[N-(3,5-difluorophenylacetyl)-L-(3,5-difluorophenyl)glycinyl]-L-(3,5difluorophenyl)glycine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-norleucine N-[N-(cyclopentaneacetyl)-L-alaninyl]-L-phenylglycine tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine iso-25 propyl ester N-(isopropyl) N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-Lphenylglycinamide N-[N-(cyclopentylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester N-[N-(cyclopropylacetyl)-L-alaninyl]-L-phenylalanine tert-butyl ester

N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-phenylglycine iso-butyl ester N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-D-phenylglycine methyl $N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-(3-\alpha-phenyl)$ proline 5 methyl ester N-[N-(3,5-Difluorophenylacetyl)-L-alaninyl]-L-azetidine methyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-2-amino-3-(5chlorobenzothiophen-2-yl)acetate methyl ester 10 N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-3-(thiazol-4yl)propionate tert-butyl ester N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide tertbutyl ester  $N\hbox{-}[N\hbox{-}(3,5\hbox{-}difluor ophenylacetyl)\hbox{-}L\hbox{-}alaninyl]\hbox{-}D\hbox{-}(thien-2\hbox{-}yl)glycinamide}$ 15 N-[N-(3,4-dichlorophenylacetyl)-L-alaninyl]-D-phenylglycinamide  $N\hbox{-}[N\hbox{-}(3\hbox{-}chlorophenylacetyl)\hbox{-}L\hbox{-}alaninyl]\hbox{-}D\hbox{-}phenylglycinamide}$ N-[N-(3-bromophenylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(3-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide20 N-[N-(4-fluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(3-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(4-methylphenylacetyl)-L-alaninyl]-D-phenylglycinamideN-[N-(3-trifluoromethylphenylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(3-methoxyphenylacetyl)-L-alaninyl]-D-phenylglycinamide  $N\hbox{-}[N\hbox{-}(2\hbox{-chlorophenylacetyl})\hbox{-}L\hbox{-}alaninyl]\hbox{-}D\hbox{-}phenylglycinamide}$ 25 N-[N-(1-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide N-[N-(2-naphthylacetyl)-L-alaninyl]-D-phenylglycinamide

	N-[N-(phenylacetyl)-L-alaninyl]-D-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycine
·	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D-phenylglycinamide
<b>5</b> ,	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-(S)-2-amino-2-(2-furanyl)acetamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide
	N'-[ $N$ -(3,4-difluorophenylacetyl)-D-alaninyl]-D-phenylglycinamide
	N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylalanin-N-methylsulfonamide
10	N''-methyl- $N''$ -phenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]-glycinamide
	N''-methyl- $N''$ -phenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-alaninamide
	N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-phenylglycinamide
15	N''-methyl- $N''$ -benzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-glycinamide
	N''-4-fluorobenzyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
20	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-(4-fluoro)phenylglycine neopentyl ester
	N-[N-(2,3,4,5,6-pentafluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(pyrid-3-yl)glycine <i>tert</i> -butyl ester
25	N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)serinyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-(O-benzyl)threoninyl]-L-phenylglycine methyl ester

	N-[N-(3,5-difluorophenylacetyl)-L-threoninyl]-L-phenylglycine methyl ester
	N-[N-(3,5-difluorophenylacetyl)-L-serinyl]-L-phenylglycine methyl ester
5	N''-4-methylphenyl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	N''-tetrahydrofurfuryl- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)- $L$ -alaninyl]- $L$ -phenylglycinamide
	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenyl-glycinamide
10	N'-[ $N$ -(3,5-difluorophenylacetyl)-L-methionyl]-L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-2-aminobutanoyl]-L-phenylglycinamide
	N'-[N-(3,5-difluorophenylacetyl)-L-phenylglycinyl]-L-phenylglycinamide
	N-[N-(3,5-difluorophenylacetyl)-L-valinyl]-L-phenylglycinamide
15	$N$ -[(R)- $\alpha$ -methylbenzyl]- $N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide
	N-[1-phenyl-2-oxo-3-methylbutan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-propan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)-L-alaninamide
20	N-[1-phenyl-2-oxo-pentan-1-yl]- $N'$ -(3,5-difluorophenylacetyl)- $L$ -alaninamide
	N-[1-phenyl-2-oxo-2-phenyl-ethan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)- $L$ -alaninamide
25	N-[1-phenyl-2-oxo-butan-1-yl]-N'-(3,5-difluorophenyl-acetyl)-L-alaninamide
	N-[1-phenyl-2-oxo-4-methylpentan-1-yl]- $N'$ -(3,5-difluorophenyl-acetyl)-L-alaninamide
	$N'$ -[ $N$ -(3,5-difluorophenylacetyl)-L-alaninyl]-L- $\alpha$ -hydroxyphenylalanine methyl ester

N''-[4-((2-hydroxy-4-azido)-phenyl)-NHC(O)-)butyl] N'-[N-(3,5difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide N-[(S)-1-phenyl-2-oxo-2-phenyl-ethan-1-yl]-N'-(3,5-difluorophenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyl-phenyl-2-oxo-2-phenyacetyl)-L-alaninamide 5 N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-fluorophenylglycine tert-butyl ester N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-phenylphenylglycine tert-butyl ester [N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-(2,3-benzo[b]proline) methyl 10 ester N''-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-4-nbutylphenylglycinamide N''-tert-butyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-4-(phenylacetenyl)phenylglycinamide 15 N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-phenylglycinthioamide N-[1,3-diphenyl-2-oxo-propan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[1-phenyl-2-oxo-2-cyclopentylethan-1-yl]-N'-(3,5difluorophenylacetyl)-L-alaninamide 20 N-[1-phenyl-2-oxo-hexan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N-[1-phenyl-2-oxo-3-methylpentan-1-yl]-N'-(3,5-difluorophenylacetyl)-Lalaninamide N''-n-hexyl-6-biotinamidyl-N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-25 D,L-phenylglycinthioamide N'-[N-(3,5-difluorophenylacetyl)-L-methioninyl]-L-methionine N'-[N-(2-tert-BOC-amino)propionyl)-L-alaninyl]-L-phenylglycine methyl ester N"-tert-butyl N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-L-2-30 fluorophenylglycinamide

N'-[N-(3,5-difluorophenylacetyl)-L-alaninyl]-D,L-2-phenylglycine methyl ester

N-[(S)-1-phenyl-2-oxo-3-phenylpropan-1-yl]-N'-(3,5-difluorophenylacetyl)-L-alaninamide

5 N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-phenylglycine

N'-[N-(3,5-difluorophenylacetyl)-D,L-thien-3-ylglycinyl]-D,L-2-phenylglycine tert-butyl ester

N'-[N-(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine

N'-[N-(3,5-difluorophenylacetyl)-L-thien-3-ylglycinyl]-L-2-phenylglycine tent-butyl ester

 $N\mbox{-}[2\mbox{-hydroxy-1-(S)phenyleth-1-yl}]-N'\mbox{-}[(3,5\mbox{-difluorophenylacetyl})-L-phenylglycinyl]-L-alaninamide$ 

N-[2-hydroxyeth-1-yl]-N'-[(3,5-difluorophenylacetyl)-L-alaninyl]-L-phenylglycinamide

N'-[N-(3,5-difluorophenyl-2-oxo-acetyl)-L-alaninyl]-L-2-phenylglycine tert-butyl ester

[N-(2,5-dichlorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3,5-difluorophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

20 [N-(3,4-dichlorothiophenoxyacetyl)-L-alaninyl]-L-phenylglycine methyl ester

[N-(3-aminoproprionyl)-L-alaninyl]-L-phenylglycine tert-butyl ester; and

[N-(3-tert-butoxycarbonylamino)propionyl)-L-alaninyl]-L-phenylglycine tert-butyl ester.

								÷
					v		;	
					1.75			
P.								
1 <sub>.,</sub>				4				
	and the second second			and the second	· *%, 3	<b>t</b> ia di	A Ministra	*
٠.								4
•				**************************************			The state of the s	
				Company of the compan		·		36
								,
•	Andrew Town							
				3.				
						8 <b>4</b>		
∯.º A					*.			
	•							
				•				
ė .								
	•							
					,			
						•		
, •			•					
} •	·				*.			
P.								
	A second second	•						
•								
			•	·	•			



## **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup>: C07K 5/06, C07C 237/22, C07D 213/38, 333/20, A61K 38/04, 31/44, 31/38

A3

(11) International Publication Number:

WO 98/22494

(43) International Publication Date:

28 May 1998 (28.05.98)

(21) International Application Number:

PCT/US97/20804

(22) International Filing Date:

21 November 1997 (21.11.97)

(30) Priority Data:

08/755,442	22 November 1996 (22.11.96)	US
08/808,528	28 February 1997 (28.02.97)	US
08/807,528	28 February 1997 (28.02.97)	US
08/807,427	28 February 1997 (28.02.97)	US

(71) Applicants (for all designated States except US): ATHENA NEUROSCIENCES, INC. [US/US]; 800 Gateway Boulevard, South San Francisco, CA 94080 (US). ELI LILLY & COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): AUDIA, James, E. [US/US]; 6449 Lakeside Woods Circle, Indianapolis, IN 46278 (US). BRITTON, Thomas, C. [US/US]; 4700 Royal Oak Lane, Carmel, IN 46033 (US). DROSTE, James, J. [US/US]; 7430 Wood Stream Drive, Indianapolis, IN 46254 (US). FOLMER, Beverly, K. [US/US]; 26 Currant Drive, Newark, DE 19702 (US). HUFFMAN, George, W. [US/US]; 12 Maple Crest Drive, Carmel, IN 46032 (US). JOHN, Varghese [US/US]; 1722 18th Avenue, San

Francisco, CA 94122 (US). LATIMER, Lee, H. [US/US]; 56 Sheridan Road, Oakland, CA 94618 (US). MABRY, Thomas, E. [US/US]; 8104 Hi-Vu Drive, Indianapolis, IN 46227 (US). NISSEN, Jeffrey, S. [US/US]; 4348 Oil Creek Drive, Indianapolis, IN 46268 (US). PORTER, Warren, J. [US/US]; 8037 Lieber Road, Indianapolis, IN 46260 (US). REEL, Jon, K. [US/US]; 14701 Alsong Court, Carmel, IN 46032 (US). THORSETT, Eugene, D. [US/US]; 571 Buena Vista Street, Moss Beach, CA 94038 (US). TUNG, Jay, S. [US/US]; 2224 Semeria Avenue, Belmont, CA 94002 (US). WU, Jing [CN/US]; Apartment 502, 10 De Sabla Road, San Mateo, CA 94402 (US). EID, Clark, Norman [US/US]; 53 Dundee Drive, Cheshire, CT 06410 (US). SCOTT, William, Leonard [US/US]; 144 Buckingham Drive, Indianapolis, IN 46208 (US).

- (74) Agents: SWISS, Gerald, F. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).
- (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report:

26 November 1998 (26.11.98)

# (54) Title: METHODS AND COMPOUNDS FOR INHIBITING β-AMYLOID PEPTIDE RELEASE AND/OR SYNTHESIS

#### (57) Abstract

Disclosed are compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. Also disclosed pharmaceutical compositions comprising a compound which inhibits  $\beta$ -amyloid peptide release and/or its synthesis as well as methods for treating Alzheimer's disease both prophylactically and therapeutically with such pharmaceutical compositions.

#### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL A	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM A	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT A	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU A	Australia	GΛ	Gabon	LV	Latvia	SZ	Swaziland
AZ A	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA E	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB E	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE E	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF E	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG E	Bulgaria	HU	Hungary .	ML	Mali	TT	Trinidad and Tobago
BJ E	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
BR E	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY E	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA C	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF C	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG C	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH S	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI C	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM C	Cameroon		Republic of Korea	PL	Poland		
CN C	China	KR	Republic of Korea	PT	Portugal Portugal		
CU C	Cuba	KZ	Kazakstan	RO	Romania		
CZ C	Czech Republic	LC	Saint Lucia	หบ	Russian Federation		
DE C	Jermany	LI	Licchtenstein	SD	Sudan		
DK I	Denmark	LK	Sri Lanka	SE	Sweden		
RE E	Estonia	LR	Liberia	SG	Singapore		

ernational Application No PCT/US 97/20804

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K5/06 C07C237/22 C07D213/38 C07D333/20 A61K38/04 A61K31/44 A61K31/38 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 6 CO7K CO7C CO7D A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. X WO 96 20949 A (SALK INST BIOTECH IND 1-89 ; MCDONALD IAN ALEXANDER (US); ALBRECHT ELISA) 11 July 1996 see the whole document X WO 96 20725 A (SALK INST BIOTECH IND 1-24; MUNOZ BENITO (US); MCDONALD IAN ALEXANDER 30-53, () 11 July 1996 59-82, 88,89 see the whole document -/--Χŀ Further documents are listed in the continuation of box C. Patent family members are tisted in annex. Special categories of cited documents : "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) involve an inventive step when the document is taken alone Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docu-"O" document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled in the art. "P" document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 14, 10, 1998 29 September 1998 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016 Groenendijk, M

ernational Application No PCT/US 97/20804

	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	Tourse and the state of the sta
Category '	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 09838 A (MERRELL DOW PHARMA; SCOIS NOVA INC (US); CORDELL BARBARA (US); SCH) 13 April 1995  see claims 1,4,6,7,11-14	1-12,16, 17, 19-23, 30, 32-41, 45,46, 48-52, 59, 61-70, 74,75, 77-81,88
x	PAPADOPOULOS A ET AL: "ANODIC OXIDATION OF N-ACYL AND N-ALKOXYCARBONYL DIPEPTIDE ESTERS ASA KEY STEP FOR THE FORMATION OF CHIRAL HETEROCYCLIC SYNTHETIC BUILDING BLOCKS" TETRAHEDRON, vol. 47, no. 4/05, 28 January 1991, pages 563-572, XP000565601 See table on page 567, compound f	61-64, 77-81, 83,84, 88,89
A	EP 0 652 009 A (LILLY CO ELI ;ATHENA NEUROSCIENCES INC (US)) 10 May 1995 see the whole document	
A	WO 95 13084 A (MILES INC ;TAMBURINI PAUL PERRY (US); BENZ GUENTER H H H (US); HAE) 18 May 1995 see the whole document	
A	EP 0 732 399 A (CEPHALON INC) 18 September 1996 see the whole document	
Ρ,Χ	WO 96 39194 A (ATHENA NEUROSCIENCES INC) 12 December 1996	1-23, 25-52, 54-81, 83-89
	see the whole document	

International application No. PCT/US 97/20804

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)						
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:						
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:						
Although claims 1-31 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.						
2. X Claims Nos.: 1-30,32-59,61-88(all partially) because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:						
see FURTHER INFORMATION sheet PCT/ISA/210						
Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).						
B x II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)						
This International Searching Authority found multiple inventions in this international application, as follows:						
see additional sheet						
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.						
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.						
As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:						
No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:						
Remark on Protest  The additional search fees were accompanied by the applicant's protest.  X No protest accompanied the payment of additional search fees.						

#### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Claims Nos.: 1-30,32-59,61-88(all partially)

The scope of the claims 1-30, 32-59 and 61-88 is very broad and speculative. A formula consisting virtually of variables which are moreover in at least part of the claims ill-defined (e.g. the use of "(hetero)aryl" and "heterocyclicl") and from which formula many embodiments are disclaimed cannot be considered to be a clear and concise definition of patentable subject-matter (Art.6 PCT). Furthermore the available experimental data actually only comprise a very small part of the compounds claimed, which part is moreover not evenly distributed over the whole claimed area. Therefore the claims can also not be considered to represent a permissible generalisation which is fairly based on experimental evidence, that is, they are also not adequately supported by the description (Art.6 PCT). Therefore a meaningful and economically feasible search could not encompass the complete subject-matter of the claims. Consequently the search had been limited to the actually synthesised examples and (closely) related analogs, that is the compounds encompassed by the claim 89 and the related claims 31 and 60 and extended to compounds with the same activity (Art.17(2)(a)(ii) PCT, PCT Guidelines III,2.1)).

### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-3,6-8,19-32,35-37,48-61,64-66, 77-89(all partially)

Compounds of Formula I, wherein R1 is an phenyl group having at least a 3,5-difluoro substitution and Z is a bond as defined in claim 89, their pharmaceutical compositions and use  $\frac{1}{2}$ 

2. Claims: 1-3,11,12,14,16,19-23,25-32,40,41,43,45,48-52, 54-61,69,70,72,74,77-81,83-89(all partially)

Compounds of Formula I, wherein R1 is a (amine substituted) alkyl or alkenyl group and Z is a bond as defined in claim 89, their pharmaceutical compositions and use

3. Claims: 1-5,19-23,25-34,48-52,54-63,77-81, 83-89(all partially)

Compounds of Formula I, wherein R1 is an unsubstituted phenyl or naphtyl group and Z is a bond as defined in claim 89, their pharmaceutical compositions and use

4. Claims: 1-3,13,15,16,19-23,25-32,42,44,45,48-52,54-61,71, 73,74,77-81,83-89(all partially)

Compounds of Formula I, wherein R1 is a cycloalkyl group and Z is a bond as defined in claim 89, their pharmaceutical compositions and use

5. Claims: 1-3,6-8,19-23,25-32,35-37,48-52,54-61,64-66,77-81, 83-89(allpartially)

Compounds of Formula I, wherein R1 is a substituted phenyl group and Z is a bond or oxygen as defined in claim 89, their pharmaceutical compositions and use as far as not covered by subject 1

6. Claims: 1-3,17-23,25-32,46-52,54-61,75-81, 83-89(all partially)

Compounds of Formula I, wherein R1 is a heterocyclic group

International Application No. PCT/US 97 /20804

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210
and Z is a bond or oxygen as defined in claim 89, their pharmaceutical compositions and use
·

Information on patent family members

ternational Application No PCT/US 97/20804

Patent document cited in search report		. Publication date	Patent family member(s)		Publication date	
WO 9620949	A	11-07-1996	US	5804560 A	08-09-1998	
			ĂÜ	4697296 A	24-07-1996	
			ΑÜ	4697396 A	24-07-1996	
			CA	2209234 A	11-07-1996	
•			EP	08 <b>00</b> 528 A	15-10-1997	
			WO	9620725 A	11-07-1996	
			US	5714471 A	03-02-1998	
WO 9620725	Α	11-07-1996	US	5804560 A	08-09-1998	
			AU	4697296 A	24-07-1996	
	,		AU	4697396 A	24-07-1996	
			CA	2209234 A	11-07-1996	
			EP	0800528 A	15-10-1997	
			WO	9620949 A	11-07-1996	
			US	5714471 A	03-02-1998	
WO 9509838	Α	13-04-1995	AU	687449 B	26-02-1998	
			ΑU	7799894 A	01-05-1995	
			CA	2171882 A	13-04-1995	
			CN	1132504 A	02-10-1996	
			EP	0721449 A	17-07-1996	
			FI	961438 A	29-03-1996	
			HU	74004 A	28-10-1996	
			JP	9505281 T	27-05-1997	
		•	NO	961326 A	01-04-1996	
			ZA	9407529 A	29-05-1995	
EP 0652009	Α	10-05-1995	AU	6897094 A	16-02-1995	
			CA	2129689 A	10-02-1995	
			CN	1120040 A	10-04-1996	
			CZ	9401841 A	15-03-1995	
-			JP	7165606 A	27-06-1995	
			NO	942883 A	10-02-1995	
			NZ	264143 A	26-11-1996	
			ZA	9405719 A	01-02-1996	
WO 9513084	Α	18-05-1995	AU	679675 B	10-07-1997	
			AU	3711093 A	18-11-1993	
			AU	5726494 A	04-07-1994	
			CA	2095888 A	12-11-1993	

information on patent family members

cernational Application No PCT/US 97/20804

Patent document cited in search report	Publication date		nt family nber(s)	Publication date	
WO 9513084 A		CA 2151927 A		23-06-1994	
			0569777 A	18-11-1993	
			0694076 A	31-01-1996	
		JP 8	3268911 A	15-10-1996	
		JP 8	3508464 T	10-09-1996	
		MX 9	9302725 A	31-08-1994	
		NZ	247575 A	25-09-1996	
•		WO S	9413319 A	23-06-1994	
		NZ	272218 A	27-07-1997	
	•	ZA S	9303249 A	11-03-1994	
		AU 7	7115094 A	29-05-1995	
			2175564 A	18-05-1995	
		EP (	0744954 A	04-12-1996	
EP 0732399 A	18-09-1996	AU	661270 B	20-07-1995	
		AU 7	7465491 A	10-10-1991	
		CA 2	2077665 A	06-09-1991	
	•	EP (	0518955 A	23-12-1992	
		FI	923983 A	04-09-1992	
		WO 9	9113904 A	19-09-1991	
WO 9639194 A	12-12-1996	US 5	783434 A	21-07-1998	
			221684 A	12-12-1996	
			831920 A	01-04-1998	